

The Effect of the Repeated Subcultures of *Helicobacter pylori* on Adhesion, Motility, Cytotoxicity, and Gastric Inflammation

In vitro subcultures of bacteria can lead to genetic and phenotypic changes. This study aimed at investigating the effect of repeated subcultures on the adhesion, motility, cytotoxicity, and gastric inflammation caused by *Helicobacter pylori*. *H. pylori* SS1 strain was subcultured 64 times on agar plates containing Brucella broth and 5% bovine calf serum. The adhesion, motility, cytotoxicity, and gastric inflammation produced in Mongolian gerbils were compared between the first and 64th subcultured strain. The adhesion rates, following 3 hr exposure of AGS cells to either the first strain or the 64th-transferred strain, were 21% and 12%, respectively. The motility of the 64th-transferred strain decreased significantly when compared to the 1st strain (9.1 mm vs. 15.1 mm). The cytotoxicity index tended to be higher in the first strain than in the 64th-transferred strain (73.7% vs. 69.2%). The initial infection rate on the gerbils showed no difference between the two strains. However, chronic gastric inflammation of the first strain-infected gerbils was somewhat more severe than that of the 64th-transferred strain-infected gerbils. Therefore, the use of repeatedly subcultured strains of *H. pylori* in virulence experiments can lead to different results from those of the original strain.

Key Words : *Helicobacter pylori*; Growth; Adhesions; Motility; Gastritis

Sung Soo Kim, Hak Sung Lee,
Young Seok Cho, Youn Soo Lee*,
Choon Sang Bhang, Hiun Suk Chae,
Sok Won Han, In Sik Chung, Doo Ho Park

Division of Gastroenterology, Department of Internal
Medicine, Department of Clinical Pathology*,
College of Medicine, Catholic University of Korea,
Seoul, Korea

Received : 7 February 2002

Accepted : 7 March 2002

Address for correspondence

In Sik Chung, Professor
Division of Gastroenterology
Department of Internal Medicine
Kangnam St. Mary's Hospital, 505 Banpo-dong,
Seocho-gu, Seoul 137-040, Korea
Tel : +82.2-590-1402, Fax : +82.2-590-1409
E-mail : isc@cmc.cuk.ac.kr

*We received financial support for this research from
the research fund of Catholic University Medical
College.

†This article was selected for poster presentation at
the XIVth International Workshop on Gastroduode-
nal Pathology and *Helicobacter pylori*, Strasbourg.

INTRODUCTION

Virtually everyone infected with *Helicobacter pylori* develops life-long chronic gastritis, but only a minority of infected individuals will develop a clinically relevant disease, such as peptic ulcer or gastric carcinoma. The mechanisms responsible for the diversity of clinical outcomes of *H. pylori* infection are not well established. Both host genetic factors and specific properties of the bacterial strains are suggested to play an important role in the development of *H. pylori*-related diseases. Putative virulence factors of *H. pylori* may include colonization factors, such as urease, adhesion, and motility, and such disease-associated factors as cytotoxicity (1-3).

A multitude of in vitro and animal studies investigating the virulence factors of *H. pylori* have been performed. Defined laboratory strains, or culture collection strains of *H. pylori* have frequently been used in such experiments. However, it is evident that such laboratory strains may have quite different properties from those of corresponding organisms under in vivo condition. An example of this was the repeated subcultures of *Streptococcus mutans*, which resulted in the loss of its adhesive qualities and changes in expression of surface proteins (4-7).

In vitro cultivation of *H. pylori* isolates on solid media may result in loss of the antigenic O side-chain (8). Also, specific mutants of the original organisms can be made by adjusting the culture condition (9, 10). Therefore, the virulence factors of laboratory strains, following repeated subcultures, may be quite different from those of the original *H. pylori* strain.

To investigate the effect of sequential subculture on the virulence factors of *H. pylori*, we have conducted in vitro experiments on a defined form of *H. pylori* and its progeny after repeated subcultures, investigating their adhesion, motility, and cytotoxicity, and the gastric inflammation they cause in the Mongolian gerbil.

MATERIALS AND METHODS

Preparation of bacteria and subculture

H. pylori SS1 strain was obtained from A. Lee (School of Microbiology and Immunology, The University of New South Wales, Sydney, Australia) in 1998. The organism was subcultured in twice sequentially and admixed with brain

heart infusion broth containing 20% (v/v) glycerol, then stored at -70°C for about 30 months. The stored organisms were thawed, and grown on Brucella agar (Difco Inc., Detroit, Mich., U.S.A.) containing 5% (v/v) bovine calf serum which was incubated at 37°C , 10% (v/v) CO_2 , 95% humidity for 3 days. These organisms were used as the first ancestor strain (first strain) in this experiment. Then, the first strain was sequentially subcultured 64 times under the same culture conditions. The first strain and 64th-transferred strain ('64th strain') used




Fig. 1. Differences in adhesion rates after exposure to the first strain and 64th-transferred strain of *H. pylori*. The adhesion rate upon AGS cells is significantly lower in the 64th-transferred strain than in the first strain.

in these experiments were incubated on the liquid media containing Brucella broth and 5% (v/v) bovine calf serum in a shaking incubator at 10% CO_2 (v/v) and 95% humidity at 37°C for 30 hr prior to use. All in vitro experiments were performed in triplicate.

Adhesion rate

The adhesion of *H. pylori* to the AGS cell line was quantified by measuring bacterial urease activity as previously described (11). Briefly, 1×10^5 AGS cells per well were cultured in 96-well tissue culture plates with medium containing 1×10^9 organisms per well of *H. pylori*. After 3 hr, the plate was washed with phosphate-buffered saline (PBS) to remove the *H. pylori* that did not adhere. Saline containing 2% (w/v) urea and 0.03% (w/v) phenol red was then added and the bacterial urease activity was assayed by measuring the optical density at 540 nm. Cells treated with culture medium without *H. pylori* served as negative controls. The index was calculated at percentage of optical density of co-cultured *H. pylori* per that of *H. pylori* alone.

Motility

Motility was determined in a semisolid medium that consisted of Brucella broth, 5% (v/v) bovine calf serum, and 0.4% (w/v) agarose (Difco, Detroit, MI, U.S.A.) as previously described (2). The melted medium was poured into petri dishes (60 mm in diameter) and solidified. The agar plates were dried for 30 min on a clean bench, and then a paper disk (6 mm in diameter) was placed in the center of the agar plate.




Fig. 2. Motility of *H. pylori* in semisolid medium. The first strain (A) spread widely in the medium. The 64th-transferred strain (B) show a narrow growth zone around a paper disk.




Fig. 3. Histologic examination of Mongolian gerbil after inoculation with either the first or 64th-transferred strain. (A), Numerous curved bacilli are visualized on the gastric pit (two weeks of infection with the first strain, Warthin-Starry silver stain, $\times 400$). The infiltration of chronic inflammatory cells on the mucosa and submucosa is more severe in the first strain-infected gerbil (B) than that in the 64th-transferred strain-infected gerbil (C) (12 weeks of infection, H&E, $\times 200$).

The culture broth of *H. pylori* ($10 \mu\text{L}$; 1×10^7 organisms) was inoculated onto the paper disk and the agar plate was incubated at 37°C , 10% (v/v) CO_2 , 95% humidity, for five days. The diameter of the growth zone was measured using a slide caliper.

Vacuolating cytotoxic activity

Cytotoxicity was quantified by measuring the cellular release of lactose dehydrogenase (LDH) into the culture medium as previously described (12). Briefly, 1×10^5 AGS cells per well were cultured in 96-well tissue culture plates with medium containing 1×10^9 organisms per well of *H. pylori*. After 9 hr, the plate was centrifuged for 10 min at $250 g$, and $100 \mu\text{L}$

of supernatant from the well was transferred into a new microplate. The concentration of LDH was then measured using a cytotoxicity kit from TaKaRa Biomedicals (Shiga, Japan). AGS cells alone in culture medium served as negative controls.

Gastric inflammation

Five-week-old male Mongolian gerbils were fasted for 24 hr and then challenged orally with 0.4 mL of culture broth containing *H. pylori* (1×10^9 CFU/mL) using a $\phi 0.9 \times 70$ mm sized Zonde needle (Dae Jong Instrument Co., Seoul, Korea), and this was repeated twice during five days. Two weeks and 12 weeks after inoculation, animals were killed and the stomach was dissected. Half of the stomach was used




Fig. 4. Differences in gastric inflammation in a gerbil infected for 12 weeks. Although there is no statistical difference, the chronic inflammatory cell infiltration tends to be higher in the first strain than in the 64th strain.

for urease testing and the culture of *H. pylori*, and the remaining stomach was fixed in 10% (v/v) neutral buffered formalin for histological examination. *H. pylori* infection was confirmed by positive culture, or when both the urease test and histology were positive. Gastric inflammation, such as infiltration of PMN neutrophils and monocytes, was assessed according to the updated Sydney system (13).

RESULTS

Adhesion rate

The repeated subculture was proven to decrease the cellular adhesion of *H. pylori*. The adhesion rate of the first strain to AGS cells was significantly higher than that of the 64th strain. The adhesion indices following 3 hr exposure of the first strain and 64th strain to the AGS cells were 21% and 12%, respectively (Fig. 1).

Motility

The change of motility was also significant following repeated subculturing of *H. pylori*. The motility of the 64th strain decreased markedly compared to that of the first strain. The diameter of the growth zone of the first strain was 15.1 ± 0.9 mm, whereas that of the 64th strain decreased to 9.1 ± 0.6 mm (Fig. 2).

Cytotoxicity

Exposure of AGS cells to either the first or 64th strains for

9 hr significantly increased LDH release into the culture medium. Exposure of AGS cells to the first strain increased LDH release somewhat more than exposure to the 64th strain. The increment of LDH in the first strain was $73.7 \pm 2.7\%$, and that in the 64th strain was $69.2 \pm 2.3\%$.

Gastric inflammation

Infection rates at two weeks after inoculation was not different between the first strain-infected gerbils and the 64th strain-infected gerbils. In both groups, 8 of the 10 inoculated gerbils were confirmed to be infected (80%). However, gastric inflammation, which was assessed at 12 weeks after infection, was somewhat different between the first strain-infected gerbils and the 64th strain-infected gerbils. The chronic inflammation in the first strain-infected gerbils tended to be more severe than that in 64th strain-infected gerbils. The mean score of monocyte infiltration in the first strain-infected gerbils was 2.0 ± 0.78 , and that in the 64th strain-infected gerbils was 1.4 ± 0.54 (Fig. 3, 4).

DISCUSSION

Although host genetic and environmental factors may have an important role in *H. pylori*-induced diseases, bacterial virulence factors such as urease, motility, adhesins, or cytotoxins are responsible for the colonization and inflammation of *H. pylori* (1, 3, 14).

These results demonstrated that repeated subculturing of *H. pylori* could change the biological activity of the virulence factors. The adhesion rate, motility, and cytotoxicity of the original strains decreased following 64 successive propagations in vitro. Gastric inflammation produced by the subcultured strain was less severe than that of the original strain. Therefore, various in vitro and in vivo virulence experiments using the old and several-times-transferred strains of *H. pylori* demonstrated different results from those of the original strains.

Our results may undoubtedly be expected when considering previous studies. In many reports, subcultures of the bacteria and cell lines can lead to genetic and phenotypic changes. For example, *Streptococcus mutans* freshly isolated from saliva showed decreased ability to adhere following a number of in vitro subcultures (4, 5). A variant strain, Xc100L, which was isolated following 100 times subcultures of *Streptococcus mutans*, strain Xc, showed a decrease in the production of a surface protein antigen with a molecular mass of 190 kDa (PAC). This was associated with changes in transcriptional regulation of the gene (6, 7).

For experimental purposes, specific mutants of the original organisms can be made by nutritional limitations and atmospheric changes during growth. A TEM-1 derived variant beta-lactamase with increased resistance to beta-lactamase inhibitors was obtained by repeated subcultures of *E. coli* J62-

2 for five days in the presence of subinhibitory concentrations of amoxicillin plus clavulanic acid (9). In vitro, spontaneous mutants resistant to fluoroquinolones are frequent in methicillin-resistant *Staphylococcus aureus* when selected on isosensitest-agar containing 1 µg/mL of ciprofloxacin (10).

In addition, the plating of successive *Staphylococcus aureus* subcultures of daily transfers proved that genetic mutants are responsible for antigen variation (15). Changes in the genetic expression of several proteins following subculture of primary rat astrocytes were also demonstrated (16).

Fresh clinical isolates of *H. pylori* expressed high-molecular weight lipopolysaccharide (LPS) with an O side-chain (S-LPS). Following numerous in vitro passages on solid media, low-molecular-weight LPS (R-LPS) was produced (8). However, this change could be reversed when grown in liquid media. Culture collection strains that were subcultured many times also produced R-LPS when grown on solid media, but when the same strains were grown under liquid conditions, S-LPS was produced. They presumed these changes in phenotypic expression were caused by a difference in the growth condition of solid media compared with in vivo conditions. However, the changes of virulent properties in our experiments were not reversed when the organisms were grown in liquid media before use.

We conducted in vivo experiments to determine whether the changes in the virulence properties, following repeated subculture, affect the inflammation of the gerbil gastric mucosa. Although the initial infection rate was not different, the chronic inflammation of the stomach tended to be different between the first strain and the 64th strain.

In summary, subcultures of *H. pylori* leads to changes in the biological activity of virulence factors, such as adhesion, motility, and cytotoxicity, and that these changes can affect the degree of gastric inflammation in animal models. Therefore, the use of repeatedly subcultured strains in virulence experiments can lead to different results from those of the original strains of *H. pylori*. Many in vivo and in vitro experiments have been performed or are still ongoing to explore the disease-specific virulence factors of *H. pylori*. Therefore, it is important to keep the culture conditions for *H. pylori* as similar as possible with the intragastric environment.

REFERENCES

- Graham DY, Yamaoka Y. Disease-specific *Helicobacter pylori* virulence factors: the unfulfilled promise. *Helicobacter* 2000; 5(Suppl. 1): S3-9.
- Iwao E, Hirayama F, Takagi S, Yokoyama Y, Ikeda Y. Virulence factors of *Helicobacter pylori* affecting its gastric colonization in Mongolian gerbils. *J Gastroenterol* 1999; 34(Suppl. 11): 47-54.
- Yoshiyama H, Nakazawa T. Unique mechanism of *Helicobacter pylori* for colonizing the gastric mucus. *Microbes Infect* 2000; 2: 55-60.
- Orstavik J, Orstavik D. Influence of in vitro propagation on the adhesive qualities of *Streptococcus mutans* isolated from saliva. *Acta Odontol Scand* 1982; 40: 57-63.
- Westergren G, Olsson J. Hydrophobicity and adherence of oral streptococci after repeated subculture in vitro. *Infect Immun* 1983; 40: 432-5.
- Koga T, Asakawa H, Okahashi N, Takahashi I. Effect of subculturing on expression of a cell-surface protein antigen by *Streptococcus mutans*. *J Gen Microbiol* 1989; 135: 3199-207.
- Yamashita Y, Tsukioka Y, Nakano Y, Shibata Y, Koga T. Molecular and genetic analysis of multiple changes in the levels of production of virulence factors in a subcultured variant of *Streptococcus mutans*. *FEMS Microbiol Lett* 1996; 144: 81-7.
- Walsh EJ, Moran AP. Influence of medium composition on the growth and antigen expression of *Helicobacter pylori*. *J Appl Microbiol* 1997; 83: 67-75.
- Thomson CJ, Amyes SG. Selection of variants of the TEM-1 beta-lactamase, encoded by a plasmid of clinical origin, with increased resistance to beta-lactamase inhibitors. *J Antimicrob Chemother* 1993; 31: 655-64.
- Witte W, Grimm H. Occurrence of quinolone resistance in *Staphylococcus aureus* from nosocomial infection. *Epidemiol Infect* 1992; 109: 413-21.
- Rosberg K, Hubinette R, Nygard G, Berglindh T, Rolfsen W. Studies of *Helicobacter pylori* in a gastric mucosa in vitro animal model. *Scand J Gastroenterol* 1991; 26: 43-8.
- Smoot DT, Wynn Z, Elliott TB, Allen CR, Mekasha G, Naab T, Ashktorab H. Effects of *Helicobacter pylori* on proliferation of gastric epithelial cells in vitro. *Am J Gastroenterol* 1999; 94: 1508-11.
- Dixon MF, Genta RM, Yardley JH, Correa P. Classification and grading of gastritis. The updated Sydney System. International Workshop on the Histopathology of Gastritis, Houston 1994. *Am J Surg Pathol* 1996; 20: 1161-81.
- Gschwanter M, Dragosics B. Physiopathology of *Helicobacter pylori* infections. *Acta Med Austriaca* 2000; 27: 117-21.
- Pereira AT. Further studies on antigen variation in *Staphylococcus aureus*. *J Clin Microbiol* 1981; 13: 245-7.
- Passaquin AC, Schreier WA, de Vellis J. Gene expression in astrocytes is affected by subculture. *Int J Dev Neurosci* 1994; 12: 363-72.