

Incidence and Risk Factors of Subsequent Hip Fractures in Korea: Multicenter Study

Kee Haeng Lee,¹ Ju Young Kim,²
Soo Jae Yim,³ Do Hyun Moon,⁴
Geun Hong Choi,⁵ and Kyoung Ho Moon⁵

¹Department of Orthopedic Surgery, Bucheon St. Mary's Hospital, The Catholic University of Korea, Bucheon; ²Department of Orthopedic Surgery, Incheon St. Mary's Hospital, The Catholic University of Korea, Incheon; ³Department of Orthopedic Surgery, Soonchunhyang University Bucheon Hospital, Bucheon; ⁴Department of Orthopedic Surgery, Gil Medical Center, College of Medicine, Gachon University, Incheon; ⁵Department of Orthopaedic Surgery, College of Medicine, Inha University, Incheon, Korea

Received: 29 October 2013

Accepted: 30 April 2014

Address for Correspondence:

Kyoung Ho Moon, MD

Department of Orthopaedic Surgery, College of Medicine, Inha University, 27 Inhang-ro, Jung-gu, Incheon 400-711, Korea
Tel: +82.32-890-3663, Fax: +82.32-890-3467
E-mail: moon@inha.ac.kr

Funding: This study was supported by a grant of Inha University.

This study analyzes the incidence of subsequent hip fractures and its risk factors in the northwestern region of Korea. We analyzed hip fracture patients who visited any of the 5 teaching hospitals in the Bucheon and Incheon area from January 2000 to December 2010. Medical records were reviewed and presence of subsequent hip fractures, alcohol history, marital status, live in solitude, dementia, dizziness, American society of anesthesiologists score, osteoporosis treatment after fracture, body mass index (BMI) and initial bone mineral density were analyzed. The average follow-up period was 12 months (range 1-130 months). A total of 2,546 patients (women 1,770, men 776) who had experienced hip fractures were included. Of these, subsequent hip fractures were found in 233 patients (9.2%) (women 187, men 46). Mean age at the time of the first fracture was 79.2 yr old (range 50-100 yr). The average interval between the first fracture and the subsequent hip fractures was 30.2 months (range 4 days-154 months). In this large-scale, retrospective, multicenter study, overall incidence of subsequent hip fractures is 9.2%. Independent risk factors of subsequent fracture are women, BMI < 22 kg/m², and being unmarried.

Keywords: Subsequent Hip Fractures; Risk Factors; Incidence

INTRODUCTION

The one-year mortality of hip fractures in elderly patients ranges from 14%-36%, and the incidence of hip fractures is increasing each year because of increasing the number of the elderly. Epidemiologic studies on hip fracture in Korea have reported high incidences of hip fracture, and confirmed that the number of hip fractures is likely to increase markedly in the near future (1, 2). It is a worldwide problem related to the aging of the population (3-7). It is expected that 63 million hip fractures will occur globally in 2050; and Melton et al. reported 6% of males and 17.5% of females will experience hip fracture (3, 4). The complications of hip fracture are death, disability, long term care needs and loss of social independency (8). Thus hip fracture appears as a significant public health problem with serious socioeconomic burden (9). Hodsman et al. (5) reported that patients with previous hip fractures will experience subsequent hip fractures (SHF) and they also suggested that the mortality is 2.7 times higher than the group without previous fractures. The incidence of asynchronous bilateral hip fractures is 1.7%-14.8%; and hip fractures alone are a major risk of contralateral fracture (10-17). So we analyzed the incidence of subsequent hip fractures and

its risk factors in the northwestern region of Korea.

All hip fracture patients (femoral head fractures, femoral neck fractures, intertrochanteric fractures, subtrochanteric fractures) treated at teaching 5 hospitals in Bucheon and Incheon area were reviewed. The patients were divided into two groups, a group that had experienced subsequent hip fractures (SHF), and a group that had unilateral hip fractures (UHF). We analyzed the incidence of SHF and the risk factors of SHF.

MATERIALS AND METHODS

We analyzed 2,546 patients (female, 1,770; male, 776). Inclusion criteria were patients who had unilateral hip fracture from January 2000 to December 2010 with no osteoporosis treatment at the time of the incident. Exclusion criteria were patients who received osteoporosis treatment at the time of the incident and causes of hip fracture were a traffic accident, a fall from a higher pointer than the patient's height, pathologic fracture and causes of follow-up loss were unknown. The average follow-up period was 12 months (1-130 months). Presence of SHF, alcohol history, marital status, dementia, dizziness, osteoporosis treatment after fracture, American society of anesthesiologists (ASA) score,

Table 1. Baseline characteristics of the patients

Parameters	No. (%) or mean \pm SD			P value
	Total (n = 2,546)	UHF (n = 2,313)	SHF (n = 233)	
Sex (female)	1,770 (69.5)	1,583 (68.4)	187 (80.3)	< 0.001
Age (yr)	79.2 \pm 10.2	78.8 \pm 10.3	83.0 \pm 8.9	0.024
Height (cm)	158.4 \pm 9.2	158.7 \pm 9.3	155.4 \pm 7.4	< 0.001
Weight (kg)	55.2 \pm 10.7	55.5 \pm 10.7	51.2 \pm 9.5	< 0.001
BMI (kg/m ²)	23.1 \pm 3.5	23.3 \pm 3.4	22.0 \pm 3.4	< 0.001
ASA score	2.3 \pm 0.5	2.3 \pm 0.5	2.3 \pm 0.5	0.671
Alcoholism	26 (1.0)	23 (1.0)	3 (1.3)	0.719
Unmarried	77 (3.0)	64 (2.8)	13 (5.6)	0.004
Live in solitude	63 (2.5)	60 (2.6)	3 (1.3)	0.506
Dementia	113 (4.4)	102 (4.4)	11 (4.7)	0.520

SD, standard deviation; ASA, american society of anesthesiologists; UHF, unilateral hip fracture; SHF, subsequent hip fracture.

body mass index (BMI) and bone mineral density (BMD) were collected through patients' medical records and telephone interviews. Patient identification (using name and birth date) was cross checked between hospitals to obtain an accurate count of incidence. We selected a SHF group with a previous history of surgery due to unilateral hip fracture and who had SHF on the contralateral side. BMD tests were performed on 1,421 patients after initial trauma with Dual-energy X-ray absorptiometry. Student's t-tests and chi-square test were performed between the two groups on their age, BMI, BMD and ASA score. We analyzed the factors known to cause SHF such as additional fractures, alcohol consumption, solitude habitation, dementia, dizziness, medical condition, and history of osteoporosis treatment by logistic regression analysis (SPSS 18.0).

Ethics statement

The study protocol was approved by the institutional review board (IRB) of Inha University Hospital (IRB Number, 13-2441), Bucheon St. Mary's Hospital (IRB Number, 1-018), Incheon St. Mary's Hospital (IRB Number, 4-002). Informed consent was waived by the boards.

RESULTS

Their average age was 79.2 (range: 50-100 yr) yr with 776 males and 1,770 females. Seventy-one (2.8%) patients were died. Average BMI was 23.1. Alcoholism was noted in 26 patients (1%). Sixty-three patients (2.5%) lived in solitude. One hundred thirteen patients (4.4%) suffered from dementia. Their ASA physical status scored 2.3 (1-4, standard deviation [SD] 0.5) on average. Two hundred thirty three patients (9.2%) experienced a SHF. Average interval between the first fracture and the subsequent hip fractures was 30.2 months (range, 4 days-154 months). There were 3 (1.3%) alcoholism cases with SHF, 3 (1.3%) lived in solitude, 11 (4.7%) had dementia. Their ASA physical status score was 2.3 (1-4, SD 0.5) on average. In the group with UHF, 60 cases

Table 2. Bone mineral density results of patients

Timing and site	No. (%) or mean \pm SD			P value
	Total (n = 1,421)	UHF (n = 1,283)	SHF (n = 138)	
First BMD after First Fx				
Neck, T score	-2.89 \pm 1.09	-2.89 \pm 1.05	-2.83 \pm 1.43	0.643
Troch, T score	-2.42 \pm 1.16	-2.42 \pm 1.13	-2.37 \pm 1.43	0.730
Hip, T score	-2.63 \pm 1.18	-2.63 \pm 1.15	-2.63 \pm 1.43	0.976
Treatment after first Fx	703 (49.5)	628 (48.9)	75 (54.3)	< 0.001

SD, standard deviation; BMD, bone mineral density; Fx, fracture; UHF, unilateral hip fracture; SHF, subsequent hip fracture.

Table 3. Multivariate analysis of risk factors for subsequent fracture

Characteristics	Exp (B)	95% CI for Exp (B)		P value
		Lower	Upper	
Sex, female (vs. male)	2.361	1.540	3.618	< 0.001
Age (y-o), \geq 83 (vs. < 83)	1.130	0.806	1.583	0.479
BMI (kg/m ²), < 22 (vs. \geq 22)	1.844	1.324	2.569	< 0.001
Marriage, no (vs. yes)	2.537	1.254	5.136	0.010

Exp, exponential function; CI, confidence interval.

(2.6%) lived in solitude, 102 cases (4.4%) had dementia, 23 (1%) alcoholism cases. Their ASA physical status score was 2.3 (1-4, SD 0.5) on average. The two groups had statistical differences in sex, age, BMI and marital status (Table 1).

T-scores at the time of initial trauma were -2.89 (SD 1.09) of femoral neck area and -2.42 (SD 1.16) of trochanteric area. A total of 703 patients (49.5%) received osteoporosis treatment after the initial fracture. A total of 138 patients with SHF had a T-score of -2.83 (SD 1.43) of femoral neck area, and -2.37 (SD 1.43) of trochanteric area, and 1,283 patients with UHF had a T-score of -2.89 (SD 1.05) of femoral neck, and -2.42 (SD 1.13) of trochanteric area. An initial T-score in the group with SHF was -2.83 (SD 1.43) while group with UHF showed a T-score of -2.89 (SD 1.05), thus showing no statistical difference ($P = 0.643$). However, more patients in the SHF group had been receiving treatment for osteoporosis ($P < 0.001$) (Table 2).

In this study, overall incidence of subsequent hip fractures was 9.2%. Multivariate analysis of risk factors found that female gender, BMI < 22 kg/m², being unmarried were statistically significant risk factors (Table 3).

DISCUSSION

The occurrence of fracture near the hip joint is increasing as our average life span increases due to medical improvement. However, mortality rates and complication rates are very high regardless of technological advances (18, 19). Boston (16) reported that in the cases of a first femoral neck fracture, death rate goes up to 13% and 30% in cases of a second fracture. Common risk factors for hip fractures are low BMI and small calcium intake in Japan and Mediterranean osteoporosis study (MEDOS).

Additionally eating fish and sleeping on Japanese mattress might exert preventive effects against hip fracture (20). The risk factors of bilateral hip fractures are dementia, neurological diseases, and Parkinson disease (10, 12, 21). Denutrition is also a risk factor (22, 23). In this study, female gender, BMI < 22 kg/m², being unmarried were found as risk factors.

Our study has some limitations. This was a multicenter study and the prescription and duration of osteoporosis medication differed between centers. Therefore the effect of medication on SHF was difficult to estimate. Second, because a number of different surgeons were included in the study, the incidence depending on operative procedures was difficult to evaluate. Third, being a retrospective study, the collection of data was irregular and BMD tests were performed on only 1,421 patients after initial trauma, not all patients. Fourth, investigation of patients' calcium profiles, hormonal status, vitamin D deficiency was not performed.

However, this is a substantially large study that included all of the teaching hospitals in the Incheon and Bucheon area. Patient identification was cross checked between hospitals leading to an accurate evaluation of the SHF patients in the community.

In this large-scale, retrospective, multicenter study, overall incidence of subsequent hip fractures is 9.2%. Independent risk factors of subsequent fracture are female gender, BMI < 22 kg/m², and being unmarried.

DISCLOSURE

The authors have no conflicts of interest to disclose.

ORCID

Do Hyun Moon <http://orcid.org/0000-0003-4806-6650>
 Ju Young Kim <http://orcid.org/0000-0003-0182-8523>
 Kee Haeng Lee <http://orcid.org/0000-0002-2137-2992>
 Soo Jae Yim <http://orcid.org/0000-0003-2107-7384>
 Kyoung Ho Moon <http://orcid.org/0000-0002-9186-0473>
 Geun Hong Choi <http://orcid.org/0000-0003-3776-1877>

REFERENCES

1. Yoon HK, Park C, Jang S, Jang S, Lee YK, Ha YC. Incidence and mortality following hip fracture in Korea. *J Korean Med Sci* 2011; 26: 1087-92.
2. Rowe SM, Song EK, Kim JS, Lee JY, Park YB, Bae BH, Hur CI. Rising incidence of hip fracture in Gwangju city and Chonnam province, Korea. *J Korean Med Sci* 2005; 20: 655-8.
3. Cooper C, Campion G, Melton LJ 3rd. Hip fractures in the elderly: a world-wide projection. *Osteoporos Int* 1992; 2: 285-9.
4. Melton LJ 3rd, Chrischilles EA, Cooper C, Lane AW, Riggs BL. Perspective: how many women have osteoporosis? *J Bone Miner Res* 1992; 7: 1005-10.
5. Hodsmann AB, Leslie WD, Tsang JF, Gamble GD. 10-year probability of recurrent fractures following wrist and other osteoporotic fractures in a large clinical cohort: an analysis from the Manitoba Bone Density Program. *Arch Intern Med* 2008; 168: 2261-7.
6. Gardner MJ, Brophy RH, Demetrakopoulos D, Koob J, Hong R, Rana A, Lin JT, Lane JM. Interventions to improve osteoporosis treatment following hip fracture: a prospective, randomized trial. *J Bone Joint Surg Am* 2005; 87: 3-7.
7. Jubay AG, De Geus-Wenceslau CM. Evaluation of osteoporosis treatment in seniors after hip fracture. *Osteoporos Int* 2002; 13: 205-10.
8. Pearse EO, Redfern DJ, Sinha M, Edge AJ. Outcome following a second hip fracture. *Injury* 2003; 34: 518-21.
9. Lee YK, Koo KH. Osteoporotic hip fracture in the elderly patients: physicians' views. *J Korean Med Sci* 2013; 28: 976-7.
10. Shabat S, Gepstein R, Mann G, Kish B, Fredman B, Nyska M. The second hip fracture: an analysis of 84 elderly patients. *J Orthop Trauma* 2003; 17: 613-7.
11. Berry SD, Samelson EJ, Hannan MT, McLean RR, Lu M, Cupples LA, Shaffer ML, Beiser AL, Kelly-Hayes M, Kiel DP. Second hip fracture in older men and women: the Framingham Study. *Arch Intern Med* 2007; 167: 1971-6.
12. Fukushima T, Sudo A, Uchida A. Bilateral hip fractures. *J Orthop Sci* 2006; 11: 435-8.
13. Chapurlat RD, Bauer DC, Nevitt M, Stone K, Cummings SR. Incidence and risk factors for a second hip fracture in elderly women: the study of osteoporotic fractures. *Osteoporos Int* 2003; 14: 130-6.
14. Lönnroos E, Kautiainen H, Karppi P, Hartikainen S, Kiviranta I, Sulkava R. Incidence of second hip fractures: a population-based study. *Osteoporos Int* 2007; 18: 1279-85.
15. Schröder HM, Petersen KK, Erlandsen M. Occurrence and incidence of the second hip fracture. *Clin Orthop Relat Res* 1993; 289: 166-9.
16. Boston DA. Bilateral fractures of the femoral neck. *Injury* 1982; 14: 207-10.
17. Dretakis KE, Dretakis EK, Papakitsou EF, Psarakis S, Steriopoulos K. Possible predisposing factors for the second hip fracture. *Calcif Tissue Int* 1998; 62: 366-9.
18. Cummings SR, Kelsey JL, Nevitt MC, O'Dowd KJ. Epidemiology of osteoporosis and osteoporotic fractures. *Epidemiol Rev* 1985; 7: 178-208.
19. Eastwood EA, Magaziner J, Wang J, Silberzweig SB, Hannan EL, Strauss E, Siu AL. Patients with hip fracture: subgroups and their outcomes. *J Am Geriatr Soc* 2002; 50: 1240-9.
20. Yoshimura N, Suzuki T, Hosoi T, Orimo H. Epidemiology of hip fracture in Japan: incidence and risk factors. *J Bone Miner Metab* 2005; 23: 78-80.
21. Chiu KY, Pun WK, Luk KD, Chow SP. Sequential fractures of both hips in elderly patients: a prospective study. *J Trauma* 1992; 32: 584-7.
22. Simon P, Gouin F, Veillard D, Laffargue P, Ehlinger M, Bel JC, Lopez R, Beaudet P, Luickx F, Molina V, et al. Femoral neck fractures in patients over 50 years old. *Rev Chir Orthop Reparatrice Appar Mot* 2008; 94: S108-32.
23. Tonetti J, Couturier P, Rémy A, Nicolas L, Merloz P, Franco A. Proximal femoral fractures in patients over 75 years: vital and functional prognosis of a cohort of 78 patients followed during 2.5 years. *Rev Chir Orthop Reparatrice Appar Mot* 1997; 83: 636-44.