

Molecular Mechanism of Reactive Oxygen Species-dependent ASK1 Activation in Innate Immunity

Shota Yamauchi, Takuya Noguchi and Hidenori Ichijo*

Laboratory of Cell Signaling, Graduate School of Pharmaceutical Sciences, Strategic Approach to Drug Discovery and Development in Pharmaceutical Sciences, Center of Excellence (COE) Program, and Core Research for Evolutional Science and Technology (CREST), Japan Science and Technology Corporation, The University of Tokyo, Tokyo 113-0033, Japan

Apoptosis signal-regulating kinase 1 (ASK1), a mitogen-activated protein kinase kinase kinase, plays pivotal roles in stress responses. In addition, ASK1 has emerged as a key regulator of immune responses elicited by pathogen-associated molecular patterns (PAMPs) and endogenous danger signals. Recent studies have demonstrated that reactive oxygen species (ROS)-dependent activation of ASK1 is required for LPS-stimulated cytokine production as well as extracellular ATP-induced apoptosis in immune cells. The mechanism of ROS-dependent regulation of ASK1 activity by thioredoxin and TRAFs has been well characterized. In this review, we focus on the molecular details of the activation of ASK1 and its involvement in innate immunity.

[Immune Network 2008;8(1):1-6]

INTRODUCTION

The mitogen-activated protein kinase (MAPK) signaling pathways consist of three classes of protein kinases: MAPK, MAPK kinase (MAP2K), and MAP2K kinase (MAP3K). MAP3K phosphorylates and thereby activates MAP2K, and activated MAP2K in turn phosphorylates and activates MAPK such as ERK, JNK and p38 (1).

Apoptosis signal-regulating kinase 1 (ASK1) was originally identified as an apoptosis-inducing MAP3K of MKK4 (SEK1)/MKK7-JNK and MKK3/MKK6-p38 MAPK signaling cascades (2). Subsequent studies have shown that ASK1 is activated by various stresses including oxidative stress, TNF α , calcium influx, and endoplasmic reticulum stress (2-6). In mammalian innate immunity, pathogen-associated molec-

ular patterns (PAMPs) are recognized by specific toll-like receptors (TLRs), culminating in activation of the ERK, JNK, and p38 MAPK pathways and the nuclear factor-kappa B (NF- κ B) pathways (7). Recently, it has been demonstrated that reactive oxygen species (ROS)-dependent activation of the ASK1-p38 pathway is required for TLR4-mediated innate immune responses (8,9).

Even in the absence of PAMPs, endogenous factors released from infected and/or damaged cells can elicit inflammatory responses. Extracellular ATP is known to serve as one such endogenous danger signal (10). ASK1 has been shown to be involved in extracellular ATP-induced macrophage apoptosis downstream of P2X₇ receptor, a plasma membrane receptor for ATP (T. Noguchi, et al, unpublished data).

Thus, ASK1 appears to be a pivotal component of both stress and immune responses. Here, we review recent studies on the mechanism of ROS-dependent ASK1 activation and its roles in host defense.

ASK1 SIGNALOSOME

Human and mouse ASK1 consist of 1374 and 1380 amino acids, respectively. ASK1 possesses a serine/threonine kinase domain in the middle portion of the molecule flanked by the N- and C-terminal coiled-coil domains (11,12). The two coiled-coil domains play critical roles in regulation of ASK1 activity (Fig. 1).

Endogenous ASK1 is constitutively oligomerized through its C-terminal coiled-coil (CCC) domain and forms a high molec-

*Corresponding Author. Tel: 81-3-5841-4859; Fax: 81-3-5841-4778; E-mail: ichijo@mol.f.u-tokyo.ac.jp

This work was supported by Grants-in-Aid for scientific research from the ministry of Education, Sciences and Culture of Japan, by CREST, Japan Science and Technology Corporation and by Center of Excellence (COE) program.

Keywords: ASK1, ROS, thioredoxin, TRAF, TLR4, P2X₇ receptor




Figure 1. (a) Domain structure of ASK1. Human and mouse ASK1 consists of 1374 and 1380 amino acids, respectively. ASK1 has a serine/threonine kinase domain in the middle portion of the molecule flanked by the N- and C-terminal coiled-coil domains (NCC and CCC, respectively). Trx binds to the N-terminal region of ASK1. (b) ROS-dependent activation of ASK1. ASK1 constitutively forms high molecular mass complex including Trx. In response to ROS, Trx is oxidized and dissociate from ASK1, leading to the recruitment of TRAFs. TRAFs promote the NCC-dependent homophilic interaction of ASK1 and thereby activate ASK1.

ular mass complex (approximately 1,500~2,000 kDa) together with its regulatory proteins and unidentified factors. The high molecular mass complex appears to act as a molecular platform regulating basal activity and stimulus-induced activation of ASK1, which we designated ASK1 signalosome (13).

THIOREDOXIN

Thioredoxin (Trx) is a ubiquitously expressed reduction/ oxidation (redox) protein that is known to act as an antioxidant and a reductant cofactor (14,15). Trx is the first identified interacting protein of ASK1 and has been established to be a signaling intermediate of the ROS-ASK1 cascade. Trx possesses two redox-active cysteine residues in its active center (-Cys-Gly-Pro-Cys-), which form intramolecular disulfide bond in response to ROS. The reduced form of Trx [Trx-(SH)₂] directly binds to the N-terminal region of ASK1 and inhibits its kinase activity. Once oxidized, Trx (Trx-S₂) dissociates from ASK1 (16).

Our group recently found that the homophilic interaction of ASK1 through the N-terminal coiled-coil domain (NCC) was necessary for ROS-induced activation of ASK1. Trx disrupts this interaction by directly binding to ASK1. Therefore, the dissociation of Trx from ASK1 is prerequisite to the activation of ASK1 in response to ROS (17,18).

TRAFs

Dissociation of Trx from ASK1 is necessary but not sufficient for the activation of ASK1. Tumor necrosis factor receptor-associated factors (TRAFs) are the major signal transducers of the TNF receptor superfamily and the interleukin (IL)-1 receptor/TLR superfamily. By mediating the signaling pathways downstream of these receptors, TRAFs regulate various physiological processes including adaptive and innate immunity, embryonic development, bone metabolism and stress response (19). Among the TRAF family members, TRAF2 and TRAF6 have been established to play crucial roles in ASK1 activation (3,13).

Following the dissociation of Trx, the ASK1 signalosome forms a higher molecular mass complex (approximately more than 3,000 kDa), at least in part, due to the recruitment of TRAF2 and TRAF6 (13,18). In contrast to Trx, TRAF2 and TRAF6 promote the NCC-dependent homophilic interaction of ASK1, which results in autophosphorylation of the threonine residue in the activation loop and thereby activation of ASK1. Accordingly, H₂O₂-induced activation of ASK1 is strongly suppressed in TRAF2- and TRAF6-deficient mouse embryonic fibroblasts (MEFs) (13,17,20).

ASK1 IN TLR4-SIGNALING

Cells involved in the innate immune system include macro-




Figure 2. ROS-dependent activation of ASK1 in TLR4- and P2X₇ receptor-signaling.

phages, dendritic cells (DCs) and neutrophils, and these cells are responsible for the detection of pathogenic microorganisms following infection. The TLR family proteins recognize the molecular signature of invading pathogens, known as PAMPs, and make an important contribution to host defense. Each PAMP such as lipids, lipoproteins, proteins and nucleic acids is sensed by distinct TLRs, culminating in activation of the ERK, JNK and p38 MAPK pathways as well as the NF- κ B pathways. In particular, TLR4 mediates the host cell responses toward the bacterial cell wall component lipopolysaccharide (LPS) (7,21).

Recent studies in mice have shown that many MAP3Ks are involved in TLR4-mediated inflammatory responses (22). For example, MEKK3 is required for the activation of JNK and p38, but not of ERK by TLR4. MEKK3-deficient MEF produces low level of IL-6 in response to LPS (23). Tpl2/Cot, another MAP3K, mediates the ERK pathway downstream of TLR4 and contributes to LPS upregulation of TNF α in splenocytes and macrophages (24).

ASK1 is also involved in mammalian innate immune responses mediated by TLR4 (Fig. 2). LPS activates ASK1 in RAW 264.7 murine macrophage cell line. In DCs and splenocytes from ASK1-deficient mice, the activation of p38 following LPS stimulation is impaired, whereas the activation of JNK and NF- κ B is not affected. LPS-induced production of proinflammatory factors including IL-6, TNF α , IL-1 β , IL-12 and NO is reduced in these cells when compared with control cells. Consistently, ASK1-deficient mice produce small amounts

of TNF α and NO after LPS injection and therefore are resistant to septic shock. These data indicate that the ASK1-p38 pathway is required for the induction of proinflammatory cytokines downstream of TLR4 in vivo (8).

Interestingly, LPS-induced activation of ASK1 is dependent on the generation of ROS. In RAW 264.7 cells, pretreatment of antioxidants abolishes the activation of ASK1 and the cytokine production in response to LPS. Furthermore, LPS stimulates the interaction of ASK1 and TRAF6 in a ROS-dependent manner. Given the importance of Trx as a redox-dependent regulator of ASK1, it is conceivable that Trx is involved in this process (8).

Although the precise molecular mechanism underlying ROS generation after LPS stimulation remains elusive, the involvement of Nox4 has been shown. Nox4 is a member of the NADPH oxidase family that catalyzes the NADPH-dependent reduction of oxygen to generate superoxide (25). In HEK293T cells, direct interaction of Nox4 with TLR4 was reported to be required for LPS-stimulated ROS generation and NF- κ B activation (26). Nox4 was also suggested to be responsible for LPS-induced ASK1 activation in U373 astrocytoma cells (27). However, the physiological relevance of these findings in immune cells remains unclear.

ASK1 IN P2X₇ RECEPTOR-SIGNALING

In recent years, it has been shown that endogenous factors released from infected and/or damaged cells can elicit in-

flammatory responses even in the absence of PAMPs. Extracellular ATP has been reported to serve as one such endogenous danger signal by acting on cell-surface ATP receptors (10).

Nucleotides including ATP are usually present at high concentrations within the cytoplasm (5~10 mM), whereas their extracellular concentrations are low (nanomolar). Acute cell injury or death are therefore thought to cause passive ATP release into extracellular milieu (10). Moreover, ATP is reported to be actively secreted via as yet unidentified pathways (28). Such extracellular ATP stimulates two classes of cell-surface receptors; the ionotropic P2X receptors and metabotropic P2Y receptors. P2X receptors and P2Y receptors are ligand-gated cation channels and G-protein-coupled receptors in the plasma membrane, respectively (29). Accumulating evidence has suggested that P2X₇ receptor, a P2X receptor expressed primarily in immune cells, plays important roles in extracellular ATP-induced inflammatory responses. For example, ATP-stimulated maturation and secretion of the proinflammatory cytokine IL-1 β have been shown to be mediated by P2X₇ receptor (30,31).

Recently, our group demonstrated that the P2X₇ receptor-ASK1-p38 pathway was necessary for ATP-induced apoptosis in macrophages (Fig. 2). In RAW 264.7 cells, ASK1 is activated in response to ATP, which is mediated by ROS derived from Nox2/gp91phox, a component of phagocytic Nox complex. ROS generation is almost completely inhibited by selective P2X₇ receptor antagonist, suggesting that P2X₇ receptor is responsible for the activation of ASK1. In spleen-derived macrophages from ASK1^{-/-} mice, ATP-induced p38 activation and consequent apoptosis are suppressed (T. Noguchi, et al, unpublished data).

Although physiological role of ATP-induced apoptosis in macrophages remains to be elucidated, it is reported that macrophages infected with certain bacteria undergo apoptosis, resulting in the death of the intracellular pathogens (32). Therefore, the P2X₇ receptor-ASK1-p38 pathway might play a critical role in host defense against invading pathogens.

CONCLUSION AND PERSPECTIVE

Although the molecular mechanism of ROS-dependent ASK1 activation has been well characterized, important questions remain unanswered. Previous studies have established that Trx is a redox-dependent negative regulator of ASK1 (13,16,17). After the dissociation of Trx in response to ROS, TRAFs are

recruited to the ASK1 signalosome. TRAFs promote homophilic interaction of ASK1 through the NCC domain, leading to the activation of ASK1. It remains unknown how Trx and TRAFs negatively and positively regulate this interaction, respectively.

TRAF6 is a RING domain-containing E3 ubiquitin ligase that synthesizes a polyubiquitin chain linked through lysine-63 of ubiquitin. TRAF6 ubiquitination has been shown to be involved in TAK1 activation. TAK1 is a member of MAP3K family capable of activating both MAPKs and NF- κ B pathways upon stimulation of IL-1 β , TNF α and ligands of several TLRs (33-35). In addition, ubiquitin-conjugating enzyme Ubc13 has been demonstrated to be a crucial component of TRAF6-mediated inflammatory responses downstream of TLR4, suggesting that TRAF6 serves as a ubiquitin ligase in TLR4-signaling (36,37). Whether TRAFs ubiquitination is necessary for ASK1 activation is yet to be determined.

Recent structural analysis of ASK1 has demonstrated that ASK1 kinase domain forms a dimer interacting in a head-to-tail orientation, which might play a pivotal role in endogenous ASK1 oligomerization (38). Structural analysis of full-length ASK1 or ASK1 complex with its regulatory proteins such as Trx, TRAF2, and TRAF6 will provide more insight into the molecular mechanism of ASK1 activation.

ASK1 appears to be important for various immune responses mediated by ROS. It has been demonstrated that ROS-dependent activation of ASK1 is required for TLR4-mediated inflammatory responses toward LPS. Furthermore, ROS-dependent P2X₇ receptor-ASK1-p38 pathway is necessary for ATP-induced apoptosis in macrophages. Nox4 and Nox2 have been suggested to be responsible for ROS generation in TLR4- and P2X₇ receptor-signaling, respectively. However, the precise mechanism of their activation remains elusive (8).

Although many MAP3Ks are identified to mediate TLR-initiated innate immune responses, their distinct roles are poorly understood (22). Notably, MEKK3 is involved in the activation of the p38 pathway downstream of TLR4 in MEFs as well as ASK1 in DCs and splenocytes (8,23). Further studies are required for elucidation of specific roles of MAP3Ks as TLR-signaling components.

Thus, we are still in the early stages of investigating ASK1-signaling in immunity. Continued research in this field will contribute to greater understanding of host defense.

ACKNOWLEDGEMENT

We thank the members of the Cell signaling Laboratory for helpful discussions.

REFERENCES

- Kyriakis JM, Avruch J. Mammalian mitogen-activated protein kinase signal transduction pathways activated by stress and inflammation. *Physiol Rev* 81:807-69, 2001
- Ichijo H, Nishida E, Irie K, ten Dijke P, Saitoh M, Moriguchi T, Takagi M, Matsumoto K, Miyazono K, Gotoh Y. Induction of apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways. *Science* 275:90-4, 1997
- Nishitoh H, Saitoh M, Mochida Y, Takeda K, Nakano H, Rothe M, Miyazono K, Ichijo H. ASK1 is essential for JNK/SAPK activation by TRAF2. *Mol Cell* 2:389-95, 1998
- Tobiome K, Matsuzawa A, Takahashi T, Nishitoh H, Morita K, Takeda K, Minowa O, Miyazono K, Noda T, Ichijo H. ASK1 is required for sustained activations of JNK/p38 MAP kinases and apoptosis. *EMBO Rep* 2:222-8, 2001
- Takeda K, Matsuzawa A, Nishitoh H, Tobiome K, Kishida S, Ninomiya-Tsuji J, Matsumoto K, Ichijo H. Involvement of ASK1 in Ca²⁺-induced p38 MAP kinase activation. *EMBO Rep* 5:161-6, 2004
- Nishitoh H, Matsuzawa A, Tobiome K, Saegusa K, Takeda K, Inoue K, Hori S, Kakizuka A, Ichijo H. ASK1 is essential for endoplasmic reticulum stress-induced neuronal cell death triggered by expanded polyglutamine repeats. *Genes Dev* 16:1345-55, 2002
- Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. *Cell* 124:783-801, 2006
- Matsuzawa A, Saegusa K, Noguchi T, Sadamitsu C, Nishitoh H, Nagai S, Koyasu S, Matsumoto K, Takeda K, Ichijo H. ROS-dependent activation of the TRAF6-ASK1-p38 pathway is selectively required for TLR4-mediated innate immunity. *Nat Immunol* 6:587-92, 2005
- Hayakawa T, Matsuzawa A, Noguchi T, Takeda K, Ichijo H. The ASK1-MAP kinase pathways in immune and stress responses. *Microbes Infect* 8:1098-107, 2006
- Skoberne M, Beignon AS, Bhardwaj N. Danger signals: a time and space continuum. *Trends Mol Med* 10:251-7, 2004
- Takeda K, Noguchi T, Naguro I, Ichijo H. Apoptosis Signal-Regulating Kinase 1 in Stress and Immune Response. *Annu Rev Pharmacol Toxicol*, 2007
- Fujino G, Noguchi T, Takeda K, Ichijo H. Thioredoxin and protein kinases in redox signaling. *Semin Cancer Biol* 16:427-35, 2006
- Noguchi T, Takeda K, Matsuzawa A, Saegusa K, Nakano H, Gohda J, Inoue J, Ichijo H. Recruitment of tumor necrosis factor receptor-associated factor family proteins to apoptosis signal-regulating kinase 1 signalosome is essential for oxidative stress-induced cell death. *J Biol Chem* 280:37033-40, 2005
- Powis G, Montfort WR. Properties and biological activities of thioredoxins. *Annu Rev Biophys Biomol Struct* 30:421-55, 2001
- Masutani H, Ueda S, Yodoi J. The thioredoxin system in retroviral infection and apoptosis. *Cell Death Differ* 12 Suppl 1:991-8, 2005
- Saitoh M, Nishitoh H, Fujii M, Takeda K, Tobiome K, Sawada Y, Kawabata M, Miyazono K, Ichijo H. Mammalian thioredoxin is a direct inhibitor of apoptosis signal-regulating kinase (ASK) 1. *EMBO J* 17:2596-606, 1998
- Fujino G, Noguchi T, Matsuzawa A, Yamauchi S, Saitoh M, Takeda K, Ichijo H. Thioredoxin and TRAF family proteins regulate reactive oxygen species-dependent activation of ASK1 through reciprocal modulation of the N-terminal homophilic interaction of ASK1. *Mol Cell Biol* 27:8152-63, 2007
- Liu H, Nishitoh H, Ichijo H, Kyriakis JM. Activation of apoptosis signal-regulating kinase 1 (ASK1) by tumor necrosis factor receptor-associated factor 2 requires prior dissociation of the ASK1 inhibitor thioredoxin. *Mol Cell Biol* 20:2198-208, 2000
- Chung JY, Park YC, Ye H, Wu H. All TRAFs are not created equal: common and distinct molecular mechanisms of TRAF-mediated signal transduction. *J Cell Sci* 115:679-88, 2002
- Tobiome K, Saitoh M, Ichijo H. Activation of apoptosis signal-regulating kinase 1 by the stress-induced activating phosphorylation of pre-formed oligomer. *J Cell Physiol* 191:95-104, 2002
- Ohto U, Fukase K, Miyake K, Satow Y. Crystal structures of human MD-2 and its complex with antiendotoxic lipid IVa. *Science* 316:1632-4, 2007
- Symons A, Beinke S, Ley SC. MAP kinase kinase kinases and innate immunity. *Trends Immunol* 27:40-8, 2006
- Huang Q, Yang J, Lin Y, Walker C, Cheng J, Liu ZG, Su B. Differential regulation of interleukin 1 receptor and Toll-like receptor signaling by MEKK3. *Nat Immunol* 5:98-103, 2004
- Dumitru CD, Ceci JD, Tsatsanis C, Kontoyannis D, Stamatakis K, Lin JH, Patriotic C, Jenkins NA, Copeland NG, Kollias G, Tsichlis PN. TNF-alpha induction by LPS is regulated posttranscriptionally via a Tpl2/ERK-dependent pathway. *Cell* 103:1071-83, 2000
- Grandvaux N, Soucy-Faulkner A, Fink K. Innate host defense: Nox and Duox on phox's tail. *Biochimie* 89:1113-22, 2007
- Park HS, Jung HY, Park EY, Kim J, Lee WJ, Bae YS. Cutting edge: direct interaction of TLR4 with NAD(P)H oxidase 4 isozyme is essential for lipopolysaccharide-induced production of reactive oxygen species and activation of NF-kappa B. *J Immunol* 173:3589-93, 2004
- Chiang E, Dang O, Anderson K, Matsuzawa A, Ichijo H, David M. Cutting edge: apoptosis-regulating signal kinase 1 is required for reactive oxygen species-mediated activation of IFN regulatory factor 3 by lipopolysaccharide. *J Immunol* 176:5720-4, 2006
- Bodin P, Burnstock G. Purinergic signalling: ATP release. *Neurochem Res* 26:959-69, 2001

29. Khakh BS, North RA. P2X receptors as cell-surface ATP sensors in health and disease. *Nature* 442:527-32, 2006
30. Ferrari D, Pizzirani C, Adinolfi E, Lemoli RM, Curti A, Idzko M, Panther E, Di Virgilio F. The P2X7 receptor: a key player in IL-1 processing and release. *J Immunol* 176:3877-83, 2006
31. Lister MF, Sharkey J, Sawatzky DA, Hodgkiss JP, Davidson DJ, Rossi AG, Finlayson K. The role of the purinergic P2X7 receptor in inflammation. *J Inflamm (Lond)* 4:5, 2007
32. Lammas DA, Stober C, Harvey CJ, Kendrick N, Panchalingam S, Kumararatne DS. ATP-induced killing of mycobacteria by human macrophages is mediated by purinergic P2Z(P2X7) receptors. *Immunity* 7:433-44, 1997
33. Wang C, Deng L, Hong M, Akkaraju GR, Inoue J, Chen ZJ. TAK1 is a ubiquitin-dependent kinase of MKK and IKK. *Nature* 412:346-51, 2001
34. Adhikari A, Xu M, Chen ZJ. Ubiquitin-mediated activation of TAK1 and IKK. *Oncogene* 26:3214-26, 2007
35. Sato S, Sanjo H, Takeda K, Ninomiya-Tsuji J, Yamamoto M, Kawai T, Matsumoto K, Takeuchi O, Akira S. Essential function for the kinase TAK1 in innate and adaptive immune responses. *Nat Immunol* 6:1087-95, 2005
36. Yamamoto M, Okamoto T, Takeda K, Sato S, Sanjo H, Uematsu S, Saitoh T, Yamamoto N, Sakurai H, Ishii KJ, Yamaoka S, Kawai T, Matsuura Y, Takeuchi O, Akira S. Key function for the Ubc13 E2 ubiquitin-conjugating enzyme in immune receptor signaling. *Nat Immunol* 7:962-70, 2006
37. Fukushima T, Matsuzawa S, Kress CL, Bruey JM, Krajewska M, Lefebvre S, Zapata JM, Ronai Z, Reed JC. Ubiquitin-conjugating enzyme Ubc13 is a critical component of TNF receptor-associated factor (TRAF)-mediated inflammatory responses. *Proc Natl Acad Sci USA* 104:6371-6, 2007
38. Bunkoczi G, Salah E, Filippakopoulos P, Fedorov O, Muller S, Sobott F, Parker SA, Zhang H, Min W, Turk BE, Knapp S. Structural and functional characterization of the human protein kinase ASK1. *Structure* 15:1215-26, 2007