

EID3 Promotes Glioma Cell Proliferation and Survival by Inactivating AMPK α 1

Yaoxian Xiang,^{1,2,3,*} Lei Zhu,^{4,*} Zijian He,^{5,*} Lei Xu,^{1,2,3} Yuhang Mao,⁵ Junjian Jiang,^{1,2,3} Jianguang Xu^{1,2,3,6}

Department of Hand Surgery,¹ Huashan Hospital, Fudan University, Shanghai, China

NHC Key Laboratory of Hand Reconstruction (Fudan University),² Ministry of Health, Shanghai, China

Shanghai Key Laboratory of Peripheral Nerve and Microsurgery,³ Shanghai, China

Department of Radiology,⁴ Beijing Electric Power Hospital, Beijing, China

Department of Neurosurgery,⁵ Minhang Hospital, Fudan University, Shanghai, China

School of Rehabilitation Science,⁶ Shanghai University of Traditional Chinese Medicine, Shanghai, China

Objective : EID3 (EP300-interacting inhibitor of differentiation) was identified as a novel member of EID family and plays a pivotal role in colorectal cancer development. However, its role in glioma remained elusive. In current study, we identified EID3 as a novel oncogenic molecule in human glioma and is critical for glioma cell survival, proliferation and invasion.

Methods : A total of five patients with glioma were recruited in present study and fresh glioma samples were removed from patients. Four weeks old male non-obese diabetic severe combined immune deficiency (NOD/SCID) mice were used as transplant recipient models. The subcutaneous tumor size was calculated and recorded every week with vernier caliper. EID3 and AMP-activated protein kinase α 1 (AMPK α 1) expression levels were confirmed by real-time polymerase chain reaction and Western blot assays. Colony formation assays were performed to evaluate cell proliferation. Methyl thiazolyl tetrazolium (MTT) assays were performed for cell viability assessment. Trypan blue staining approach was applied for cell death assessment. Cell Apoptosis DNA ELISA Detection Kit was used for apoptosis assessment.

Results : EID3 was preferentially expressed in glioma tissues/cells, while undetectable in astrocytes, neuronal cells, or normal brain tissues. EID3 knocking down significantly hindered glioma cell proliferation and invasion, as well as induced reduction of cell viability, apoptosis and cell death. EID3 knocking down also greatly inhibited tumor growth in SCID mice. Knocking down of AMPK α 1 could effectively rescue glioma cells from apoptosis and cell death caused by EID3 absence, indicating that AMPK α 1 acted as a key downstream regulator of EID3 and mediated suppression effects caused by EID3 knocking down inhibition. These findings were confirmed in glioma cells generated patient-derived xenograft models. AMPK α 1 protein levels were affected by MG132 treatment in glioma, which suggested EID3 might down regulate AMPK α 1 through protein degradation.

Conclusion : Collectively, our study demonstrated that EID3 promoted glioma cell proliferation and survival by inhibiting AMPK α 1 expression. Targeting EID3 might represent a promising strategy for treating glioma.

Key Words : Glioma · EP300-interacting inhibitor of differentiation · AMP activated protein kinase α 1.

• Received : December 7, 2021 • Revised : February 7, 2022 • Accepted : February 26, 2022

• Address for correspondence : **Jianguang Xu**

Department of Hand Surgery, Huashan Hospital, Fudan University, 12 Wulumuqi Middle Road, Shanghai 200040, China

Tel : +86-21-52887090, Fax : +86-21-52888055, E-mail : jianguangxu@fudan.edu.cn, ORCID : <https://orcid.org/0000-0003-2077-2588>

Junjian Jiang

Department of Hand Surgery, Huashan Hospital, Fudan University, 12 Wulumuqi Middle Road, Shanghai 200040, China

Tel : +86-21-52887092, Fax : +86-21-52888055, E-mail : aizhuzhu20140626@163.com, ORCID : <https://orcid.org/0000-0002-1133-0473>

*These authors contributed equally to this work.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Gliomas comprise about one third of all brain tumors and more than three quarter of all malignant brain tumors^{1,2,32}. It is considered as one of the most aggressive human malignancies^{5,14}. The prognosis of glioma patients is extremely poor due to lack of approaches for early-diagnosis^{18,24}. And rapid progression of glioma also greatly contributes to its dismal outcomes^{26,38}. Currently, molecule-targeted therapeutic strategy has become the focus of glioma research, and showed promising prospects, which leads to an urgent demand for novel therapeutic targets of glioma.

EID3 (EP300-interacting inhibitor of differentiation) was identified as a novel member of EID family, which inhibits CREB binding protein (CBP)-dependent co-activation³. Previous studies revealed that it could inhibit cellular differentiation by binding to class I histone deacetylases (HDAC) or CBP/P300 complex^{3,15,28}. A pilot study reported that EID3 serves as a key regulator of cancer stem cell in colorectal cancer²⁷. With decreased proteasome activity, EID3 is transcriptionally up-regulated, resulting in activation of Wnt- β -catenin signaling pathway, which plays a vital role in proliferation, invasion, and self-renew of colorectal cancer stem cells²⁷. However, the function of EID3 in glioma cell remained elusive.

AMP-activated protein kinase (AMPK) is identified as a master regulator of energy and metabolism with a highly-conserved trait^{4,23}. Moreover, AMPK can restore energy balance during metabolic stress^{31,39}. Previous investigations reported that AMPK acted as a vital suppressor in cancer through various ways^{7,36,41}. Activated AMPK could inhibit the activation of mechanistic target of rapamycin complex 1 (mTORC1), which is considered as a key regulator of oncogenic cascade in tumor development and progression^{8,10,16}. Moreover, recent studies revealed that activated AMPK could trigger autophagy to inhibit cancer cell proliferation and induce cell death^{19,25}. Additionally, AKT-MDM2-Foxo3 cascade is blocked after AMPK activation⁹. AMPK was also reported to be highly suppressed in glioma, and restoration of AMPK expression in glioma showed promising therapeutic potential⁴⁰. The identifying of AMPK as a novel key upstream regulator in glioma might imply a novel strategy to glioma therapy. We also search AMPK α 1, one of major regulators in AMPK signal pathway, in the Cancer Genome Atlas data set and found that it was a mutant gene in the pathogenesis of glioma, and the mutation rate was 57.14%. The

type of mutation is substitution, and the result of mutation is missense.

In this study, we explored the function of EID3 in glioma cell proliferation, survival, death, and invasion. Our data identified EID3 as a novel upstream inhibitor of APMK, which provide a new insight into the mechanism of glioma progression and indicated that EID3 might be a promising target for glioma therapy.

MATERIALS AND METHODS

The protocols of using human cells and tissues were in accordance with the principle of Declaration of Helsinki, and were approved by the Ethics Review Board (ERB) of Minhang Hospital.

Cell culture

A172 glioma cell line was purchased from the Cell Bank of Chinese Academy of Sciences (Shanghai, China). A172 cells were cultured in DMEM medium (Gibco, New York, NY, USA) supplemented with 10% fetal bovine serum (FBS) (Invitrogen, Carlsbad, CA, USA), 100 U/mL penicillin and 100 μ g/mL streptomycin (Gibco) at 37°C in a humidified incubator (Thermo-Fisher, Waltham, MA, USA) with 5% CO₂. All experiments were conducted with cells passage less than eight times. DNA fingerprinting and profiling were performed every 5 months to verify the cell line phenotype. Human astrocyte culture was performed according to previous studies^{11,21}.

Clinical glioma tissues

A total of five patients with glioma were recruited in present study (male, World Health Organization grade III, age 54–68) at Minhang Hospital. Right after surgery, tumor tissues and surrounding normal brain tissues were collected, and separated carefully under microscope. Glioma was defined by pathological tests with both fresh frozen section tissue and formalin-fixed paraffin-embedded (anaplastic astrocytoma). Tissues were carefully washed with pre-cold PBS, and then minced for homogenization with tissue lysis buffer (Invitrogen). Written informed consent was obtained from each participant.

Cell viability, death, and apoptosis evaluations

Methyl thiazolyl tetrazolium (MTT) assays were performed

for cell viability assessment. After different treatments, 0.5 mg/mL MTT reagent was added into the culture medium and then incubated for 4 hours at 37°C. Purple formazan salt crystals were dissolved by adding the solubilization solution with 10% sodium dodecyl sulfate (SDS) and 0.01 M HCl. The absorption at 490 nm was measured using a multi-well plate reader.

For cell death assessment, trypan blue staining approach was applied. Attached cells were digested with trypsin/ethylenediamine tetraacetic acid (EDTA), suspended in 1X PBS, and mixed with 0.4% trypan blue dye (Sigma, St. Louis, MO, USA). Dead cells would take up trypan blue due to compromised cell membranes, while viable cells would not due to the membrane integrity. The percentages of dead cell were recorded for further assessments. Cell Apoptosis DNA ELISA Detection Kit (Roche, San Francisco, CA, USA) was used for apoptosis assessment according to manufacturer's instructions.

Cell proliferation, migration, and invasion evaluations

To evaluate cell proliferation, colony formation assays were performed. A172 cells were harvested with trypsin/EDTA and re-suspended in pre-cold PBS before being seeded in a 6-well plate at a density of 500 cells per well with completed DMEM medium (10% FBS). The cells were cultured for 21 days before the cell colonies were stained with crystal violet solution (Sigma). Number of viable colonies was recorded.

EID3, and AMPK α 1 knocking down

Control lentivirus pLKO.1 vector and pLKO.1 vectors containing shRNA for EID3 (RHS4531-EG493861) were purchased from Thermo Fisher Scientific (Waltham, MA, USA). Glioma cells were infected with viral particles twice for the establishment of stable EID3 knocking down clones. Stable transfected cells were selected with puromycin (2 μ g/mL). Knocking down efficiency were confirmed by real-time polymerase chain reaction (RT-PCR) and Western blot (WB) assays.

Two lentiviral AMPK α 1 shRNAs were purchased from Santa Cruz Biotech (shAMPK α 1-1; Santa Cruz, CA, USA) and GeneChem (shAMPK α 1-2; Montreal, Canada). Each shRNA was added into A172 cells separately for 24 hours. The cell culture medium was then replaced by fresh medium for an additional 24 hours. Stable clones expressing AMPK α shRNA were

selected by puromycin (0.5 μ g/mL; Sigma) for a period of 10 days. Control cells were infected with lentiviral scramble shRNA (Santa Cruz Biotech). Knocking down efficiencies were evaluated by RT-PCR and WB assays.

Construction of AMPK α 1 dominant negative mutation

The dominant negative AMPK α (DN-AMPK α , T172A) construct was designed as mentioned before. Glioma cells were cultured in a medium without antibiotics when reached 60–70% confluence. Then DN-AMPK α cDNA (0.10 μ g/mL) was transfected into the glioma cells followed by Lipofectamine 2000 protocol. Afterwards, stable cells were selected via neomycin (1.0 μ g/mL; Sigma). Transfection efficiency was verified via WB.

RT-PCR assays

Total RNA was extracted from cells using RNeasy Mini Kit (Qiagen, Dusseldorf, Germany), and reversed transcript using Quantitect Reverse Transcription Kit (Qiagen) according to the manufacturer's instructions. The RT-PCR mixtures contained 2 \times QuantiTect SYBR Green PCR Master Mix (Qiagen), 10 \times QuantiTect primer assay mix, and synthesized cDNA. Amplification was performed in triplicate for each sample on a Lightcycler 480 platform (Roche, Mannheim, Germany). EID3 protein expression levels were normalized to glyceraldehyde-3-phosphate dehydrogenase (GAPDH) levels. Primer sequences were shown as follows: GAPDH, forward (F), TGCACCACCAACTGCTTAGC; reverse (R), GGCATGGACTGTGGTCATGAG; EID3, F, GGAGGCACCTCAGTCATGATA; R, GTCCACAAA-CAGTTCTCTGAC.

WB assays

Cells were lysed in RIPA lysis buffer (Beyotime, Shanghai, China) with PMSF (Roche, Mannheim, Germany). Supernatant was collected after centrifugation at 13000 \times g at 4°C for 10 minutes. Protein concentration was measured with a BCA protein kit (Beyotime) and the whole lysates were mixed with 4 \times SDS loading buffer (125 mmol/Tris-HCl, 4% SDS, 20% glycerol, 100 mmol/L DTT, and 0.2% bromophenol blue). Protein samples were heated to 100°C for 5 minutes. For the WB, 30 μ g total protein samples were loaded to a SDS-polyacrylamide gel. After electrophoresis, the protein was trans-

ferred to a polyvinylidene fluoride membrane blocked in 5% fat-free milk and incubated with antibodies against EID3 (1 : 800; Abcam, Burlingame, CA, USA), AMPK α 1 (1 : 1000; CST, Chicago, IL, USA), pS6K1 (1 : 1000; Abcam), S6K1 (1 : 750; Abcam), LC3B-I (1 : 1000; Abcam), LC3B-II (1 : 800; Abcam), Beclin (1 : 750; Pharmingen, San Diego, CA, USA), ATG-5 (1 : 800; Abcam), ATG-7 (1 : 1000; Abcam), p62 (1 : 1000; Abcam), pAMPK α 1 (1 : 1000; Abcam), ACC (1 : 1000; Abcam), pACC (1 : 1000; Abcam), and Tubulin (1 : 1000, Proteintech, Wuhan, China) overnight at 4°C, followed by another 2 hours incubation with horseradish peroxidase-conjugated anti-rabbit secondary antibody (1 : 5000; Jackson ImmunoResearch Labs, West Grove, PA, USA). GAPDH was served as a loading control. NIH Image J software (Bethesda, MD, USA) was used for the densitometric analysis of the WB. The protein levels were normalized with GAPDH.

Glioma Xenograft assays

Specific pathogen-free BALB/c nude mice (4–6 weeks old) purchased from the Chinese Academy of Medical Science were injected with 5×10^6 glioma cells subcutaneously into the left upper flank regions. The subcutaneous tumor size was calculated and recorded every week with vernier caliper as follows : tumor volume (mm^3) = $(L \times W^2) / 2$, where L=long axis and W=short axis, the measurements were repeated three times.

Construction of patients-derived xenograft models and primary glioma cells

Glioma patient-derived xenograft (PDX) models were generated according to a previous study¹⁷. Briefly, 4 weeks old male non-obese diabetic severe combined immune deficiency (NOD/SCID) mice were used as transplant recipient models and raised in the aseptic environment. Fresh glioma samples were cut into 1mm pieces within 1 hour after removal from patients. Tissue fragments were incubated in DMEM medium mixed with 50% Matrigel™ (BD, Franklin Lakes, NJ, USA), 10 ng/mL epidermal growth factor (Gibco), 10 ng/mL basic fibroblast growth factor (Gibco), 100 U/mL penicillin, and 100 U/mL streptomycin for 30 minutes. The whole tumor tissue mixture was then transplanted into the right flanks of mice (n=3; 4–5 weeks old; Shanghai Institute of Material Medicine, Chinese Academy of Science) subcutaneously with a #20 trocar. Animal welfare and experimental protocols were ap-

proved by the Shanghai Medical Experimental Animal Care Commission. Primary EID3 expression levels were detected by RT-PCR and WB prior to tissue transplantation. Single glioma cell suspension was made according to a previous study⁶. All the samples were mechanically disaggregated and digested with type IV collagenase (Gibco) within one hour, and resuspended in DMEM medium. The treated tissue was filtrated through a 40 μm filter to obtain the single-cell suspension. Red blood cells were lysed with ACK buffer (Invitrogen). The number of viable cells was counted and analyzed by trypan blue analysis (Procell, Wuhan, China).

Statistical analysis

Statistical analyses were performed with SPSS ver. 20.0 for Mac (IBM, Armonk, NY, USA). Quantitative data presented as mean \pm standard error are from at least three independent experiments. Continuous data were analyzed by one-way analysis of variance and Student's t-test, and categorical data were analyzed by Fisher's exact test or chi-square test. A *p*-value <0.05 was considered statistically significant.

RESULTS

EID3 is preferentially expressed in glioma cells and correlated with AMPK α 1

EID3 expression level in glioma cells was evaluated by quantitative PCR and WB. Results of RT-PCR assays showed that EID3 only expressed in glioma cells, and undetectable in human astrocyte and neuronal cells (HCN-1a cell line) (Fig. 1A). WB assays further validated the findings of RT-PCR (Fig. 1B). Since AMPK was identified as a key suppressor in glioma, we next investigate the correlation between EID3 and AMPK α 1, an active form of AMPK. The WB results showed that EID3 expression was correlated with AMPK α 1 downregulation (Fig. 1B). AMPK inhibition could result in mTORC1 activation process, during which phosphorylation of S6K1 was considered as a hallmark. Hence, we further explore the S6K1 activation status. Our data showed that the S6K1 phosphorylation was increased in AMPK α 1-low A172 cells. However, the pS6K1 level was low in EID3-null astrocytes and HCN-1a cells which with a high AMPK α 1 expression (Fig. 1B).

EID3 and AMPK α 1 expression status were also evaluated in human glioma tissues. As shown in Fig. 1C, EID3 expression

could be detected in four of five patients (80%) enrolled, however, it could not be detected in surrounding normal brain tissues. Also, EID3 expression was correlated with downregulation of AMPK α 1, which consistent with the results in A172 cells.

EID3 knocking down inhibited proliferation, survival, and invasion of glioma cell

EID3 expression was knocking down with two shRNAs targeting different sites of EID3 cDNA, the efficiencies of knocking down were validated through RT-PCR and WB (Fig. 2A). Remarkably, EID3 knocking down could restore AMPK α 1 expression, resulting in inhibition of S6K1 (Fig. 2A). Colony formation assays demonstrated that EID3 knocking down significantly inhibited A172 cell proliferation ($p < 0.05$, Fig. 2B). Meanwhile, Transwell assays indicated that the number of migrated and invaded cells was significantly decreased when EID3 expression is knocked down (Fig. 2C and Supplementary Fig. 1). Based on the MTT results, we found the number of viable cells was significantly decreased after EID3 knocking

down. We also found that EID3 shRNA could induce cell death according to trypan blue staining. Additionally, Histone DNA apoptosis ELISA assay revealed EID3 shRNA could induce apoptosis in glioma cells (Fig. 2D), while the scramble non-sense control shRNA exhibited no effects on glioma cell proliferation, death, and apoptosis.

According to a previous report²²⁾, AMPK activation would trigger autophagy by activating Ulk1 or inhibiting mTOR signaling. Since our results indicated that EID3 could restore mTOR activation by downregulating AMPK α 1, we further investigated the effects of EID3 on autophagy in glioma cells. Our data indicated that EID3 shRNA could induce increased Beclin-1, ATG-5, and ATG-7 expression, LC3B-I to LC3B-II switch, and p62 degradation, which confirmed that EID3 knocking down glioma cells exhibited autophagy-like changes (Fig. 2E).

We also validated *in vivo* proliferation inhibition of EID3. A172 cells that stable express EID3 shRNAs and parental control A172 cells were injected into the flanks of SCID mice, and tumor volumes and weights were recorded. As Fig. 2F demonstrated, tumors generated from EID3 knocking down A172 cells grew significantly slower than control tumors (sh-C). Moreover, weights of tumor generated from EID3 knocking down A172 cells were significantly lighter than those of control tumors.

Validation of AMPK α 1 as the key downstream molecule of EID3 pathway, EID3 might decrease AMPK α 1 level by promoting protein degradation

Our results indicated that EID3 knocking down could restore AMPK α 1 expression and inhibit tumor growth and invasion. We further explored whether AMPK α 1 served as the key inhibitor that mediated the inhibition effects following EID3 knocking down. For this purpose, AMPK α 1 was knocked down in A172 cells expressing shEID3 (Fig. 3A). We found that AMPK α 1 inhibition could restore the activation of S6K1 in EID3 knocking down cells (Fig. 3A). However, AMPK α 1 knocking down did not affect expression status of EID3. Moreover, cell viability reduction and cell apoptosis caused by EID3 knocking down could be largely attenuated by AMPK α 1 knocking down inhibition (Fig. 3B).

To further study the role of AMPK α 1, a dominant negative AMPK α 1 mutant (DN-AMPK α 1, T172A) were introduced into A172 cells, aiming to block AMPK α 1 activation following




Fig. 1. EID3 preferentially expressed in glioma cells and correlated with AMP-activated protein kinase α 1 (AMPK α 1) downregulation. EID3 preferentially expressed in glioma cells and correlated with AMPK α 1 downregulation. A : mRNA expression of EID3 in primary astrocyte, neuronal cell (HCN-1a), and A172 glioma cells assessed by real-time polymerase chain reaction assays. B : Protein expression of EID3, AMPK α 1, pS6K1, and S6K1 in primary astrocyte, neuronal cell (HCN-1a), and A172 glioma cells assessed by WB assays. Tubulin was used as an internal control. C : EID3 and AMPK α 1 protein expression in human glioma tissues and paired surrounding normal brain tissues assessed by WB assays. EID3 : EP300-interacting inhibitor of differentiation, WB : Western blot.

EID3 knocking down. WB assays confirmed exogenous expression of the DN-AMPK α 1 in A172 cells (Fig. 3C). As expected, EID3 knocking down inhibition greatly activated AMPK α 1, which can be proven by increased expression levels of p-AMPK α 1 and p-ACC (Fig. 3D). However, such effects

were significantly hindered by DN-AMPK α 1 (Fig. 3D). Consistently, forced expression of DN-AMPK α 1 did not rescue the inhibition effects of EID3 knocking down on cell survival (Fig. 3E) and death (Fig. 3F). Meanwhile, all control shRNAs showed no significant effects on A172 cells. Through RT-PCR




Fig. 2. Biological functions of EID3 in glioma. A : Stable A172 cells expressing EID3 shRNAs (sh-1, sh-2), and scramble control shRNA were subjected to real-time polymerase chain reaction and WB assays. EID3 expression was knocking down efficiently with two shRNAs targeting different sites of EID3 cDNA. EID3 knocking down could restore AMPK α 1 and inhibit S6K1 expression. B : Colony formation assays were used to evaluate the effects of EID3 knocking down on glioma cell proliferation. C : Transwell assays were conducted to evaluate the effects of EID3 knocking down on glioma cell migration and invasion. D : Effects of EID3 knocking down on glioma cell apoptosis (left), survival (middle), and death (right) were evaluated by certain function assays. EID3 shRNA could induce apoptosis in glioma cells compared to the scramble non-sense control shRNA. E : Expressions of listed autophagy-associated genes were detected by WB assays after EID3 knocking down. F : *In vivo* growth curves (left) and weights (right) of glioma tumors in SCID mice. The tumors generated from EID3 knocking down A172 cells grew significantly slower than control tumors. * $p < 0.05$ vs. sh-C. EID3 : EP300-interacting inhibitor of differentiation, AMPK α 1 : AMP-activated protein kinase α 1, ELISA : enzyme linked immunosorbent assay, OD : optical density, MTT : methyl thiazolyl tetrazolium, WB : Western blot, SCID : severe combined immune deficiency.

assays, we found that neither EID3 shRNA could decrease AMPK α 1 mRNA expression (Fig. 3G), which suggested EID3 regulates AMPK α 1 expression through a non-transcriptional way. Therefore, MG132, a proteasome inhibitor, was intro-

duced to test our hypothesis. WB assays showed that MG132 restored AMPK α 1 expression in A172 cells with high EID3 expression (Fig. 3H). Also, MG132 reactivated the mTOR signaling, supported by increased pS6K1 level after treatment




Fig. 3. AMP-activated protein kinase (AMPK) signaling activation mediated cell death and apoptosis caused by EID3 knocking down. A : Validation of the transfection efficiencies of AMPK α 1 silence in stable A172 cells that expressing EID3 shRNAs. B : Effects of AMPK α 1 knocking down on apoptosis (left), survival (middle), and death (right) of glioma cell expressing EID3 shRNAs. The cell viability reduction and cell apoptosis could be largely attenuated by AMPK α 1 knocking down. C : Validation of the transfection efficiencies of DN-AMPK α 1. D : Effects of DN-AMPK α 1 transfection on the activation status of AMPK and ACC were evaluated by WB assays. E and F : Effects of DN-AMPK α 1 transfection on glioma survival and death in A172 cells expressing control shRNA or EID3 shRNAs. The forced expression of DN-AMPK α 1 did not rescue the inhibition effects of EID3 knocking down on cell survival and death. G : mRNA expression of AMPK α 1 assessed by real-time polymerase chain reaction (RT-PCR) after EID3 knocking down. Neither EID3 shRNA could decrease AMPK α 1 mRNA expression. H : Protein expression of AMPK α 1 assessed by WB after MG132 treatment. MG132 showed no significant influence on the mRNA expression of AMPK α 1 ($p=0.677$). I : mRNA expression of AMPK α 1 assessed by RT-PCR after MG132 treatment. MG132 showed no significant influence on the mRNA expression of AMPK α 1 ($p=0.677$). J : Protein expression of AMPK α 1 assessed by WB after cycloheximide treatment in EID3 knocking down A172 cells. * $p<0.05$ vs. sh-C. † $p<0.05$ vs. no shRNA. ‡ $p<0.05$ vs. parental or vector in shEID3. EID3 : EP300-interacting inhibitor of differentiation, DN : dominant negative, ACC : acetyl-CoA carboxylase, MTT : methyl thiazolyl tetrazolium, OD : optical density, WB : Western blot.

(Fig. 3H). However, MG132 showed no significant influence on the mRNA expression of AMPK α 1 (Fig. 3I). To exclude the possibility that EID3 inhibits the protein synthesis process of AMPK α 1, cycloheximide (CHX), an inhibitor of protein synthesis was used. Results showed that CHX treatment could

not affect the expression level of AMPK α 1 in EID3 knocking down A172 cells (Fig. 3J), which indicated that increased expression of AMPK α 1 in EID3 knocking down cells might not due to promoted protein synthesis. Collectively, these data implied that EID3 might induce ubiquitination and protea-




Fig. 4. AMP-activated protein kinase (AMPK) mediated viability reduction and cell death in EID3 knocking down primary glioma cells. A : Protein expression of EID3, AMPK α 1, pS6K1, and S6K1 in three tumor with descending EID3 level from PDX models. B : Growth curves indicated that EID3 expression level correlated with glioma growth speed in PDX models. T1 (high EID3) grew significantly faster than T2 and T3. T2 also showed significantly faster growth than T3 did. C : Effects of EID3 knocking down on the expression level of AMPK α 1, pS6K1, and S6K1 in primary glioma cells generated from PDX model T1 (high EID3 expression). D : Expressions of listed autophagy-associated genes were detected by Western blot assays after EID3 knocking down in primary glioma cells generated from PDX model T1. E : Effects of AMPK α 1 knocking down on S6K1 activation status in primary glioma cells expressing EID3 shRNA3. F : Effects of AMPK α 1 knocking down on cell survival (left) and death (right) in primary glioma cells expressing EID3 shRNA3. AMPK α 1 knocking down largely attenuated the viability reduction and cell death caused by EID3 inhibition ($p < 0.05$). * $p < 0.05$. EID3 : EP300-interacting inhibitor of differentiation, MTT : methyl thiazolyl tetrazolium, OD : optical density, PDX : patient-derived xenograft.

somal degradation of AMPK α 1.

AMPK mediated viability reduction and cell death in EID3 knocking down primary glioma cells

To validate the function of EID3 in primary glioma, we generated three patient-derived xenograft model with descending EID3 expression (T1, high; T2, moderate; T3, low). The EID3 expression status were confirmed by WB assays (Fig. 4A). Accordingly, primary glioma with high EID3 expression showed lower level of AMPK α 1 but high pS6K1 expression (T1), while tumor with undetectable EID3 exhibited high level of AMPK α 1 and low level pS6K1 (T3, Fig. 4A). We observed that T1 (high EID3) grew significantly faster than T2 and T3, based on bigger tumor volumes (Fig. 4B). Moreover, T2 also showed significantly faster growth than T3 did (Fig. 4B).

We then separated single glioma cells from T1 tumor tissue and transfected it with EID3 shRNA to explore the effects of EID3 knocking down in primary glioma. WB assays confirmed the efficiencies of EID3 knocking down and demonstrated that EID3 knocking down resulted in upregulation of AMPK α 1 and inactivation of S6K1 (Fig. 4C). Furthermore, we also observed an autophagy-like phenotype in primary glioma cells with EID3 knocking down, which were similar to those findings in A172 cells (Fig. 4D). To further validate the role of AMPK α 1, primary T1 glioma cells were transfected with AMPK α 1 shRNA. Notably, AMPK α 1 knocking down could effectively restore the pS6K1 level in primary glioma cells transfected with EID3 shRNAs (Fig. 4E). Moreover, AMPK α 1 knocking down largely attenuated the viability reduction and cell death caused by EID3 inhibition (Fig. 4F). These results in primary glioma cells further confirmed that AMPK mediated viability reduction and cell death by EID3 knocking down.

DISCUSSION

Present study identified EID3 as a novel oncogenic molecule in human glioma, and was critical for glioma cell survival, proliferation and invasion. EID3 was preferentially expressed in glioma tissues/cells, while undetectable in astrocytes, neuronal cells, or normal brain tissues. Importantly, EID3 knocking down significantly inhibited *in vivo* tumor growth in SCID mice. Moreover, we found EID3 expression positively correlated with tumor growth in PDX models. AMPK α 1 was

identified as the key downstream regulator underlying EID3 that mediated its suppression effects when being knocked down, and EID3 might downregulate AMPK α 1 through protein degradation.

EID3 was identified as a negative regulator of cellular differentiation, as it binding to class I HDAC or CBP/p300^{15,27}. However, its function in human cancer, especially in glioma, remained largely unclear. A pilot study revealed that EID3 might be a critical promoter for the carcinogenesis and development of colorectal cancer²⁷, which led to our hypothesis that EID3 might also serve as an oncogene in glioma. Our results showed that EID3 knocking down significantly hindered glioma cell proliferation and invasion, induced cell viability reduction, apoptosis, and cell death. Therefore, we believed EID3 acted as a vital regulator in glioma that promoting tumor progression. This characteristic and novel function of EID3 was shown for the first time in human glioma.

AMPK has been considered as a critical tumor suppressor for a long time, and AMPK signaling has been reported to be dysregulated in various kinds of solid tumors²⁰. In lung and cervical cancer, AMPK signaling was inhibited by loss-of-function or deletion of upstream activator^{13,29}. In hepatocellular carcinoma (HCC), AMPK signaling was inhibited by insulin growth factor pathway activation, and metformin showed anti-HCC effect via reactivating AMPK signaling^{35,37}. In glioma, AMPK signaling was also constantly inhibited, and several upstream regulators, such as MAGEA6, have been identified³⁰. In present study, we showed that EID3 served as a novel AMPK signaling suppressor, since EID3 expressed glioma tissues/cells showed decreased expression of AMPK α 1. Meanwhile, AMPK α 1 knocking down could effectively rescue glioma cells from apoptosis and death caused by EID3 silence. However, we only figured out that EID3 might downregulate AMPK α 1 expression through proteasomal degradation, while detailed molecular mechanism involved in this process remained unclear. Further investigation is needed to explore EID3 induced AMPK α 1 degradation and key molecules involved in this process.

mTOR hyper-activation was considered as a hallmark in glioma, which greatly contributed to carcinogenesis³³. Therefore, great efforts have been made to develop novel regimens targeting mTOR or it regulated signaling pathways. Previous study showed that AMPK activation could suppress mTOR signaling through various ways, such as phosphorylation of

TSC2, which was a key upstream inhibitor of mTOR signaling³⁴). AMPK could also phosphorylate and inactivate raptor to block mTOR signaling activation¹²). Restoration of AMPK signaling might be an ideal strategy for preventing glioma progression. In present study, we showed that AMPK expression recovered from EID3 silencing, and recovery of AMPK could effectively inhibited mTOR activation, resulting in viability reduction, cell death and apoptosis. Significantly, mTOR signaling could be restored in EID3 knocking down cells through transfecting AMPK α 1 shRNA or plasmid containing dominant negative AMPK α 1 mutation. Our data implied that AMPK activation caused by EID3 silence was responsible for the inhibition of mTOR signaling in glioma.

CONCLUSION

Taken together, our study demonstrated that EID3 promoted glioma cell proliferation and survival through inducing AMPK α 1 downregulation. These findings strongly implied EID3 as a novel biomarker and target for glioma. Notably, targeting EID3 might represent a promising strategy for treating mTOR in glioma. Further investigations aiming to screen and identify selective inhibitor EID3 are urgently needed.

AUTHORS' DECLARATION

Conflicts of interest

No potential conflict of interest relevant to this article was reported.

Informed consent

Informed consent was obtained from all individual participants included in this study.

Author contributions

Conceptualization : YX; Data curation : LZ; Formal analysis : LZ, ZH; Funding acquisition : JX, JJ; Methodology : YX, YM; Project administration : JJ; Visualization : LX; Writing - original draft : YX; Writing - review & editing : JX

Data sharing

None

Preprint

None

ORCID

Yaoxian Xiang	https://orcid.org/0000-0003-0594-7458
Lei Zhu	https://orcid.org/0000-0002-5012-3610
Zijian He	https://orcid.org/0000-0003-1765-1558
Lei Xu	https://orcid.org/0000-0003-1750-9164
Yuhang Mao	https://orcid.org/0000-0001-9951-077X
Junjian Jiang	https://orcid.org/0000-0002-1133-0473
Jianguang Xu	https://orcid.org/0000-0003-2077-2588

• Acknowledgements

This study was supported by Shanghai Municipal Key Clinical specialty (shslczdzk05601).

• Supplementary materials

The online-only data supplement is available with this article at <https://doi.org/10.3340/jkns.2021.0298>.

References

1. Almiron Bonnin DA, Havrda MC, Israel MA : Glioma cell secretion: a driver of tumor progression and a potential therapeutic target. **Cancer Res** **78** : 6031-6039, 2018
2. Barthel FP, Wesseling P, Verhaak RGW : Reconstructing the molecular life history of gliomas. **Acta Neuropathol** **135** : 649-670, 2018
3. Bävner A, Matthews J, Sanyal S, Gustafsson JA, Treuter E : EID3 is a novel EID family member and an inhibitor of CBP-dependent co-activation. **Nucleic Acids Res** **33** : 3561-3569, 2005
4. Bilanges B, Alliouachene S, Pearce W, Morelli D, Szabadkai G, Chung YL, et al. : Vps34 PI 3-kinase inactivation enhances insulin sensitivity through reprogramming of mitochondrial metabolism. **Nat Commun** **8** : 1804, 2017
5. Boussiotis VA, Charest A : Immunotherapies for malignant glioma. **Oncogene** **37** : 1121-1141, 2018
6. Cai S, Li Y, Bai JY, Zhang ZQ, Wang Y, Qiao YB, et al. : Gai3 nuclear translocation causes irradiation resistance in human glioma cells. **Oncotarget** **8** : 35061-35068, 2017
7. Carling D : AMPK signalling in health and disease. **Curr Opin Cell Biol** **45** : 31-37, 2017
8. Carroll B, Dunlop EA : The lysosome: a crucial hub for AMPK and

- mTORC1 signalling. **Biochem J** 474 : 1453-1466, 2017
9. Chou CC, Lee KH, Lai IL, Wang D, Mo X, Kulp SK, et al. : AMPK reverses the mesenchymal phenotype of cancer cells by targeting the Akt-MDM2-Foxo3a signaling axis. **Cancer Res** 74 : 4783-4795, 2014
 10. Cork GK, Thompson J, Slawson C : Real talk: the inter-play between the mTOR, AMPK, and hexosamine biosynthetic pathways in cell signaling. **Front Endocrinol (Lausanne)** 9 : 522, 2018
 11. Cui Y, Zhao J, Yi L, Jiang Y : microRNA-153 targets mTORC2 component rictor to inhibit glioma cells. **PLoS One** 11 : e0156915, 2016
 12. Dunlop EA, Tee AR : The kinase triad, AMPK, mTORC1 and ULK1, maintains energy and nutrient homeostasis. **Biochem Soc Trans** 41 : 939-943, 2013
 13. Fu L, Zhang S, Zhang L, Tong X, Zhang J, Zhang Y, et al. : Systems biology network-based discovery of a small molecule activator BL-AD008 targeting AMPK/ZIPK and inducing apoptosis in cervical cancer. **Oncotarget** 6 : 8071-8088, 2015
 14. Ghinda DC, Wu JS, Duncan NW, Northoff G : How much is enough-can resting state fMRI provide a demarcation for neurosurgical resection in glioma? **Neurosci Biobehav Rev** 84 : 245-261, 2018
 15. Guerinéau M, Kriz Z, Kozakova L, Bednarova K, Janos P, Palecek J : Analysis of the Nse3/MAGE-binding domain of the Nse4/EID family proteins. **PLoS One** 7 : e35813, 2012
 16. He L, Zhou X, Huang N, Li H, Tian J, Li T, et al. : AMPK regulation of glucose, lipid and protein metabolism: mechanisms and nutritional significance. **Curr Protein Pept Sci** 18 : 562-570, 2017
 17. He S, Hu B, Li C, Lin P, Tang WG, Sun YF, et al. : PDXliver: a database of liver cancer patient derived xenograft mouse models. **BMC Cancer** 18 : 550, 2018
 18. Hombach-Klonisch S, Mehrpour M, Shojaei S, Harlos C, Pitz M, Hamai A, et al. : Glioblastoma and chemoresistance to alkylating agents: involvement of apoptosis, autophagy, and unfolded protein response. **Pharmacol Ther** 184 : 13-41, 2018
 19. Hsieh FS, Chen YL, Hung MH, Chu PY, Tsai MH, Chen LJ, et al. : Palbociclib induces activation of AMPK and inhibits hepatocellular carcinoma in a CDK4/6-independent manner. **Mol Oncol** 11 : 1035-1049, 2017
 20. Jeon SM : Regulation and function of AMPK in physiology and diseases. **Exp Mol Med** 48 : e245, 2016
 21. Jiang H, Liu W, Zhan SK, Pan YX, Bian LG, Sun B, et al. : GSK621 targets glioma cells via activating AMP-activated protein kinase signalings. **PLoS One** 11 : e0161017, 2016
 22. Kim J, Kundu M, Viollet B, Guan KL : AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. **Nat Cell Biol** 13 : 132-141, 2011
 23. Konagaya Y, Terai K, Hirao Y, Takakura K, Imajo M, Kamioka Y, et al. : A highly sensitive FRET biosensor for AMPK exhibits heterogeneous AMPK responses among cells and organs. **Cell Rep** 21 : 2628-2638, 2017
 24. Lapointe S, Perry A, Butowski NA : Primary brain tumours in adults. **Lancet** 392 : 432-446, 2018
 25. Liu X, Hu X, Kuang Y, Yan P, Li L, Li C, et al. : BCLB, methylated in hepatocellular carcinoma, is a starvation stress sensor that induces apoptosis and autophagy through the AMPK-mTOR signaling cascade. **Cancer Lett** 395 : 63-71, 2017
 26. Miller JJ, Shih HA, Andronesi OC, Cahill DP : Isocitrate dehydrogenase-mutant glioma: evolving clinical and therapeutic implications. **Cancer** 123 : 4535-4546, 2017
 27. Munakata K, Uemura M, Tanaka S, Kawai K, Kitahara T, Miyo M, et al. : Cancer stem-like properties in colorectal cancer cells with low proteasome activity. **Clin Cancer Res** 22 : 5277-5286, 2016
 28. Neguembor MV, Xynos A, Onorati MC, Caccia R, Bortolanza S, Godio C, et al. : FSHD muscular dystrophy region gene 1 binds suv4-20h1 histone methyltransferase and impairs myogenesis. **J Mol Cell Biol** 5 : 294-307, 2013
 29. O'Brien AJ, Villani LA, Broadfield LA, Houde VP, Galic S, Blandino G, et al. : Salicylate activates AMPK and synergizes with metformin to reduce the survival of prostate and lung cancer cells ex vivo through inhibition of de novo lipogenesis. **Biochem J** 469 : 177-187, 2015
 30. Pan SJ, Ren J, Jiang H, Liu W, Hu LY, Pan YX, et al. : MAGEA6 promotes human glioma cell survival via targeting AMPK α 1. **Cancer Lett** 412 : 21-29, 2018
 31. Pei S, Minhajuddin M, Adane B, Khan N, Stevens BM, Mack SC, et al. : AMPK/FIS1-mediated mitophagy is required for self-renewal of human AML stem cells. **Cell Stem Cell** 23 : 86-100, 2018
 32. Peng Z, Liu C, Wu M : New insights into long noncoding RNAs and their roles in glioma. **Mol Cancer** 17 : 61, 2018
 33. Quinones A, Le A : The multifaceted metabolism of glioblastoma. **Adv Exp Med Biol** 1063 : 59-72, 2018
 34. Sharples AP, Hughes DC, Deane CS, Saini A, Selman C, Stewart CE : Longevity and skeletal muscle mass: the role of IGF signalling, the sirtuins, dietary restriction and protein intake. **Aging Cell** 14 : 511-523, 2015
 35. Smith BK, Marcinko K, Desjardins EM, Lally JS, Ford RJ, Steinberg GR : Treatment of nonalcoholic fatty liver disease: role of AMPK. **Am J Physiol Endocrinol Metab** 311 : E730-E740, 2016
 36. Umezawa S, Higurashi T, Nakajima A : AMPK: therapeutic target for diabetes and cancer prevention. **Curr Pharm Des** 23 : 3629-3644, 2017
 37. Wang C, Tong Y, Wen Y, Cai J, Guo H, Huang L, et al. : Hepatocellular carcinoma-associated protein TD26 interacts and enhances sterol regulatory element-binding protein 1 activity to promote tumor cell proliferation and growth. **Hepatology** 68 : 1833-1850, 2018
 38. Weller M, Van den Bent M, Tonn JC, Stupp R, Preusser M, Cohen-Jonathan-Moyal E, et al. : European Association for Neuro-Oncology (EANO) guideline on the diagnosis and treatment of adult astrocytic and oligodendroglial gliomas. **Lancet Oncol** 18 : e315-e329, 2017
 39. Xu L, Kong L, Wang J, Ash JD : Stimulation of AMPK prevents degeneration of photoreceptors and the retinal pigment epithelium. **Proc Natl Acad Sci U S A** 115 : 10475-10480, 2018
 40. Zhao W, Peng F, Shu M, Liu H, Hou X, Wang X, et al. : Isogambogenic acid inhibits the growth of glioma through activation of the AMPK-mTOR pathway. **Cell Physiol Biochem** 44 : 1381-1395, 2017
 41. Zhao Y, Hu X, Liu Y, Dong S, Wen Z, He W, et al. : ROS signaling under metabolic stress: cross-talk between AMPK and AKT pathway. **Mol Cancer** 16 : 79, 2017