

# Novel Genome-Wide Interactions Mediated via *BOLL* and *EDNRA* Polymorphisms in Intracranial Aneurysm

Eun Pyo Hong,<sup>1</sup> Dong Hyuk Youn,<sup>1</sup> Bong Jun Kim,<sup>1</sup> Jae Jun Lee,<sup>1</sup> Sehyeon Nam,<sup>2</sup> Hyojong Yoo,<sup>2</sup> Heung Cheol Kim,<sup>3</sup> Jong Kook Rhim,<sup>4</sup> Jeong Jin Park,<sup>5</sup> Jin Pyeong Jeon<sup>6</sup>

*Institute of New Frontier Research,<sup>1</sup> Hallym University College of Medicine, Chuncheon, Korea*  
*Department of Materials Science and Chemical Engineering,<sup>2</sup> Hanyang University, Ansan, Korea*  
*Department of Radiology,<sup>3</sup> Hallym University College of Medicine, Chuncheon, Korea*  
*Department of Neurosurgery,<sup>4</sup> Jeju National University College of Medicine, Jeju, Korea*  
*Department of Neurology,<sup>5</sup> Konkuk University Medical Center, Seoul, Korea*  
*Department of Neurosurgery,<sup>6</sup> Hallym University College of Medicine, Chuncheon, Korea*

**Objective :** The association between *boule* (*BOLL*) and endothelin receptor type A (*EDNRA*) loci and intracranial aneurysm (IA) formation has been reported via genome-wide association studies. We sought to identify genome-wide interactions involving *BOLL* and *EDNRA* loci for IA in a Korean adult cohort.

**Methods :** Genome-wide pairwise interaction analyses of *BOLL* and *EDNRA* involving 250 patients with IA and 296 controls were performed using the additive effect model after adjusting for confounding factors.

**Results :** Among 512575 single-nucleotide polymorphisms (SNPs), 23 and 11 common SNPs suggested a genome-wide interaction threshold ( $p < 1.25 \times 10^{-8}$ ) involving rs700651 (*BOLL*) and rs6841581 (*EDNRA*). Rather than single SNP effect of *BOLL* or *EDNRA* on IA development, they showed a synergistic effect on IA formation via multifactorial pair-wise interactions. The rs1105980 of *PTCH1* gene showed the most significant interaction with rs700651 (natural log-transformed odds ratio [lnOR], 1.53;  $p = 6.41 \times 10^{-11}$ ). The rs74585958 of *RYK* gene interacted strongly with rs6841581 (lnOR, -19.91;  $p = 1.64 \times 10^{-9}$ ). Although, there was no direct interaction between *BOLL* and *EDNRA* variants, two *EDNRA*-interacting gene variants of *TNIK* (rs11925024 and rs1231) and *FTO* (rs9302654), and one *BOLL*-interacting *METTL4* gene variant (rs549315) exhibited marginal interaction with *BOLL* gene.

**Conclusion :** *BOLL* or *EDNRA* may have a synergistic effect on IA formation via multifactorial pair-wise interactions.

**Key Words :** *Boule* · Endothelin receptor type A · Genome-wide association study · Intracranial aneurysm.

## INTRODUCTION

Intracranial aneurysm (IA) refers to a bulge in the wall of in-

tracranial arteries due to endothelial dysfunction and extracellular matrix remodeling of the hemodynamic response. Although the prevalence of IA is approximately 3% in the general

• Received : February 8, 2022 • Revised : June 20, 2022 • Accepted : October 20, 2022

• Address for correspondence : Jin Pyeong Jeon

Department of Neurosurgery, Hallym University College of Medicine, 77 Sakju-ro, Chuncheon 24253, Korea

Tel : +82-33-240-5171, Fax : +82-33-240-9970, E-mail : jjs6553@daum.net, ORCID : https://orcid.org/0000-0001-8543-6855

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

population, the mortality rate due to subarachnoid hemorrhage following aneurysm rupture is close to 50%<sup>25,31</sup>. Clinical and radiological studies showed that IA formation and rupture were related to female gender, hypertension, smoking, larger size and posterior circulation aneurysm<sup>3</sup>.

The plethora of genome-wide association studies (GWASs) during late 2000's has increased the number of investigations into IA. These genetic studies have identified several candidate genes and loci associated with IA such as *EDNRA*, *GBA*, *CDKN2A/B*, *RBBP8*, *STARD13/KL*, and *SOX17*<sup>1,2,11,29</sup>. GWAS reported differences in the frequency of single-nucleotide polymorphisms (SNPs) based on case-control studies. Accordingly, the missing heritability is explained by independent SNPs involving complex diseases in human including cerebrovascular diseases (CVDs). To address this issue, robust analytical approaches such as meta-analyses, multifactorial interactions, and polygenic risk scoring systems have been performed in many populations<sup>9</sup>. A GWA meta-analysis reported IA-associated risk loci, including new loci (*SLC22A5*, *6q16.1*, *12q21.33*, *PSMA4*, and *NT5C2*) based on 10754 cases and 306882 controls<sup>11</sup>. Two large-scale meta-analyses also reported the successful validation of two IA-associated loci *BOLL* and endothelin receptor type A (*EDNRA*) in an East Asian population of nearly 20000 individuals<sup>12,13</sup>. However, few studies have reported gene-gene interactions or polygenic risk scores for IA patients. Considering to two previous findings of *BOLL* and *EDNRA* loci, here we estimated the effects of genome-wide gene-gene interactions on IA in a prospective hospital-based cohort study.

## MATERIALS AND METHODS

All the study protocols have been approved by the Institutional Review Board and Ethics Committee of Hallym University Chuncheon Sacred Heart Hospital (No. 2016-3, 2019-06-006). The study protocol and design are described in detail elsewhere<sup>11</sup>.

### Study populations

The study subjects were enrolled from the multi-institutional biobanks comprising five university hospitals constituting "The First Korean Stroke Genetics Association Research", and including patients diagnosed with CVDs such as IA be-

tween March 2015 and December 2020 (<https://www.1ksgh.org/>)<sup>11,18</sup>. Data derived from 250 patients with IA and 296 controls, which were also used in the first Korean IA GWAS<sup>11</sup>, were used in the analysis. The inclusion of patients with IA was based on the following criteria : 1) adult patients more than 18 years of age; 2) patients without other types of CVD such as ischemic stroke, hemorrhagic stroke, and vascular malformation; and 3) patients without any other genetic disorders such as polycystic kidney and moyamoya disease. Control subjects were defined as adults without CVD. Medical and radiological data were collected and updated.

### Genotyping and quality controls

Genomic DNA derived from the peripheral blood of the study population was genotyped using the Axiom<sup>TH</sup> Asian Precision Medicine Research Array (APMRA) (Thermo Fisher Scientific, Waltham, MA, USA). High quality plates were defined by a plate pass rate higher than 95% for samples and the average call rate of passing samples was greater than 99%. Out of 798148 SNPs, 512575 SNPs passed the quality control including genotyping call rate of 95% or higher, minor allele frequency (MAF) of at least 1%, and Hardy-Weinberg equilibrium (HWE) with  $p\text{-value} \geq 1 \times 10^{-6}$ <sup>11</sup>.

### Genome-wide SNP-SNP interactions via *BOLL* and *EDNRA* genes


We investigated genome-wide SNP-SNP interactions using either *BOLL* or *EDNRA* loci associated with IA in previous GWASs and meta-analyses<sup>11-13</sup>. The multivariate analyses of the SNP interactions between rs700651 intron SNP (*BOLL*, 2q33.1) and rs6841581 upstream SNP (*EDNRA*, 4q31.22) on 512574 SNPs were performed using the Contrived Acronym of software for SNP Interactions (CASSI ver. 2.5; <https://www.staff.ncl.ac.uk/richard.howey/cassi/index.html>)<sup>27</sup>. SNP-SNP interactions were analyzed by choosing target SNP from two given SNP windows (possibly from different pedigree files). Each pair of SNPs that interaction test passes a given significance level (i.e., minimized  $p\text{-value}=1$ ) is returned in the output file with possible extra information such as beta coefficient, standard error, chi-square, and  $p\text{-value}$ . The CASSI accepted only PLINK binary files in order to perform the calculations as efficiently as possible. The logistic regression epistasis test was available for SNP-SNP interactions in this study even though this program can provide the maximum number

of 1M terms. Subsequent regression analyses were carried out under the additive effect model and adjusted for 10 covariates including age, sex, hypertension, diabetes mellitus, hyperlipidemia, smoking status, and four genetic ancestry factors. The effect coefficient was estimated using the natural log-transformed scale of odds ratio (i.e., natural log-transformed odds ratio [lnOR]). The multiple comparisons were adjusted for a genome-wide interaction threshold with a  $p$ -value less than  $1.25 \times 10^{-8}$  (genome-wide  $p$ -value =  $5 \times 10^{-8}$  divided by four tests for interaction). A subsequent power and sample size calculations for each interaction term were estimated under the assumption with GW interaction significance threshold, 1 : 1.2 case-control ratio and information of each SNP (i.e., allele frequency and effect size) through performing the Quanto ver. 1.2.4 program (<https://bio.tools/QUANTO>). The performance of the large-scale interaction analyses was completed by the multi-tasking supper computer that has a capacities of Intel(R) Xeon(R) CPU E5-2667 v4 (3.20 GHz), 256 GB RAM, and 15 Cores. We performed Manhattan plots of *BOLL* and *EDNRA* interaction using the package of “*qqman*” in R v3.6.1 (<https://cran.r-project.org/web/packages/qqman>) and regional visualizations of the target SNP's base-pair position  $\pm 400$  kb regions using LocusZoom ver. 1.3 ([https://genome.sph.umich.edu/wiki/LocusZoom\\_Standalone](https://genome.sph.umich.edu/wiki/LocusZoom_Standalone)) written in Python and R<sup>23</sup>.

## RESULTS

Detailed information including SNP genotype distribution and HWE  $p$ -value, is presented in Supplementary Table 1. Out

of 512574 SNP interaction terms, 23 and 11 SNPs reached a genome-wide interaction threshold ( $p < 1.25 \times 10^{-8}$ ) with rs700651 intron SNP (*BOLL*, 2q33.1) and rs6841581 upstream SNP (*EDNRA*, 4q31.22), respectively (Fig. 1, Tables 1 and 2). All 34 SNPs showed an MAF above 1% and an HWE  $p$ -value greater than 0.01 (Supplementary Table 2). Most of the *BOLL*- or *EDNRA*-interacting SNPs showed shared alleles in both patient and control groups (i.e., average of MAF  $> 0.23$ ) without significant association in a single SNP analysis ( $0.0 < p < 1$ ) (Table 1). These findings suggest that *BOLL* or *EDNRA* may have a synergistic effect on IA formation, via multifactorial pair-wise interactions, rather than involved alone in the IA formation. Among the interactions, the rs1105980 upstream SNP of *PTCH1* gene (9q22.32) showed the most significant interaction with rs700651 (effect, 1.53;  $p = 6.41 \times 10^{-11}$ ) (Fig. 2A). The rs74585958 of *RYK* gene (3q22.2) interacted strongly with rs6841581 (effect, -19.91;  $p = 1.64 \times 10^{-9}$ ) (Fig. 2B). Two strong pair-wise linkage disequilibrium ( $r^2 > 0.95$ ) were observed in the interaction of rs328025 with rs700855 (*RGPD4*, 2q12.3) and between rs11925024 and rs1231 (*TNIK*, 3q26.31) ( $r^2 > 0.95$ , data not shown). Interestingly, two *EDNRA*-interacting gene variants of *TNIK* (rs11925024 and rs1231,  $p = 1.04 \times 10^{-8}$  and  $1.22 \times 10^{-9}$ , respectively) and *FTO* (rs9302654,  $p = 3.78 \times 10^{-9}$ ), and one *BOLL*-interacting *METTL4* gene variant (rs549315,  $p = 4.80 \times 10^{-10}$ ) showed marginal interaction with *BOLL* gene ( $0.001 < p < 0.05$ ). However, there was no direct interaction between *BOLL* and *EDNRA* variants (effect, -0.27;  $p = 0.301$ ). When power calculation was estimated by the basis of detail SNP information (Supplementary Tables 2 and 3), two SNPs such as rs74585958 (*RYK*) and rs150664966 (*EIF4H*) showed



**Fig. 1.** Manhattan plots of genome-wide interactions with (A) rs700651 (*BOLL*, 2q33.1) and (B) rs6841581 (*EDNRA*, 4q31.22) and their effect on intracranial aneurysm based on the additive effect model. Red line indicates genome-wide significance threshold (interaction  $p = 1.25 \times 10^{-8}$ ). SNP : single nucleotide polymorphism, ENDRA : endothelin receptor type A.

**Table 1.** Genome-wide interaction terms by *BOLL* or *EDNRA* polymorphisms on intracranial aneurysm according to *BOLL* interaction *p*-value ranking

Gene	Chr	SNP	BP	M/m	MAF, case/ control	lnOR*	<i>p</i> -value for GWAS*	lnOR <sup>†</sup>	<i>p</i> -value for interaction <sup>†</sup>
<i>BOLL</i>	2q33.1	rs700651	198631714	A/G	0.476/0.449	1.42	0.0079	NA	NA
<i>PTCH1</i>	9q22.32	rs1105980	98113635	G/C	0.27/0.294	-0.08	0.5821	1.53	6.41E-11
<i>CCDC3</i>	10p13	rs12412014	12911725	G/C	0.281/0.291	-0.09	0.5145	1.47	6.63E-11
<i>LINC00457</i>	13q13.2	rs1536847	35106975	G/T	0.295/0.329	-0.18	0.2033	1.39	2.37E-10
<i>C5orf60</i>	5q35.3	rs62405726	179069468	G/A	0.318/0.287	0.13	0.3646	1.46	4.21E-10
<i>METTL4</i>	18p11.32	rs549315	2183055	G/A	0.378/0.429	-0.29	0.0389	-1.32	4.80E-10
<i>RGPD4</i>	2q12.3	rs700855	108368694	T/C	0.372/0.328	0.27	0.0454	1.31	9.91E-10
<i>RGPD4</i>	2q12.3	rs328025	108355045	G/A	0.377/0.324	0.31	0.0219	1.25	1.06E-09
<i>MALL</i>	2q13	rs117802391	110862084	C/T	0.036/0.061	-0.77	0.0198	19.57	1.06E-09
<i>LINC01978</i>	17q25.3	rs57851800	77896371	A/C	0.345/0.326	0.16	0.2657	-1.27	1.08E-09
<i>RREB1</i>	6p24.3	rs9505086	7232186	T/C	0.286/0.307	-0.11	0.4279	1.24	1.20E-09
<i>DST</i>	6p12.1	rs117021265	56628021	T/C	0.034/0.024	0.25	0.5494	-18.96	1.75E-09
<i>RPRM</i>	2q23.3	rs5005908	154003680	G/T	0.344/0.27	0.32	0.022	1.26	3.03E-09
<i>FOXP1</i>	3p13	rs878118	71246228	T/G	0.238/0.255	-0.15	0.323	-1.39	3.85E-09
<i>LINC01344</i>	1q25.3	rs12033118	182229747	C/T	0.022/0.027	-0.15	0.7192	-19.75	5.00E-09
<i>RBMS3</i>	3p24.1	rs1979271	29607405	T/A	0.406/0.39	0.07	0.5841	-1.17	6.47E-09
<i>CXCR4</i>	2q22.1	rs189432614	136809235	A/G	0.016/0.025	-1.63	0.0131	-34.81	7.59E-09
<i>CDH13</i>	16q23.3	rs3848296	82550548	G/A	0.192/0.231	-0.16	0.3193	1.46	7.69E-09
<i>RUFY1</i>	5q35.3	rs4075890	178997373	T/C	0.216/0.2	0.1	0.5444	1.46	8.47E-09
<i>EIF2B5</i>	3q27.1	rs4350902	184352200	T/C	0.472/0.492	-0.09	0.5253	-1.18	8.83E-09
<i>PLEKHA1</i>	10q26.13	rs10510110	124192430	C/T	0.399/0.372	0.17	0.2098	1.26	8.96E-09
<i>PFKP</i>	10p15.2	rs58183624	3107217	C/T	0.066/0.044	0.15	0.592	-2.68	9.52E-09
<i>TRIM22</i>	11p15.4	rs7480654	5722839	T/C	0.317/0.284	0.18	0.22	-1.21	9.59E-09
<i>LINC00879</i>	3q11.2	rs4411883	94549686	T/G	0.09/0.111	-0.34	0.1404	-1.91	1.24E-08
<i>TNIF</i>	3q26.31	rs11925024	171014067	A/C	0.145/0.151	-0.15	0.4431	-0.9	0.0008
<i>TNIF</i>	3q26.31	rs1231	171031233	A/T	0.144/0.154	-0.18	0.3537	-0.8	0.0021
<i>FTO</i>	16q12.2	rs9302654	54009545	C/T	0.114/0.144	-0.34	0.088	0.58	0.0334
<i>SLFN11</i>	17q12	rs77814639	33678827	A/G	0.184/0.153	0.28	0.1329	-0.43	0.108
<i>SAP18</i>	13q12.11	rs9509543	21692404	C/T	0.346/0.356	-0.05	0.705	-0.2	0.2731
<i>EDNRA</i>	4q31.22	rs6841581	148401190	A/G	0.13/0.217	0.53	0.0006	-0.27	0.301
<i>SLC7A10</i>	19q13.11	rs11672303	33726375	T/C	0.171/0.154	0.08	0.6556	0.23	0.3238
<i>CACUL1</i>	10q26.11	rs11198727	120767097	A/G	0.382/0.429	-0.13	0.3308	0.17	0.3558
<i>MPDZ</i>	9p23	rs1332064	12942764	T/C	0.354/0.309	0.19	0.1724	0.1	0.5916
<i>UNC13C</i>	15q21.3	rs4774715	55140204	C/T	0.432/0.441	-0.01	0.9362	-0.09	0.6185
<i>RYK</i>	3q22.2	rs74585958	133773362	G/A	0.054/0.041	0.57	0.0817	-0.21	0.6852
<i>EIF4H</i>	7q11.23	rs150664966	73594157	T/C	0.016/0.022	-0.17	0.7152	-0.05	0.9384


\*These were estimated by generalized linear model after adjusting for age, sex, hypertension, diabetes, hyperlipidemia, and smoking in the previous GWAS. <sup>†</sup>These were estimated after *BOLL* by 500 K SNPs interactions by performing CASSI (Contrived Acronym of software for SNP Interactions) program after adjusting for age, sex, hypertension, diabetes, hyperlipidemia, and smoking. Chr : chromosome, SNP : single-nucleotide polymorphism, BP : base-pair position, M/m : major/minor allele type, MAF : minor allele frequency, lnOR : natural log-transformed odds ratio, GWAS : genome-wide association study, NA : not available

**Table 2.** Genome-wide interaction terms by *BOLL* or *EDNRA* polymorphisms on intracranial aneurysm according to *EDNRA* interaction *p*-value ranking

Gene	Chr	SNP	BP	M/m	MAF, case/ control	lnOR*	<i>p</i> -value for GWAS*	lnOR <sup>†</sup>	<i>p</i> -value for interaction <sup>†</sup>
<i>EDNRA</i>	4q31.22	rs6841581	148401190	A/G	0.13/0.217	0.53	0.0006	NA	NA
<i>RYK</i>	3q22.2	rs74585958	133773362	G/A	0.054/0.041	0.57	0.0817	-19.91	1.64E-09
<i>SAP18</i>	13q12.11	rs9509543	21692404	C/T	0.346/0.356	-0.05	0.705	1.85	2.87E-09
<i>SLC7A10</i>	19q13.11	rs11672303	33726375	T/C	0.171/0.154	0.08	0.6556	2.16	3.55E-09
<i>FTO</i>	16q12.2	rs9302654	54009545	C/T	0.114/0.144	-0.34	0.088	-3.1	3.78E-09
<i>SLFN11</i>	17q12	rs77814639	33678827	A/G	0.184/0.153	0.28	0.1329	-18.5	4.48E-09
<i>EIF4H</i>	7q11.23	rs150664966	73594157	T/C	0.016/0.022	-0.17	0.7152	20.91	4.80E-09
<i>MPDZ</i>	9p23	rs1332064	12942764	T/C	0.354/0.309	0.19	0.1724	1.75	5.10E-09
<i>UNC13C</i>	15q21.3	rs4774715	55140204	C/T	0.432/0.441	-0.01	0.9362	-1.68	7.74E-09
<i>CACUL1</i>	10q26.11	rs11198727	120767097	A/G	0.382/0.429	-0.13	0.3308	1.72	8.06E-09
<i>TNII</i>	3q26.31	rs11925024	171014067	A/C	0.145/0.151	-0.15	0.4431	-2.71	1.04E-08
<i>TNII</i>	3q26.31	rs1231	171031233	A/T	0.144/0.154	-0.18	0.3537	-2.86	1.22E-08
<i>METTL4</i>	18p11.32	rs549315	2183055	G/A	0.378/0.429	-0.29	0.0389	0.59	0.033
<i>PTCH1</i>	9q22.32	rs1105980	98113635	G/C	0.27/0.294	-0.08	0.5821	-0.6	0.058
<i>MALL</i>	2q13	rs117802391	110862084	C/T	0.036/0.061	-0.77	0.0198	-1.58	0.0762
<i>CXCR4</i>	2q22.1	rs189432614	136809235	A/G	0.016/0.025	-1.63	0.0131	1.02	0.2116
<i>LINC01344</i>	1q25.3	rs12033118	182229747	C/T	0.022/0.027	-0.15	0.7192	0.79	0.2679
<i>BOLL</i>	2q33.1	rs700651	198631714	A/G	0.476/0.449	1.42	0.0079	-0.27	0.301
<i>LINC01978</i>	17q25.3	rs57851800	77896371	A/C	0.345/0.326	0.16	0.2657	-0.29	0.3135
<i>PLEKHA1</i>	10q26.13	rs10510110	124192430	C/T	0.399/0.372	0.17	0.2098	-0.26	0.335
<i>PFKP</i>	10p15.2	rs58183624	3107217	C/T	0.066/0.044	0.15	0.592	0.56	0.3378
<i>LINC00457</i>	13q13.2	rs1536847	35106975	G/T	0.295/0.329	-0.18	0.2033	0.25	0.3692
<i>TRIM22</i>	11p15.4	rs7480654	5722839	T/C	0.317/0.284	0.18	0.22	-0.26	0.3718
<i>CDH13</i>	16q23.3	rs3848296	82550548	G/A	0.192/0.231	-0.16	0.3193	-0.31	0.3982
<i>FOXP1</i>	3p13	rs878118	71246228	T/G	0.238/0.255	-0.15	0.323	0.25	0.4035
<i>LINC00879</i>	3q11.2	rs4411883	94549686	T/G	0.09/0.111	-0.34	0.1404	-0.37	0.4182
<i>RGPD4</i>	2q12.3	rs328025	108355045	G/A	0.377/0.324	0.31	0.0219	-0.19	0.4897
<i>RUFY1</i>	5q35.3	rs4075890	178997373	T/C	0.216/0.2	0.1	0.5444	-0.21	0.5058
<i>RGPD4</i>	2q12.3	rs700855	108368694	T/C	0.372/0.328	0.27	0.0454	-0.18	0.5209
<i>CCDC3</i>	10p13	rs12412014	12911725	G/C	0.281/0.291	-0.09	0.5145	-0.14	0.6273
<i>RPRM</i>	2q23.3	rs5005908	154003680	G/T	0.344/0.27	0.32	0.022	-0.13	0.6388
<i>RBMS3</i>	3p24.1	rs1979271	29607405	T/A	0.406/0.39	0.07	0.5841	0.13	0.6541
<i>RREB1</i>	6p24.3	rs9505086	7232186	T/C	0.286/0.307	-0.11	0.4279	0.06	0.825
<i>DST</i>	6p12.1	rs117021265	56628021	T/C	0.034/0.024	0.25	0.5494	0.16	0.8283
<i>C5orf60</i>	5q35.3	rs62405726	179069468	G/A	0.318/0.287	0.13	0.3646	-0.02	0.9398
<i>EIF2B5</i>	3q27.1	rs4350902	184352200	T/C	0.472/0.492	-0.09	0.5253	-0.01	0.9793

\*These were estimated by generalized linear model after adjusting for age, sex, hypertension, diabetes, hyperlipidemia, and smoking in the previous GWAS. <sup>†</sup>These were estimated after *BOLL* by 500 K SNPs interactions by performing CASSI (Contrived Acronym of software for SNP Interactions) program after adjusting for age, sex, hypertension, diabetes, hyperlipidemia, and smoking. Chr : chromosome, SNP : single-nucleotide polymorphism, BP : base-pair position, M/m : major/minor allele type, MAF : minor allele frequency, lnOR : natural log-transformed odds ratio, GWAS : genome-wide association study, NA : not available





**Fig. 2.** Regional association plots of (A) rs1105980 (*PTCH1*, 9q22.32) interacting with rs700651 (*BOLL*, 2q33.1) and (B) rs74585958 (*RYK*, 3q22.2) interacting with rs6841581 (*EDNRA*, 4q31.22) at position  $\pm 400$  kb and the effect on intracranial aneurysm (IA). X-axis indicates the chromosomal position (mega base, Mb) and Y-axis did  $-\log_{10}$  transformed *p*-value and recombination rate, respectively. Purple triangles of rs1105980 ( $p=6.41 \times 10^{-11}$ ) and rs74585958 ( $p=6.41 \times 10^{-9}$ ) represent the most significant interactions with *BOLL* and *EDNRA*, respectively. Other up or down triangles denote other variants within the target variant  $\pm 400$  kb regions. Up and down triangles indicate positive and negative effect sizes on IA formation, respectively. Each color shows pair-wise linkage disequilibrium with either rs1105980 or rs74585958. *PTCH1*: patched 1, *RYK*: receptor-like tyrosine kinase.

sufficient statistical power among *EDNRA*-interacting loci (i.e., 0.885 and 0.805, respectively). The rs11672303 (*SLC7A10*) of *EDNRA*-interacting loci showed a marginal statistical power of 78.9%. In contrast, no SNP reached sufficient statistical power threshold of 80% in interaction terms by the *BOLL* locus (i.e., power <50%).

## DISCUSSION

Although *BOLL* gene, the G allele of rs700651 associated with risk showed a significant genome-wide overall effect on IA in multi-ethnic integrative meta-analyses ( $p=1.05 \times 10^{-8}$ )<sup>12</sup>, its pathogenic mechanism remains to be identified. Most studies related to *BOLL* involved spermatogenesis due to its role in germ cell development or cancer<sup>17,22</sup>. *BOLL* is a well-known gene associated with normal germ cell development<sup>26</sup>. The gene is predominantly expressed in secondary spermatocytes<sup>28</sup>. However, few studies investigated the role of *BOLL* in CVD including IA. Harrod et al.<sup>10</sup> reported that estrogen deficiencies may lead to IA by interrupting the inflammatory response. In reality, earlier age at menopause increased the risk of IA, suggesting the association between estrogen deficiency and IA pathogenesis<sup>6</sup>. Thus, in the case of *BOLL* gene, additional studies are needed to determine the protective effect of

male hormones against the IA development. The *EIF2B5* gene that interacts with *BOLL* in the current GW interaction and network analyses exhibited a homologous inhibition of cell translation. The differential expression of *EIF2B5* was moderate in human tissues and cells. Brady et al.<sup>4</sup> reported that intron retention in *EIF2B5* inhibited protein translation in hypoxic cancer cells. In the case of abdominal aortic aneurysms, there was no meaningful *EIF2B5* network<sup>16</sup>. Based on a review of the current literature, it is difficult to elucidate the contribution of the two genes to IA pathogenesis. Thus, a further *in vivo* study is needed to investigate the function of the two genes in IA formation.

The G allele of the rs6841581 located near the 5'-untranslated region of *EDNRA* (4q31.22) gene was associated with IA<sup>13</sup>. However, detailed mechanisms of IA mediated by *EDNRA* have yet to be reported to determine the direct effect or an indirect effect. Rats with pulmonary hypertension showed higher expression of *EDNRA* genes<sup>21</sup>. Endothelial injury, followed by disruption of collagen and elastin synthesis contribute to IA<sup>15</sup>. Chronic hypertension per se may induce the structural changes. Thus, the inflammatory response to increased hemodynamic stress following the disruption of cerebral arteries mediated by *EDNRA* may result in IA. In our study, among the *EDNRA*-interacting genes, the *MPDZ*-centered interaction between *EIF4H*, *SAP18*, and *UNC13C* was

observed. *MPDZ* is a tight junction protein, which modulates notch signaling during angiogenesis by controlling ligand recruitment to adherent junctions<sup>8,24</sup>. Feldner et al.<sup>8</sup> reported that loss of *MPDZ* decreased ependymal cell integrity and caused hydrocephalus. Ependymal cells are mainly responsible for electrolyte transport between brain parenchyma and the CSF. Adult ependymal cells are highly differentiated. Ependymal cells lining the lateral ventricles are quiescent under normal physiological conditions<sup>5</sup>. However, after stroke, adult ependymal cells are transformed into radial glial cells in the subventricular zone<sup>30</sup>. IA formation and growth occur within the CSF space surrounding the cerebral arteries. Accordingly, a further study is required to investigate the relationship between IA and CSF mediated by ependymal cells.

In our study, most variants did not show a significant association with IA via single SNP-based GWAS, although they exhibited significant associations with *BOLL* or *EDNRA* via multiple interactions terms. It is widely believed that a single SNP often has small effect on disease phenotypes including IA, thus it cannot fully account for the genetic susceptibility, in particular stroke. Therefore, identification of SNP interactions that are associated with disease is increasing to interpret the genetic basis of the disease susceptibility<sup>20</sup>. Although several loci related to *BOLL* or *EDNRA* did not pass genome-wide significance in a single SNP analysis, the loci might have a synergy effect on IA development by interacting with the genes. In this study, we aimed to investigate gene-gene interaction using the previous GWAS data for Korean patients with IA for the first time<sup>11</sup>. However, due to the relative small number of the enrolled patients and possible false positives<sup>14</sup>, we did not perform all possible pairwise SNP-SNP interaction and inevitably focused on two IA candidates of *BOLL* and *EDNRA* by referring to previous studies<sup>12,13</sup>. Nevertheless, we required further replication GWAS and exhaustive searching for SNP-SNP interaction in a large dataset of GWAS<sup>19</sup>. In addition, further molecular functional study including the estimation of protein levels to validate our interaction results between *BOLL* (or *EDNRA*) and several loci in the future.

The study has some limitations. First, although we identified novel loci for IA, their functional role was not investigated. The role of most of the 34 novel genes interacting with *BOLL* and *EDNRA* in our study has yet to be analyzed in IA or other CVDs. Second, the study had a potential sample size limitation of multiple interaction terms with either *BOLL* or

*EDNRA* loci, which have been replicated in IA susceptibility involving Korean adults including 250 patients with IA and 296 controls. In addition, there is a possibility that the results of this IA genetic study are limited to the Korean population. Therefore, we may warrant these findings in the second stage GWA meta-analysis and interaction test. Nevertheless, this study evaluates the first multiple genome-wide SNP-SNP interactions by IA-targeting genes such as *BOLL* and *EDNRA*, which have been validated in previous studies<sup>12,13</sup>. Disease is caused by various biological pathways and it is difficult to explain it based on a GWAS, which focused on differences between individual paired loci<sup>7</sup>. Accordingly, a study of genetic interactions and additional insights into various compensatory functional modules is needed to elucidate complex diseases such as IA. In summary, it is necessary to develop a general framework for mapping complex genetic networks of IA using GWAS data combined with clinically relevant risk factors.

## CONCLUSION

Genome-wide interaction between IA and *BOLL* or *EDNRA* revealed 34 novel loci, which were likely to be associated with IA. Common susceptibility variants and their interacting factors can be used to determine the inter-individual status of IA formation. The novel gene-gene interactions reported in this study need to be corroborated via larger prospective cohort studies.

## AUTHORS' DECLARATION

### Conflicts of interest

No potential conflict of interest relevant to this article was reported.

### Informed consent

This type of study does not require informed consent.

### Author contributions

Conceptualization : JPI, EPH; Data curation : JIL, SN, HCK, JKR, JJP; Formal analysis : EPH; Funding acquisition : JPI; Methodology : DHY, BJK, HY; Project administration : JPI; Visualization : EPH; Writing - original draft : JPI, EPH; Writ-

ing - review & editing : JPJ

## Data sharing

None

## Preprint

None

## ORCID

Eun Pyo Hong <https://orcid.org/0000-0001-7789-686X>  
 Dong Hyuk Youn <https://orcid.org/0000-0003-2259-1844>  
 Bong Jun Kim <https://orcid.org/0000-0003-3374-5554>  
 Jae Jun Lee <https://orcid.org/0000-0002-5418-500X>  
 Sehyeon Nam <https://orcid.org/0000-0001-6591-1022>  
 Hyojong Yoo <https://orcid.org/0000-0002-4932-3885>  
 Heung Cheol Kim <https://orcid.org/0000-0002-3573-1108>  
 Jong Kook Rhim <https://orcid.org/0000-0003-2538-8374>  
 Jeong Jin Park <https://orcid.org/0000-0001-9325-164X>  
 Jin Pyeong Jeon <https://orcid.org/0000-0001-8543-6855>

## • Acknowledgements

This research was supported by a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea (grant number : HR21C0198) and Hallym University Research Fund.

All genotype and phenotype resources are managed by “The First Korean Stroke Genetics Association Research” study constructed from the participating hospital stroke database.

## • Supplementary materials

The online-only data supplement is available with this article at <https://doi.org/10.3340/jkns.2022.0026>.

## References

- Bakker MK, van der Spek RAA, van Rheenen W, Morel S, Bourcier R, Hostettler IC, et al. : Genome-wide association study of intracranial aneurysms identifies 17 risk loci and genetic overlap with clinical risk factors. **Nat Genet** 52 : 1303-1313, 2020
- Bilguvar K, Yasuno K, Niemelä M, Ruigrok YM, von Und Zu Fraunberg M, van Duijn CM, et al. : Susceptibility loci for intracranial aneurysm in European and Japanese populations. **Nat Genet** 40 : 1472-1477, 2008
- Boehme AK, Esenwa C, Elkind MS : Stroke risk factors, genetics, and prevention. **Circ Res** 120 : 472-495, 2017
- Brady LK, Wang H, Radens CM, Bi Y, Radovich M, Maity A, et al. : Transcriptome analysis of hypoxic cancer cells uncovers intron retention in EIF2B5 as a mechanism to inhibit translation. **PLoS Biol** 15 : e2002623, 2017
- Carlén M, Meletis K, Göritz C, Darsalia V, Evergren E, Tanigaki K, et al. : Forebrain ependymal cells are notch-dependent and generate neuroblasts and astrocytes after stroke. **Nat Neurosci** 12 : 259-267, 2009
- Ding C, Toll V, Ouyang B, Chen M : Younger age of menopause in women with cerebral aneurysms. **J Neurointerv Surg** 5 : 327-331, 2013
- Fang G, Wang W, Paunic V, Heydari H, Costanzo M, Liu X, et al. : Discovering genetic interactions bridging pathways in genome-wide association studies. **Nat Commun** 10 : 4274, 2019
- Feldner A, Adam MG, Tetzlaff F, Moll I, Komljenovic D, Sahm F, et al. : Loss of Mpdz impairs ependymal cell integrity leading to perinatal-onset hydrocephalus in mice. **EMBO Mol Med** 9 : 890-905, 2017
- Génin E : Missing heritability of complex diseases: case solved? **Hum Genet** 139 : 103-113, 2020
- Harrod CG, Batjer HH, Bendok BR : Deficiencies in estrogen-mediated regulation of cerebrovascular homeostasis may contribute to an increased risk of cerebral aneurysm pathogenesis and rupture in menopausal and postmenopausal women. **Med Hypotheses** 66 : 736-756, 2006
- Hong EP, Kim BJ, Cho SS, Yang JS, Choi HJ, Kang SH, et al. : Genomic variations in susceptibility to intracranial aneurysm in the Korean population. **J Clin Med** 8 : 275, 2019
- Hong EP, Kim BJ, Jeon JP : Genome-wide association between the 2q33.1 locus and intracranial aneurysm susceptibility: an updated meta-analysis including 18,019 individuals. **J Clin Med** 8 : 692, 2019
- Hong EP, Kim BJ, Jeon JP, Yang JS, Choi HJ, Kang SH, et al. : Association of endothelin receptor type a with intracranial aneurysm in 20,609 East Asians: an updated meta-analysis. **World Neurosurg** 130 : e804-e814, 2019
- Hong EP, Youn DH, Kim BJ, Ahn JH, Park JJ, Rhim JK, et al. : Fine-mapping of intracranial aneurysm susceptibility based on a genome-wide association study. **Sci Rep** 12 : 2717, 2022
- Inci S, Spetzler RF : Intracranial aneurysms and arterial hypertension: a review and hypothesis. **Surg Neurol** 53 : 530-540; discussion 540-542, 2000
- Jones GT, Tromp G, Kuivaniemi H, Gretarsdottir S, Baas AF, Giusti B, et al. : Meta-analysis of genome-wide association studies for abdominal aortic aneurysm identifies four new disease-specific risk loci. **Circ Res** 120 : 341-353, 2017
- Kang KJ, Pyo JH, Ryu KJ, Kim SJ, Ha JM, Choi K, et al. : Oncogenic role of BOLL in colorectal cancer. **Dig Dis Sci** 60 : 1663-1673, 2015



18. Kim BJ, Kim Y, Youn DH, Park JJ, Rhim JK, Kim HC, et al. : Genome-wide blood DNA methylation analysis in patients with delayed cerebral ischemia after subarachnoid hemorrhage. **Sci Rep** **10** : 11419, 2020
19. Lee KY, Leung KS, Ma SL, So HC, Huang D, Tang NL, et al. : Genome-wide search for SNP interactions in GWAS data: algorithm, feasibility, replication using schizophrenia datasets. **Front Genet** **11** : 1003, 2020
20. Li P, Guo M, Wang C, Liu X, Zou Q : An overview of SNP interactions in genome-wide association studies. **Brief Funct Genomics** **14** : 143-155, 2015
21. Lim KA, Kim KC, Cho MS, Lee BE, Kim HS, Hong YM : Gene expression of endothelin-1 and endothelin receptor a on monocrotaline-induced pulmonary hypertension in rats after bosentan treatment. **Korean Circ J** **40** : 459-464, 2010
22. Luetjens CM, Xu EY, Rejo Pera RA, Kamischke A, Nieschlag E, Gromoll J : Association of meiotic arrest with lack of BOULE protein expression in infertile men. **J Clin Endocrinol Metab** **89** : 1926-1933, 2004
23. Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, et al. : LocusZoom: regional visualization of genome-wide association scan results. **Bioinformatics** **26** : 2336-2337, 2010
24. Tetzlaff F, Adam MG, Feldner A, Moll I, Menuchin A, Rodriguez-Vita J, et al. : MPDZ promotes DLL4-induced Notch signaling during angiogenesis. **Elife** **7** : e32860, 2018
25. Vlak MH, Algra A, Brandenburg R, Rinkel GJ : Prevalence of unruptured intracranial aneurysms, with emphasis on sex, age, comorbidity, country, and time period: a systematic review and meta-analysis. **Lancet Neurol** **10** : 626-636, 2011
26. Westerveld GH, Repping S, Leschot NJ, van der Veen F, Lombardi MP : Mutations in the human BOULE gene are not a major cause of impaired spermatogenesis. **Fertil Steril** **83** : 513-515, 2005
27. Wu X, Dong H, Luo L, Zhu Y, Peng G, Reveille JD, et al. : A novel statistic for genome-wide interaction analysis. **PLoS Genet** **6** : e1001131, 2010
28. Xu EY, Moore FL, Pera RA : A gene family required for human germ cell development evolved from an ancient meiotic gene conserved in metazoans. **Proc Natl Acad Sci U S A** **98** : 7414-7419, 2001
29. Yasuno K, Bilguvar K, Bijlenga P, Low SK, Krischek B, Auburger G, et al. : Genome-wide association study of intracranial aneurysm identifies three new risk loci. **Nat Genet** **42** : 420-425, 2010
30. Zhang RL, Zhang ZG, Wang Y, LeTourneau Y, Liu XS, Zhang X, et al. : Stroke induces ependymal cell transformation into radial glia in the subventricular zone of the adult rodent brain. **J Cereb Blood Flow Metab** **27** : 1201-1212, 2007
31. Zhou S, Dion PA, Rouleau GA : Genetics of intracranial aneurysms. **Stroke** **49** : 780-787, 2018