

Laboratory Investigation

Neuroprotective Effects of Quetiapine on Neuronal Apoptosis Following Experimental Transient Focal Cerebral Ischemia in Rats

Muhammet Bahadır Yılmaz, M.D.,¹ Mehmet Töngel, M.D.,² Hakan Emmez, M.D.,³ Figen Kaymaz, M.D.,⁴ Memduh Kaymaz, M.D.³

Department of Neurosurgery,¹ Ministry of Health, Training and Research Hospital, Kayseri, Turkey

Department of Neurosurgery,² Çankırı State Hospital, Çankırı, Turkey

Department of Neurosurgery,³ Gazi University Faculty of Medicine, Ankara, Turkey

Department of Histology and Embryology,⁴ Hacettepe University Faculty of Medicine, Ankara, Turkey

Objective : This study was undertaken in the belief that the atypical antipsychotic drug quetiapine could prevent apoptosis in the penumbra region following ischemia, taking into account findings that show 5-hydroxytryptamine-2 receptor blockers can prevent apoptosis.

Methods : We created 5 groups, each containing 6 animals. Nothing was done on the K-I group used for comparisons with the other groups to make sure adequate ischemia had been achieved. The K-II group was sacrificed on the 1st day after transient focal cerebral ischemia and the K-III group on the 3rd day. The D-I group was administered quetiapine following ischemia and sacrificed on the 1st day while the D-II group was administered quetiapine every day following the ischemia and sacrificed on the 3rd day. The samples were stained with the immunochemical TUNEL method and the number of apoptotic cells were counted.

Results : There was a significant difference between the first and third day control groups (K-II/K-III : $p=0.004$) and this indicates that apoptotic cell death increases with time. This increase was not encountered in the drug groups (D-I/D-II : $p=1.00$). Statistical analysis of immunohistochemical data revealed that quetiapine decreased the apoptotic cell death that normally increased with time.

Conclusion : Quetiapine is already in clinical use and is a safe drug, in contrast to many substances that are used to prevent ischemia and are not normally used clinically. Our results and the literature data indicate that quetiapine could help both as a neuronal protector and to resolve neuropsychiatric problems caused by the ischemia in cerebral ischemia cases.

Key Words : Quetiapine · Neuronal apoptosis · Neuroprotective · Transient focal cerebral ischemia.

INTRODUCTION

Stroke is the third most common cause of death globally after coronary heart disease and cancer. The incidence of stroke and the number of related deaths are increasing because of the increased percentage of the elderly in the population. Stroke also causes disability and significant socioeconomic damage in the population. It is the cause of neurological damage and many psychiatric and systemic complications such as depression and anxiety. The financial damage caused by decreased productivity in addition to the treatment costs of the disease and rehabilitation period are quite significant⁽²³⁾. The psychological damage

that develops after stroke is another aspect of the event and delays healing. Many drugs that show their effect through various mechanisms and various surgical methods have therefore been tried to prevent the ischemic damage caused by the primary ischemic or traumatic cerebrovascular events or the secondary injury. However, there is still no method or drug that can completely solve of this problem. Many of the drugs that have been tested are effective on one or several mechanisms but this chain of complicated and multi-factorial events cannot be broken and there is only a limited effect⁽⁵⁶⁾.

Ischemia-related apoptotic death and the penumbra area have recently become more popular subjects and are now the focal

• Received : March 10, 2013 • Revised : April 28, 2013 • Accepted : July 17, 2013

• Address for reprints : Muhammet Bahadır Yılmaz, M.D.

Sanayi Mah. Atatürk Bulvarı Hastane Cad. No:78, 38010. Ministry of Health, Training and Research Hospital, Department of Neurosurgery, Kocasinan, Kayseri, Turkey
Tel : +90-505-319-23-65, Fax : +90-352-320-73-13, E-mail : mbahadiryilmaz@yahoo.com.tr

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

point of many studies. Several articles emphasized the relationship between 5-hydroxytryptamine-2 (5-HT₂) agonist and antagonists and apoptotic cell death^{16,17,27,49,55,65}. We therefore planned our study believing that quetiapine, a dibenzothiazepine derivative atypical antipsychotic drug, could decrease apoptotic cell death and the negative effects of secondary injury in cerebral ischemia while also preventing the neuropsychiatric complications caused by this ischemia.

MATERIALS AND METHODS

This study was performed on a total of 30 rats in 5 experimental groups each containing 6 randomly selected rats obtained from the Gazi University Experimental Animals Research Center (GUDAM). The groups were as K-I, the control group where no surgical procedure was performed (n=6); K-II, the ischemia control group that lived for 1 day (n=6); K-III, the ischemia control group that lived for 3 days (n=6); D-I, the ischemia-medication group that lived for 1 day (n=6); and D-II, the ischemia-medication group that lived for 3 days (n=6). The K-I group was sacrificed to compare the apoptosis in the brain of healthy rats with those that had undergone ischemia. We performed the same surgical procedures in the other 4 groups to create ischemia and to sacrifice the animals. Taking the result of previous studies into account, the medication groups received quetiapine 10 mg/kg/day intraperitoneally 30 minutes after the trauma was caused, and then at the same time every day for 3 days^{24,51}.

Animals and ischemia surgery

Male Wistar type albino rats (n : 30) weighing 200-220 gr were kept in a room at 21±1°C temperature at 12 hours darkness and 12 hours light with 4 rates in a cage and ad lib water and feed.

General anesthesia was administered intramuscularly by using 10 mg/kg xylazine (Rompun, Bayer, Istanbul, Turkey) and 50 mg/kg ketamine hydrochloride (Ketalar, Eczacibasi, Istanbul, Turkey). Anesthesia depth was evaluated with a painful stimulus through the tail every 15 minutes. All surgical procedures were performed under the microscope (Opmi 99, Carl Zeiss, Germany).

Cerebral blood flow (CBF) monitorization was with a laser Doppler probe (Vasamedics, Blood Perfusion Monitor, Model BPM 433-1, St Paul, MN, USA) after a burr-hole was opened in the left middle fossa using a high-speed drill (Aesculap Microtron GD 412, Tuttlingen, Germany). After the 0 minute measurement, the left carotid artery bifurcation was dissected with a ventral midline neck incision for the MCA occlusion procedure. Transient focal cerebral ischemia was created using the MCA intraluminal filament method defined by Kawamura et al.³⁷ in 1991. We observed decreased blood flow on the CBF monitor when full occlusion was achieved. We accepted that the suture closed the ostium when the 0 minute CBF measurement had decreased at least 80%. All surgical procedures took

25 minutes on average. Reperfusion was achieved by removing the suture under anesthesia after 120 minutes of occlusion.

Animals in the K-II and D-I groups were sacrificed on the 24th hour of trauma. Animals in the K-III and D-II groups were sacrificed after being followed-up for 3 days. The K-I group, where no surgical intervention causing ischemia had taken place or medication had been given, was sacrificed to use as a comparison for apoptosis in healthy brain tissue.

Immunohistochemical study

All the histological and immunohistochemistry experimental studies were performed at the Hacettepe University Histology and Embryology Department laboratories.

The total brain tissue obtained from the rats was fixed with 10% buffer formaldehyde for 3 days. The dehydration and paraffinization procedures were performed with a tissue processing device (Leica TP 1020 CH, Germany). The paraffin embedding and block formation procedures were performed on the paraffin station (Leica EG 1150 CH, Germany).

Sections 5 µm thick were obtained from the paraffin blocks on special silanized slides. Following deparaffinization with heat for 1 hour once and chemical deparaffinization twice with xylol for 15 minutes, they were stained with hematoxylin-eosin and the accuracy of the field was checked. Other sections were marked with the immunohistochemical TUNEL method using the ApopTag Plus Peroxidase *In Situ* Apoptosis Detection Kit (Chemicon, USA, cat no : 7101). Background staining was provided by keeping the specimens in 0.5% methyl green for 10 minutes. Differentiation was with 100% n-butanol. We finally applied closure balsam and covered the material with a coverslip.

Evaluation of sections

The immunohistochemically stained sections of the group were evaluated on the light microscope (Leica CTR 6000, Germany). Nuclei stained with Methyl Green were evaluated as normal while cells with brown nuclear staining were considered ApopTag positive. Sections for all animals in each group were evaluated under 40× magnification. Five random areas that showed the most intensive ischemia were photographed (Leica DFC 490 camera, Germany) and the ApopTag positive cells were counted in each field. The apoptotic cells of the groups are shown in Fig. 1, 2.

Statistical methods

The Kruskal-Wallis test was used to compare the groups while the Dunn test was used to determine the group(s) that caused a difference during two-way comparisons. A *p* value <0.05 was considered statistically significant.

RESULTS

Table 1 presents the data on ApopTag positive cells stained

brown on a pale green background that were seen with the evaluation of 5 different randomly selected areas from the ischemia region for each animal in the groups (Table 1, Fig. 3).

Table 1 shows that the number of apoptotic cells increases significantly in the control group with time (in the K-III group compared with the K-II group). This time-related marked apoptotic cell increase is not seen in the medication group (between the D-I and D-II groups). The 1-day results show less apoptosis in the D-I group than the K-II group but with no statistical significance. The 3-day results show that the D-II group had markedly less apoptosis than the K-III group and that this is statistically significant.

Statistical analysis

Statistical analysis results are shown in Table 1. When all the groups were evaluated the p value was <0.001 indicating a significant difference between the groups for the number of apoptotic cells. The control group where no ischemic procedure was performed (K-I) was found to be statistically significantly different when compared with two-way comparisons with the other two groups that indicate that the method used for ischemia was successful ($p=0.002$). We did not find a significant difference when comparing the K-II, D-I, and D-II groups (D-I/D-II : $p=1.000$, K-II/D-I : $p=0.699$, K-II/D-II : $p=0.699$). The K-III group had the largest number of apoptotic cells and we found a statistically significant difference when we compared it with the other groups (K-III/D-I : $p=0.002$, K-III/D-II : 0.004 , K-II/K-III : $p=0.004$). There was a statistically significant difference between the K-II and K-III groups but none between the D-I and D-II groups indicating that the medication treatment was effective.

DISCUSSION

Stroke can lead to many neurological, psychiatric and systemic complications and has social and socioeconomic significance. It can develop as a result of ischemia, bleeding, or secondary injury. The central part of the ischemic area following stroke develops irreversible necrosis within minutes while the

Table 1. The number of apoptotic cells in 5 randomly selected fields from the ischemic area of each animal in the groups

	Mean	Std. Deviation	Median	Minimum	Maximum
K-I	1.17	0.41	1	1	2
K-II	72.33	40.32	55.5	40	140
K-III	156.17	31.44	154.5	109	203
D-I	60.00	23.26	59.5	35	91
D-II	63.50	37.40	50	36	137




Fig. 1. A : Neuron with brown marking following ApopTag staining in the K-II group. The black arrow shows an apoptotic cell sample (original magnification 40 \times). B : Neuron marked brown after ApopTag staining in the K-III group. The ApopTag (+) neurons are seen to be denser in the K-II group. The black arrow shows an apoptotic cell (original magnification 40 \times).




Fig. 2. A : Normal cells stained with methyl green and ApopTag (+) neurons stained brown in the D-I group (original magnification 40 \times). B : Normal cells stained with methyl green (blue arrow) are seen among the ApopTag (+) cells (black arrow) in the D-II group ischemic area (original magnification 40 \times). Significant apoptotic cell increase is not visible in D-I and D-II groups.

cell integrity is not yet structurally damaged in the penumbra area and the reestablishment of cerebral blood flow and use of neuronal protective agents can prevent ischemic injury in this area as the ion pumps are still healthy enough to continue their function, thus making this a target area. Energy-dependent apoptosis is seen especially in this penumbra area^{10,50,56,72}.

Apoptosis has first been used by the Scottish investigators Kerr et al.³⁸) in 1972 to mean “fall separately”. Apoptosis is currently defined as cell suicide and is a type of cell death where the organism can eliminate cells that have completed their biological function or are injured without damaging the surrounding cells, and it also plays a role in keeping the number of cells in tissues




Fig. 3. Graphical view of median apoptotic cell counts detected in each group.

under control. It is effective in both pathological and physiological events, is genetically controlled and consists of programmed biochemical and morphological changes. Apoptosis is an active function and requires high ATP levels^{7,15,26,32,36,40,44,50,56,58}.

Recent studies have selected the penumbra area that shows apoptosis as the target tissue that can be saved^{9,56}. Experimental studies have aimed to decrease or slow down this apoptotic process to gain time and ensure that some functions are regained with reperfusion. Despite this positive clinical picture and the possibility to recover some functions with reperfusion, the damage can also increase with the formation of free oxygen radicals^{21,56}.

Several recent studies with glutamate receptor antagonists^{20,43}, calcium channel blockers²⁵, free oxygen radical eliminating agents^{21,60}, citicholine^{1-4,56}, hypothermia^{5,47} and sometimes a combination of these agents have provided positive neuron protective results.

Experimental and clinical cerebral ischemia studies have shown that moderate hypothermia (2-6°C decrease) has a neuron protective effect by itself or when used in combination and is well tolerated^{55,56}. Although the mechanism of action has not been fully elucidated, it is thought to show its effect by decreasing the metabolic rate, protein synthesis and ATP usage, preventing the formation of free oxygen radicals and blocking the release of excitatory neurotransmitters^{41,42}. It has been reported that particularly positive results can be obtained with mutual reinforcement when it is used together with neuron protective substances^{5,56}.

Citicholine (CDP-choline), a necessary nucleoside for phospholipid synthesis and repair that play a role in the continuity in the cells membranes, has been shown to decrease the DNA fragmentation that appears following ischemia and the pro-caspase 1, 2, 3, 6 and 8 expression that plays a role in the apoptotic process^{2,4,47,56,59,67}. Some reports indicate that, when combined with hypothermia, it can prevent the apoptotic cell death stages without the creation of a apoptosome, and can decrease excitatory amino acid (EAA) accumulation and therefore neuronal damage that develops following a lack of energy^{45,56,59,69}.

An experimental study by Luo and Shi⁴⁶ has shown that polyethylene glycol can decrease apoptotic cell death following spinal trauma by suppressing cytochrome C release, one of the most important steps of apoptosis. The preventive effect against apoptosis in the posttraumatic spinal cord has been shown for 17-beta estradiol by Yune et al.⁷³, for erythropoietin by Arishima et al.⁸ and Vitellaro-Zuccarello et al.⁶⁴ and for antioxidants by Bao and Liu^{11,12}.

Gupta et al.³⁰, in their 2002 study, have shown that adenosine, and endogenous inhibitory neuromodulator in the central nervous system, decreases the release of EAA secretion during ischemia and protects the neurons against oxidative stress.

Caserta et al.¹⁹ have found that the irreversible general caspase inhibitor QVD-OPH prevents Caspase 8 and 10 activation related to the Fas/CD95 death receptors and Caspase 9 and 3 related to cytochrome C, thus preventing apoptosis in both neuronal and glial cells, decreasing secondary injury and contributing to functional recovery.

Studies by Muto et al.⁴⁹ in 2005 and Rajesh et al.⁵⁵ in 2006 have shown that Sarpogrelate, a 5-HT₂ receptor blocker, prevents the apoptotic death of cells in the cardiac tissue with its antiapoptotic effect and also shows a protective effect against myocardial dysfunction. Capela et al.^{16,17} have studied ecstasy (MDMA) and shown that it causes apoptotic cell death through the 5-HT_{2A} receptor pathways and that this apoptosis can be prevented using the 5-HT₂ blocker agent Ketanserine. Tascadda et al.⁶³ in 1999 and Tarazi et al.⁶² in 2003 have shown a neuroprotective effect of quetiapine through NMDA receptors with chronic usage.

We planned our study taking these articles on 5-HT₂ agonists and antagonists into account and used the dibenzothiazepine derivative atypical antipsychotic drug quetiapine believing that it may prevent apoptotic cell death and decrease the secondary damage caused by trauma in cerebral ischemia. The statistical results of our study show that comparison of K-I with the other groups has shown a significance of $p=0.002$ indicating that the ischemic procedure used to create apoptosis was successful. There was also a significant difference between the 1st day and 3rd day control groups (K-II/K-III : $p=0.004$), indicating increasing apoptotic cell death with time. Statistically, there was no increased apoptotic cell death with time in the groups administered medication (D-I/D-II : $p=1.00$). Our main purpose was to show whether our medication was effective and there was no statistically significant difference between the medication group on day 1 and the control group (K-II/D-I : $p=0.699$) while a statistical difference was present between the medication group on day 3 and the control group (K-III/D-II : 0.004). All these results have demonstrated immunohistochemically that quetiapine decreased time-related apoptotic cell death.

Yan et al.^{70,71} have reported that global cerebral ischemia causes hippocampal neurodegeneration, neuropsychiatric disorders and memory loss and that quetiapine can prevent these developments with its neuroprotective effect¹⁸. Administering quetiapine before ischemia has also been found to markedly

decrease depression and anxiety-type behavioral changes triggered with global cerebral ischemia in mice¹⁸). The use of quetiapine and olanzapine has been shown to have a significant effect on decreasing PC12 cell death and this cytoprotective effect is regulated through the translocation and expression of pro-apoptotic bax and antiapoptotic Bcl-X_L or through the increase of the antioxidant enzyme superoxide dismutase-1^{53,66,71}.

Another mechanism in ischemic events is the glutamate-related excitotoxicity. It is thought that ischemia and oxidative stress increase glutamate-related excitotoxicity and neuronal injury²⁸) and that quetiapine shows a neuroprotective effect by regulating glutamate receptor expression in the hippocampus and other cerebral regions and also decreases schizophrenia symptoms through this system^{54,62,63}).

Various studies have shown that atypical antipsychotic drugs including quetiapine prevent the apoptosis caused by N-methyl-4-phenylpyridinium ions⁵³) while another study has reported that they block the caspase-3 activation caused by beta-amyloid peptide and decrease apoptosis⁶³). Atypical antipsychotics including quetiapine have been shown to increase antiapoptotic Bcl-2 but Jarskog et al.³⁵) have found that subchronic antipsychotic usage increases caspase-3 activity without increasing DNA fragmentation and that it does not support Bcl-2 mediated neuroprotective efficacy despite what is currently believed.

Motor, sensory, visual, and speech disturbances are dominant among the clinical problems caused by microscopic cerebral ischemia. Some neuropsychiatric diseases, with depression being the most common, can be encountered in the early and late periods but may be missed or not emphasized due to the severe neurological picture of the patient. These conditions can also limit physical and cognitive recovery^{13,14,22,68,71}). From this point of view, apoptotic cell death, hippocampal neurodegeneration and effect of atypical antipsychotics on these disorders and especially schizophrenia have become popular study subjects in ischemia.

Most of the medications and methods used for neuroprotection and to prevent ischemia-related problems are not related to clinical use despite their beneficial effects. The quetiapine that we used in our study had marked efficacy especially when we look at our 3rd day results. It is also currently in use for schizophrenia, and unipolar^{18,48}) and bipolar depression^{15,18}) and is an effective medication with a better side effect profile than antipsychotics. It is also safe regarding extrapyramidal side effects^{6,33,52,61}). Clinical studies have revealed that quetiapine is also a suitable agent for the treatment of aggression and agitation due to traumatic cerebral damage³⁹). Quetiapine causes blockage with its high binding affinity for serotonin 5-HT₂ and 5-HT₆ receptors and the more moderate affinity for dopamine D₂ receptors. It also has affinity towards α_1 and α_2 receptors and H₁ receptors like clozapine. It shows negligible binding to dopamine D₁ and Muscarinic M₁ receptors^{33,34,57,61}). Although it possesses clozapine-like antipsychotic effects, its lower affinity for muscarinic and dopaminergic receptors ensures less anticholinergic and extrapyramidal

system side effects and less effect on prolactin levels^{29,31}). Some studies suggest a wider clinical use due to its many neuroprotective and neuropsychological effects as mentioned in our study¹⁸). Both our results and the fact that quetiapine is already used clinically also support a wider clinical perspective for this medication. However, further clinical studies are needed to support the efficacy of quetiapine in ischemia. Despite all these beneficial features, the US Food and Drug Administration has reported possible increased mortality with the use of atypical antipsychotic medications in the elderly with cardiovascular and infection problems in addition to the dementia-related psychosis¹⁸).

Studies mostly performed by psychiatry departments have emphasized many different mechanisms while studying the effect of ischemia and apoptosis in neuropsychiatric disorders and the mechanism of action of atypical antipsychotics in these disorders. In conclusion, quetiapine has been shown to decrease ischemia-related apoptosis in both our study and other studies.

CONCLUSION

Our results implicate that quetiapine which is emphasized effectiveness of prevention in neuropsychiatric diseases and enlarge its usage area by psychiatrists, also should be discussed by neurologists and neurosurgeons study on ischemia, be taken part in studies on clinical reflection of reducing apoptosis after ischemia, and remember its beneficial effect on overlooked ischemic neuropsychiatric problems.

References

1. Adibhatla RM, Hatcher JF : Cytidine 5'-diphosphocholine (CDP-choline) in stroke and other CNS disorders. *Neurochem Res* 30 : 15-23, 2005
2. Adibhatla RM, Hatcher JF, Dempsey RJ : Citicoline : neuroprotective mechanisms in cerebral ischemia. *J Neurochem* 80 : 12-23, 2002
3. Adibhatla RM, Hatcher JF, Dempsey RJ : Cytidine-5'-diphosphocholine affects CTP-phosphocholine cytidylyltransferase and lyso-phosphatidylcholine after transient brain ischemia. *J Neurosci Res* 76 : 390-396, 2004
4. Adibhatla RM, Hatcher JF, Dempsey RJ : Phospholipase A₂, hydroxyl radicals, and lipid peroxidation in transient cerebral ischemia. *Antioxid Redox Signal* 5 : 647-654, 2003
5. Alkan T, Kahveci N, Buyukuysal L, Korfali E, Ozluk K : Neuroprotective effects of MK 801 and hypothermia used alone and in combination in hypoxic-ischemic brain injury in neonatal rats. *Arch Physiol Biochem* 109 : 135-144, 2001
6. Altamura AC, Salvadori D, Madaro D, Santini A, Mundo E : Efficacy and tolerability of quetiapine in the treatment of bipolar disorder : preliminary evidence from a 12-month open-label study. *J Affect Disord* 76 : 267-271, 2003
7. Antonsson B : Bax and other pro-apoptotic Bcl-2 family "killer-proteins" and their victim the mitochondrion. *Cell Tissue Res* 306 : 347-361, 2001
8. Arishima Y, Setoguchi T, Yamaura I, Yone K, Komiya S : Preventive effect of erythropoietin on spinal cord cell apoptosis following acute traumatic injury in rats. *Spine (Phila Pa 1976)* 31 : 2432-2438, 2006
9. Back T : Pathophysiology of the ischemic penumbra--revision of a con-

- cept. *Cell Mol Neurobiol* 18 : 621-638, 1998
10. Balla A, Tóth B, Timár G, Bak J, Krajcsi P : Molecular targets for pharmacological cytoprotection. *Biochem Pharmacol* 61 : 769-777, 2001
 11. Bao F, Liu D : Hydroxyl radicals generated in the rat spinal cord at the level produced by impact injury induce cell death by necrosis and apoptosis : protection by a metalloporphyrin. *Neuroscience* 126 : 285-295, 2004
 12. Bao F, Liu D : Peroxynitrite generated in the rat spinal cord induces apoptotic cell death and activates caspase-3. *Neuroscience* 116 : 59-70, 2003
 13. Bokura H, Robinson RG : Long-term cognitive impairment associated with caudate stroke. *Stroke* 28 : 970-975, 1997
 14. Brådvik B, Sonesson B, Holtås S : Spatial impairment following right hemisphere transient ischaemic attacks in patients without carotid artery stenosis. *Acta Neurol Scand* 80 : 411-418, 1989
 15. Calabrese JR, Keck PE Jr, Macfadden W, Minkwitz M, Ketter TA, Weisler RH, et al. : A randomized, double-blind, placebo-controlled trial of quetiapine in the treatment of bipolar I or II depression. *Am J Psychiatry* 162 : 1351-1360, 2005
 16. Capela JP, Fernandes E, Remião F, Bastos ML, Meisel A, Carvalho F : Ecstasy induces apoptosis via 5-HT(2A)-receptor stimulation in cortical neurons. *Neurotoxicology* 28 : 868-875, 2007
 17. Capela JP, Ruscher K, Lautenschlager M, Freyer D, Dirnagl U, Gaio AR, et al. : Ecstasy-induced cell death in cortical neuronal cultures is serotonin 2A-receptor-dependent and potentiated under hyperthermia. *Neuroscience* 139 : 1069-1081, 2006
 18. Carta MG, Zairo F, Mellino G, Hardoy MC : Add-on quetiapine in the treatment of major depressive disorder in elderly patients with cerebrovascular damage. *Clin Pract Epidemiol Ment Health* 3 : 28, 2007
 19. Caserta TM, Smith AN, Gultice AD, Reedy MA, Brown TL : Q-VD-OPh, a broad spectrum caspase inhibitor with potent antiapoptotic properties. *Apoptosis* 8 : 345-352, 2003
 20. Childs AM, Evans RH, Watkins JC : The pharmacological selectivity of three NMDA antagonists. *Eur J Pharmacol* 145 : 81-86, 1988
 21. Cho S, Park EM, Febbraio M, Anrather J, Park L, Racchumi G, et al. : The class B scavenger receptor CD36 mediates free radical production and tissue injury in cerebral ischemia. *J Neurosci* 25 : 2504-2512, 2005
 22. Desmond DW, Moroney JT, Sano M, Stern Y : Mortality in patients with dementia after ischemic stroke. *Neurology* 59 : 537-543, 2002
 23. Easton JD : Current advances in the management of stroke. *Neurology* 51 (3 Suppl 3) : S1-S2, 1998
 24. Ellenbroek BA, Lubbers LJ, Cools AR : Activity of "seroquel" (ICI 204,636) in animal models for atypical properties of antipsychotics : a comparison with clozapine. *Neuropsychopharmacology* 15 : 406-416, 1996
 25. Franke CL, Palm R, Dalby M, Schoonderwaldt HC, Hantson L, Eriksson B, et al. : Flunarizine in stroke treatment (FIST) : a double-blind, placebo-controlled trial in Scandinavia and the Netherlands. *Acta Neurol Scand* 93 : 56-60, 1996
 26. Friedlander RM : Apoptosis and caspases in neurodegenerative diseases. *N Engl J Med* 348 : 1365-1375, 2003
 27. Ginsberg MD : Adventures in the pathophysiology of brain ischemia : penumbra, gene expression, neuroprotection : the 2002 Thomas Willis Lecture. *Stroke* 34 : 214-223, 2003
 28. Green AR, Hainsworth AH, Jackson DM : GABA potentiation : a logical pharmacological approach for the treatment of acute ischaemic stroke. *Neuropharmacology* 39 : 1483-1494, 2000
 29. Gunasekara NS, Spencer CM : Quetiapine : a review of its use in schizophrenia. *CNS Drugs* 9 : 325-340, 1998
 30. Gupta YK, Sinha K, Chaudhary G, Jagannathan NR : Protective effect of adenosine against neuronal injury induced by middle cerebral artery occlusion in rats as evidenced by diffusion-weighted imaging. *Pharmacol Biochem Behav* 72 : 569-574, 2002
 31. Hamner MB, Arvanitis LA, Miller BG, Link CG, Hong WW : Plasma prolactin in schizophrenia subjects treated with Seroquel (ICI 204,636). *Psychopharmacol Bull* 32 : 107-110, 1996
 32. Hengartner MO : The biochemistry of apoptosis. *Nature* 407 : 770-776, 2000
 33. Hirsch SR, Link CG, Goldstein JM, Arvanitis LA : ICI 204,636 : a new atypical antipsychotic drug. *Br J Psychiatry Suppl* : 45-56, 1996
 34. Horacek J, Bubenikova-Valesova V, Kopecek M, Palenicek T, Dockery C, Mohr P, et al. : Mechanism of action of atypical antipsychotic drugs and the neurobiology of schizophrenia. *CNS Drugs* 20 : 389-409, 2006
 35. Jarskog LE, Gilmore JH, Glantz LA, Gable KL, German TT, Tong RI, et al. : Caspase-3 activation in rat frontal cortex following treatment with typical and atypical antipsychotics. *Neuropsychopharmacology* 32 : 95-102, 2007
 36. Kanduc D, Mittelman A, Serpico R, Sinigaglia E, Sinha AA, Natale C, et al. : Cell death : apoptosis versus necrosis (review). *Int J Oncol* 21 : 165-170, 2002
 37. Kawamura S, Yasui N, Shirasawa M, Fukasawa H : Rat middle cerebral artery occlusion using an intraluminal thread technique. *Acta Neurochir (Wien)* 109 : 126-132, 1991
 38. Kerr JF, Wyllie AH, Currie AR : Apoptosis : a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br J Cancer* 26 : 239-257, 1972
 39. Kim E, Bijlani M : A pilot study of quetiapine treatment of aggression due to traumatic brain injury. *J Neuropsychiatry Clin Neurosci* 18 : 547-549, 2006
 40. Korenkov AI, Pahnke J, Frei K, Warzok R, Schroeder HW, Frick R, et al. : Treatment with nimodipine or mannitol reduces programmed cell death and infarct size following focal cerebral ischemia. *Neurosurg Rev* 23 : 145-150, 2000
 41. Krieger DW, Yenari MA : Therapeutic hypothermia for acute ischemic stroke : what do laboratory studies teach us? *Stroke* 35 : 1482-1489, 2004
 42. Krupinski J, Ferrer I, Barrachina M, Secades JJ, Mercadal J, Lozano R : CDP-choline reduces pro-caspase and cleaved caspase-3 expression, nuclear DNA fragmentation, and specific PARP-cleaved products of caspase activation following middle cerebral artery occlusion in the rat. *Neuropharmacology* 42 : 846-854, 2002
 43. LeBlanc MH, Li XQ, Huang M, Patel DM, Smith EE : AMPA antagonist LY293558 does not affect the severity of hypoxic-ischemic injury in newborn pigs. *Stroke* 26 : 1908-1914; discussion 1914-1915, 1995
 44. Liu X, Kim CN, Yang J, Jemmerson R, Wang X : Induction of apoptotic program in cell-free extracts : requirement for dATP and cytochrome c. *Cell* 86 : 147-157, 1996
 45. Love S : Apoptosis and brain ischaemia. *Prog Neuropsychopharmacol Biol Psychiatry* 27 : 267-282, 2003
 46. Luo J, Shi R : Polyethylene glycol inhibits apoptotic cell death following traumatic spinal cord injury. *Brain Res* 1155 : 10-16, 2007
 47. Maier CM, Ahern Kv, Cheng ML, Lee JE, Yenari MA, Steinberg GK : Optimal depth and duration of mild hypothermia in a focal model of transient cerebral ischemia : effects on neurologic outcome, infarct size, apoptosis, and inflammation. *Stroke* 29 : 2171-2180, 1998
 48. McIntyre A, Gendron A, McIntyre A : Quetiapine reduces residual depressive and prominent anxiety symptoms in partial responders to selective serotonin reuptake inhibitors (SSRIs) or serotonin norepinephrine reuptake inhibitors (SNRIs) with major depression : an 8-week, double-blind, randomized, placebo-controlled study. *European Psychiatry* 21 Suppl 1 : 164, 2006
 49. Muto T, Hotta Y, Miyazeki K, Ando H, Ishikawa N, Hasegawa T, et al. : Protective effects of sarpogrelate, a 5-HT2A antagonist, against post-ischemic myocardial dysfunction in guinea-pig hearts. *Mol Cell Bio-*

- chem 272 : 119-132, 2005
50. Onténiente B, Couriaud C, Braudeau J, Benchoua A, Guégan C : The mechanisms of cell death in focal cerebral ischemia highlight neuroprotective perspectives by anti-caspase therapy. *Biochem Pharmacol* 66 : 1643-1649, 2003
 51. Orsetti M, Canonico PL, Dellarole A, Colella L, Di Brisco F, Ghi P : Quetiapine prevents anhedonia induced by acute or chronic stress. *Neuropsychopharmacology* 32 : 1783-1790, 2007
 52. Peuskens J, Link CG : A comparison of quetiapine and chlorpromazine in the treatment of schizophrenia. *Acta Psychiatr Scand* 96 : 265-273, 1997
 53. Qing H, Xu H, Wei Z, Gibson K, Li XM : The ability of atypical antipsychotic drugs vs. haloperidol to protect PC12 cells against MPP+-induced apoptosis. *Eur J Neurosci* 17 : 1563-1570, 2003
 54. Raggi MA, Mandrioli R, Sabbioni C, Pucci V : Atypical antipsychotics : pharmacokinetics, therapeutic drug monitoring and pharmacological interactions. *Curr Med Chem* 11 : 279-296, 2004
 55. Rajesh KG, Suzuki R, Maeda H, Murio Y, Sasaguri S : 5-HT₂ receptor blocker sarpogrelate prevents downregulation of antiapoptotic protein Bcl-2 and protects the heart against ischemia-reperfusion injury. *Life Sci* 79 : 1749-1755, 2006
 56. Sahin S, Alkan T, Temel SG, Tureyen K, Tolunay S, Korfali E : Effects of citicoline used alone and in combination with mild hypothermia on apoptosis induced by focal cerebral ischemia in rats. *J Clin Neurosci* 17 : 227-231, 2010
 57. Saller CF, Salama AI : Seroquel : biochemical profile of a potential atypical antipsychotic. *Psychopharmacology (Berl)* 112 : 285-292, 1993
 58. Saraste A, Pulkki K : Morphologic and biochemical hallmarks of apoptosis. *Cardiovasc Res* 45 : 528-537, 2000
 59. Sastry PS, Rao KS : Apoptosis and the nervous system. *J Neurochem* 74 : 1-20, 2000
 60. Schmidley JW : Free radicals in central nervous system ischemia. *Stroke* 21 : 1086-1090, 1990
 61. Small JG, Hirsch SR, Arvanitis LA, Miller BG, Link CG : Quetiapine in patients with schizophrenia. A high- and low-dose double-blind comparison with placebo. Seroquel Study Group. *Arch Gen Psychiatry* 54 : 549-557, 1997
 62. Tarazi FI, Baldessarini RJ, Kula NS, Zhang K : Long-term effects of olanzapine, risperidone, and quetiapine on ionotropic glutamate receptor types : implications for antipsychotic drug treatment. *J Pharmacol Exp Ther* 306 : 1145-1151, 2003
 63. Tascetta F, Lovati E, Blom JM, Muzzioli P, Brunello N, Racagni G, et al. : Regulation of ionotropic glutamate receptors in the rat brain in response to the atypical antipsychotic seroquel (quetiapine fumarate). *Neuropsychopharmacology* 21 : 211-217, 1999
 64. Vitellaro-Zuccarello L, Mazzetti S, Madaschi L, Bosisio P, Gorio A, De Biasi S : Erythropoietin-mediated preservation of the white matter in rat spinal cord injury. *Neuroscience* 144 : 865-877, 2007
 65. Wang H, Xu H, Dyck LE, Li XM : Olanzapine and quetiapine protect PC12 cells from beta-amyloid peptide(25-35)-induced oxidative stress and the ensuing apoptosis. *J Neurosci Res* 81 : 572-580, 2005
 66. Wei Z, Mousseau DD, Richardson JS, Dyck LE, Li XM : Atypical antipsychotics attenuate neurotoxicity of beta-amyloid(25-35) by modulating Bax and Bcl-X(l/s) expression and localization. *J Neurosci Res* 74 : 942-947, 2003
 67. Weiss GB : Metabolism and actions of CDP-choline as an endogenous compound and administered exogenously as citicoline. *Life Sci* 56 : 637-660, 1995
 68. Williams LS, Ghose SS, Swindle RW : Depression and other mental health diagnoses increase mortality risk after ischemic stroke. *Am J Psychiatry* 161 : 1090-1095, 2004
 69. Winfree CJ, Baker CJ, Connolly ES Jr, Fiore AJ, Solomon RA : Mild hypothermia reduces penumbral glutamate levels in the rat permanent focal cerebral ischemia model. *Neurosurgery* 38 : 1216-1222, 1996
 70. Yan B, Bi X, He J, Zhang Y, Thakur S, Xu H, et al. : Quetiapine attenuates spatial memory impairment and hippocampal neurodegeneration induced by bilateral common carotid artery occlusion in mice. *Life Sci* 81 : 353-361, 2007
 71. Yan B, He J, Xu H, Zhang Y, Bi X, Thakur S, et al. : Quetiapine attenuates the depressive and anxiolytic-like behavioural changes induced by global cerebral ischemia in mice. *Behav Brain Res* 182 : 36-41, 2007
 72. Yu L, Miao H, Hou Y, Zhang B, Guo L : Neuroprotective effect of A20 on TNF-induced postischemic apoptosis. *Neurochem Res* 31 : 21-32, 2006
 73. Yune TY, Kim SJ, Lee SM, Lee YK, Oh YJ, Kim YC, et al. : Systemic administration of 17beta-estradiol reduces apoptotic cell death and improves functional recovery following traumatic spinal cord injury in rats. *J Neurotrauma* 21 : 293-306, 2004