

Erratum



Erratum: Correction of Text in the Article “Seroprevalence of Hepatitis C Virus Infection in North Korean Defectors Residing in Korea”

Young Mi Hong ^{1,2} Ki Tae Yoon ^{1,2} Young Joo Park ^{1,3} Hyun Young Woo ^{1,3}
and Jeong Heo ^{1,3}

¹Department of Internal Medicine, College of Medicine, Pusan National University, Busan, Korea

²Department of Internal Medicine, Pusan National University Yangsan Hospital, Yangsan, Korea

³Biomedical Research Institute, Pusan National University Hospital, Busan, Korea

OPEN ACCESS

► This corrects the article “Seroprevalence of Hepatitis C Virus Infection in North Korean Defectors Residing in Korea” in volume 38, number 34, e270.

Published online: Sep 21, 2023

Address for Correspondence:

Jeong Heo, MD, PhD

Department of Internal Medicine, College of Medicine, Pusan National University and Biomedical Research Institute, Pusan National University Hospital, 179 Gudeok-ro, Seo-gu, Busan 49241, Republic of Korea.
Email: jheo@pusan.ac.kr

© 2023 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Young Mi Hong 

<https://orcid.org/0000-0001-5427-7505>

Ki Tae Yoon 

<https://orcid.org/0000-0002-8580-0239>

Young Joo Park 

<https://orcid.org/0000-0001-8416-0615>

Hyun Young Woo 

<https://orcid.org/0000-0002-0605-6318>

Jeong Heo 

<https://orcid.org/0000-0003-0961-7851>

To the Editor:

The authors regret that there were errors in text. This notice corrects the percentage of participants who were unaware of HCV disease.

Content of correction:

<Before>

Of the participants, 24.6% were unaware of HCV disease, and only 23.7% thought that the treatment was necessary for HCV infection before participating in the study.

<After>

Of the participants, 75.4% were unaware of HCV disease, and only 23.7% thought that the treatment was necessary for HCV infection before participating in the study.