

Original Article

Carbon monoxide releasing molecule-2 suppresses stretch-activated atrial natriuretic peptide secretion by activating large-conductance calcium-activated potassium channels

Weijian Li^{1,#}, Sun Hwa Lee^{2,3,#}, and Sun Hee Kim^{1,3,*}

¹Departments of Physiology and ²Internal Medicine, Jeonbuk National University Medical School, ³Research Institute of Clinical Medicine of Jeonbuk National University Jeonju 54907, Korea

ARTICLE INFO

Received January 3, 2022
Revised January 18, 2022
Accepted January 18, 2022

*Correspondence

Suhn Hee Kim
E-mail: shkim@jbnu.ac.kr

Key Words

ANP
BK_{Ca} channel
Carbon monoxide
CORM-2
Hypoxia
K_{ATP} channel

#These authors contributed equally to this work.

ABSTRACT Carbon monoxide (CO) is a known gaseous bioactive substance found across a wide array of body systems. The administration of low concentrations of CO has been found to exert an anti-inflammatory, anti-apoptotic, anti-hypertensive, and vaso-dilatory effect. To date, however, it has remained unknown whether CO influences atrial natriuretic peptide (ANP) secretion. This study explores the effect of CO on ANP secretion and its associated signaling pathway using isolated beating rat atria. Atrial perfusate was collected for 10 min for use as a control, after which high atrial stretch was induced by increasing the height of the outflow catheter. Carbon monoxide releasing molecule-2 (CORM-2; 10, 50, 100 μM) and hemin (HO-1 inducer; 0.1, 1, 50 μM), but not CORM-3 (10, 50, 100 μM), decreased high stretch-induced ANP secretion. However, zinc porphyrin (HO-1 inhibitor) did not affect ANP secretion. The order of potency for the suppression of ANP secretion was found to be hemin > CORM-2 >> CORM-3. The suppression of ANP secretion by CORM-2 was attenuated by pretreatment with 5-hydroxydecanoic acid, paxilline, and 1*H*-[1,2,4] oxadiazolo [4,3-*a*] quinoxalin-1-one, but not by diltiazem, wortmannin, LY-294002, or NG-nitro-L-arginine methyl ester. Hypoxic conditions attenuated the suppressive effect of CORM-2 on ANP secretion. In sum, these results suggest that CORM-2 suppresses ANP secretion via mitochondrial K_{ATP} channels and large conductance Ca²⁺-activated K⁺ channels.

INTRODUCTION

Carbon monoxide (CO), an important gasotransmitter endogenously produced via the degradation of heme by heme-oxygenases (HOs) [1], possesses similar biological and pharmacological properties to nitric oxide (NO) and hydrogen sulfide [2-4]. As active signaling molecules involved in a variety of physiological and pathological processes, these gasotransmitters have been the focus of increasing attention over the past decade [2-6]. CO has been previously regarded only as a highly toxic gas for its role in preventing O₂ transport and delivery [7], and involvement in the

inducement of cardiac dysfunction [8-10]. However, CO is now known to be a neurotransmitter, as well as a cardiovascular and immune regulator [2,5]. Numerous studies have shown that low concentrations of CO exhibit pleiotropic biological functions, including anti-apoptotic, cytoprotective, and anti-inflammatory effects [11-13]. Cardiovascular tissues in normal conditions in particular express very low levels of HO-1 (an inducible form of HO), which is upregulated following stress symptoms such as high temperature and ischemia/reperfusion (I/R) or in certain pathological conditions [4,14-16]. The HO/CO system has been reported as being responsible for the numerous positive effects in



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.
Copyright © Korean J Physiol Pharmacol, pISSN 1226-4512, eISSN 2093-3827

Author contributions: W.L. performed the experiments, analyzed the data, wrote the original draft. S.H.L. wrote the review & editing. S.H.K. supervised and wrote the review & editing.

studies of cardiovascular [13] and lung disorders [6,17].

The cardiac hormone atrial natriuretic peptide (ANP) is released primarily from the cardiac atria and plays an important role in various physiological and pathological processes [18,19]. ANP decreases blood pressure by regulating extracellular fluid (ECF) volume and vasodilation, while exerting numerous beneficial effects on cardiovascular disorders and metabolic syndromes [20]. In addition, ANP levels serve as a potential biomarker for the diagnosis of the acute stages of heart failure. The factors that determine the amount of circulating ANP are intravascular volume change [21,22], solute concentration, and presence of several hormones [23,24]. Therefore, increased plasma ANP levels are an important trigger for the compensatory mechanisms of the cardiovascular system that respond to physiological and pathological conditions. The effects of CO vary across tissues, but involve the modulation of multiple pathways that contain NO, reactive oxygen species, and P38 MAP kinases [4]. What remains unknown are the details of the relationship between CO's beneficial effects and ANP.

Several boron- or ruthenium-based compounds (carbon monoxide releasing molecules, CORMs) have been synthesized in the last decade. These exhibit pharmaceutical traits superior to those of gaseous CO by virtue of their chemistry and structure [25,26]. We performed this study using CORM and isolated perfused beating rat atria to determine the direct effect of CO on ANP secretion and the underlying signaling mechanisms involved.

METHODS

Animals

Male Sprague–Dawley rats (Dae Han Bio Link, Eumsung, Korea), weighing 200–250 g each, were housed in temperature-controlled rooms with a 12 h light–dark cycle. Animals were given free access to standard laboratory chow (5L79 Purina rat and mouse 18% chow; Charles River Laboratories Inc., Wilmington, MA, USA) and water. All experimental protocols conformed to the guidelines established by the National Institutes of Health Guide for the Care and Use of Laboratory Animals (NIH publication number 85-23, revised 1996) and were approved by our institution.

Chemicals

ANP was purchased from Bachem Holding AG (Bubendorf, Switzerland). CORM-2, CORM-3, Hemin, zinc porphyrin (ZnPP), pinacidil, 5-hydroxydecanoate (5-HD), diltiazem, NG-nitro-L-arginine methyl ester (L-NAME), paxilline, wortmannin, LY-294002 (LY), and 1*H*-[1,2,4] oxadiazolo [4,3-*a*] quinoxalin-1-one (ODQ) were purchased from Sigma-Aldrich (St. Louis, MO, USA).

Preparation of perfused beating atria

Isolated perfused beating atria were prepared as previously described elsewhere [27,28]. In brief, hearts were rapidly excised from rats and a cannula was fixed into the left atria. The cannulated atria were fixed in an organ chamber, perfused with oxygenated 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) buffered saline (10 mM HEPES, 118 mM NaCl, 4.7 mM KCl, 2.5 mM CaCl₂, 1.2 mM MgSO₄, 25 mM NaHCO₃, 10 mM glucose, and 0.1% bovine serum albumin) at 37°C, and then paced at 1.2 Hz (30 V, 0.4 ms). The intra-atrial pressure was recorded using a Power Lab (ML-820, ADInstruments Pvt. Ltd., Sydney, Australia); atrial pulse pressure (APP) is defined as the difference between the systolic (SAP) and diastolic atrial pressure (DAP). The pericardial buffer solution that contained [³H]-inulin (Amersham Biosciences, Uppsala, Sweden) and that was used for measuring the translocation of ECF was also oxygenated through silicone tube coils inside the organ chamber. After ANP secretion stabilization and ECF translocation for 80 min, atrial perfusate was collected at 2 min intervals over 50 min at 4°C.

The groups were divided as follows. Group 1 was comprised of the control atria (n = 10). After a 10 min control period, atria were high-stretched by increasing the height of the outflow catheter from 5 cmH₂O to 7.5 cmH₂O by connecting a 2.5 cm long catheter, and atrial perfusate was continuously collected. The loaded volume to atria during diastole was 736 μl. Group 2 was comprised of atria perfused with CORM-2. After a 10 min control period, atria were perfused with various doses of CORM-2 (10, 50, 100 μM; n = 10 for each group) with high stretch. Group 3 was comprised of atria perfused with CORM-3, HO activator (hemin), and HO inhibitor (ZnPP). After a 10 min control period, atria were perfused with CORM-3 (10, 50, 100 μM; n = 10 for each group), hemin (0.1, 1, 50 μM; n = 10 for each group), ZnPP (50 μM; n = 10 for each group) with high stretch. Group 4 was comprised of atria perfused with CORM-2 (50 μM) in the presence of regulators, including K_{ATP} channel activator (pinacidil, 50 μM), mitochondrial K_{ATP} channel blocker (5-hydroxydecanoic acid, 5-HD, 100 μM), L-type Ca²⁺ channel blocker (diltiazem, 50 μM), high-conductance Ca²⁺-activated K⁺ (BK_{Ca}) channel blocker (paxilline, 25 μM), nitric oxide synthase (NOS) inhibitor (L-NAME, 100 μM), PI3K inhibitor (wortmannin, 10 μM; LY-294002, 10 μM), and soluble guanylate cyclase (sGC) inhibitor (ODQ, 100 μM). Atria were perfused with a regulator 20 min before sample collection, and CORM-2 (50 μM, n = 10 for each group) was simultaneously perfused into the high-stretched atria after a 10 min control period. Group 5 was comprised of atria perfused with CORM-2 (10 nM, 50 μM) in a hypoxic condition, as opposed to a normoxic condition. Atria were perfused with oxygen-free HEPES buffered saline, and CORM-2 (10 nM or 50 μM, n = 7 for each group) was simultaneously perfused into the high-stretched atria after a 10 min control period.

Measurement of ANP concentration and ECF translocation

The ANP concentration in the perfusate was measured using a specific radioimmunoassay (RIA) method previously described elsewhere [21]. The ANP in the plasma and coronary effluent was extracted using Sep-Pak C₁₈ cartridges, dried, and measured by RIA [29]. The intra- and inter-assay coefficients of variation were 6.3% (n = 9) and 7.8% (n = 11), respectively. The amount of secreted ANP was presented as ng of ANP/g atrial wet weight/min. The molar concentration of ANP released is determined by:

$$\text{ANP concentration } (\mu\text{M}) = \frac{\text{ANP secretion (ng/min/g)} \times 1,000}{\text{ECF translocation } (\mu\text{l/min/g}) \times 3,063}$$

in which 3,063 is the molecular mass of ANP [30,31].

The radioactivity of [³H]-inulin in the perfusates and pericardial buffer solution was measured with a 2 ml Ultima Gold cocktail solution (PerkinElmer, Waltham, MA, USA) using a liquid scintillation system (Tri-Card 300C; Packard, Downers Grove, IL, USA). ECF translocation was calculated as follows [30,31]:

$$\text{ECF translocation } (\mu\text{l/min/g}) = \frac{\text{Radioactivity in perfusate (cpm/min)} \times 1,000}{\text{Radioactivity in pericardial solution (cpm/100 } \mu\text{l)} \times \text{atrial wet weight (g)}}$$




Fig. 1. Effects of CORM-2 on ANP secretion and atrial contractility. (A) Effects of CORM-2 (10, 50, 100 μM) on ANP secretion (a), SAP (b), DAP (c) and APP (d) as a function of time in high-stretched atria. (B) Relative percent changes in atrial parameters by CORM-2. Values are expressed by the relative percent change from the mean of the first five control values (fraction no. 1–5) and the last five peak values (fraction no. 21–25). Values are expressed as the means ± SEM. Arrow indicates the time to start to perfuse CORM-2 with high atrial stretch. CORM, carbon monoxide releasing molecule; ANP, atrial natriuretic peptide; SAP, systolic atrial pressure; DAP, diastolic atrial pressure; APP, atrial pulse pressure. *** vs. vehicle group, p < 0.005; # vs. corresponding group, p < 0.05, ## p < 0.01.

Statistical analysis

All values are expressed as means \pm SEM. Differences in hemodynamic variables over time between the control and the treatment groups were assessed using ANOVA followed by Bonferroni's multiple comparison test (Sigmaplot; Systat Software, Inc., Erkrath, Germany). The significance level was set at $p < 0.05$.

RESULTS

Effects of CORM-2 on ANP secretion and atrial contractility

After a 10-min collection period, each dose of CORM-2 (10, 50, or 100 μ M) or vehicle was perfused into high-stretched atria, and atrial pressure and ANP secretion were measured. Fig. 1A shows ANP secretion, SAP, DAP, and APP over time. There was no significant difference in basal values of ANP secretion and atrial pressure among groups in this study. When the atria were high-stretched by increasing the height of outflow catheter from 5 cmH₂O to 7.5 cmH₂O, ANP secretion was gradually increased from 4.52 ± 0.73 ng/min/g of atrial tissue (at 10 min) to 11.54 ± 2.42 ng/min/g of atrial tissue (at 50 min) (Fig. 1Aa). SAP, DAP, and APP were increased suddenly and the higher level was maintained throughout the experiment (Fig. 1Ab–d). When 10 μ M of CORM-2 was perfused into the high-stretched atria, ANP secre-

tion was increased from 6.81 ± 0.91 ng/min/g of atrial tissue (at 10 min) to 16.88 ± 2.41 ng/min/g of atrial tissue (at 50 min) (Fig. 1Aa). When the dose of CORM-2 was increased to 50 μ M, ANP secretion was rose slightly from 5.20 ± 0.62 ng/min/g of atrial tissue (at 10 min) to 8.67 ± 0.57 ng/min/g of atrial tissue (at 50 min). However, 100 μ M of CORM-2 decreased ANP secretion from 8.08 ± 0.49 ng/min/g of atrial tissue to 6.09 ± 1.40 ng/min/g of atrial tissue.

To quantitatively compare the effects of CORM-2 on ANP secretion and atrial pressure, the relative percent change between the mean of the first five control values (fraction no. 1 to 5) and the mean of the last five peak values (fraction no. 21 to 25) was calculated (Fig. 1A). The ANP secretion in atria perfused with 10 μ M CORM-2 changed by $121.7 \pm 14.6\%$, which was similar to that observed in the control atria ($124.9 \pm 17.6\%$) (Fig. 1Ba). However, changes in ANP secretion in atria perfused with 50 μ M or 100 μ M CORM-2 were $63.9 \pm 12.1\%$ and $-29.1 \pm 14.3\%$, respectively, which were significantly lower than the amount of secretion observed in control atria ($124.9 \pm 17.6\%$) and in the 10 μ M CORM-2. There was a significant difference in ANP secretion induced by 10 μ M CORM-2 and 100 μ M CORM-2 (Fig. 1Ba). Changes in SAP, DAP, and APP after three doses of CORM-2 were no different from those observed in the control group (Fig. 1Bb–d).




Fig. 2. Comparison of changes in ANP secretion (A) and atrial contractility (B) in response to (a) CORM-2, (b) CORM-3, (c) hemin, and (d) ZnPP. Values are expressed by the relative percent change from the mean of the first five control values (fraction no. 1–5) and the last five peak values (fraction no. 21–25). CORM, carbon monoxide releasing molecule; ANP, atrial natriuretic peptide; ZnPP, zinc porphyrin; APP, atrial pulse pressure. Legends are the same as in Fig. 1. Values are expressed as the means \pm SEM. *** vs. vehicle group, $p < 0.005$

Effects of CORM-3, hemin, and zinc protoporphyrin on ANP secretion and atrial contractility

The effects on ANP secretion of other CO-related chemicals—CORM-3, hemin (a HO activator), and ZnPP (a HO inhibitor)—were compared with CORM-2. The same doses of CORM-3 (10, 50, and 100 μ M) as CORM-2 were used but changes in ANP secretion were not significantly different from those observed in the control atria ($121.7 \pm 14.5\%$, $120.7 \pm 7.7\%$, $103.5 \pm 14.5\%$ vs. $124.8 \pm 17.6\%$) (Fig. 2). Hemin 0.1 μ M did not suppress ANP secretion, but higher doses of hemin (1 and 50 μ M) markedly reduced ANP secretion ($45.3 \pm 1.6\%$, $-2.6 \pm 0.2\%$ vs. $124.8 \pm 17.6\%$) (Fig. 2A). No significant differences in APP were found among samples administered CORM-2, CORM-3, and hemin (Fig. 2B). The order of potency in suppressing ANP secretion was hemin > CORM-2 >> CORM-3. Interestingly, ZnPP (50 μ M) did not cause any significant change in ANP secretion (Fig. 2Ad) or APP (Fig. 2Bd).

To compare the effect of ANP secretion, CORM-2 or hemin (0.1 nM to 100 μ M) was perfused into isolated rat atria. As shown in Fig. 3, between 0.1 nM and 10 μ M CORM-2 did not result in any significant effects on ANP secretion. Nor did 0.1 nM to 100 nM hemin. CORM-2 and hemin began to suppress ANP secretion once administered at levels of 50 μ M and 1 μ M, respectively. Hemin appears to be stronger to suppress ANP secretion compare to CORM-2.

Modulation of the suppressive effect of CORM-2 on high stretch-induced ANP secretion

Previous reports have indicated that the ability of CO to relax

blood vessels occurs through cGMP-dependent and independent pathways [4]. To determine the mechanisms by which the suppressive effect of CORM-2 (50 μ M) on high stretch-induced ANP secretion operate, atria were pretreated with either mitochondrial K_{ATP} channel blocker (5-HD), K_{ATP} channel opener (pinacidil), L-type Ca²⁺ channel blocker (diltiazem), BK_{Ca} channel blocker (paxilline), NOS inhibitor (L-NAME), sGC inhibitor (ODQ), or PI3K inhibitor (wortmannin, LY). 5-HD (500 μ M) itself decreased ANP secretion ($56.1 \pm 8.9\%$, Fig. 4A) relative to the control group ($124.9 \pm 17.6\%$). Pretreatment with 5-HD blocked CORM-2-induced suppression of ANP secretion and concentration (Fig. 4). Pretreatment with pinacidil did not block CORM-2-induced suppression of ANP secretion (Fig. 4A), while pretreatment with paxilline was effective at blocking CORM-2-induced suppression of ANP secretion and concentration. ODQ blocked CORM-2-induced suppression of ANP concentration but not ANP secretion (Fig. 4D). Diltiazem, L-NAME, and PI3K inhibitors (LY, wortmannin) had no significant effect on CORM-2-induced suppression of ANP secretion (Fig. 4A).

Modulation of ANP secretion induced by CORM-2 during hypoxic condition

Low (10 nM) and high doses (50 μ M) of CORM-2 were used to compare the effect of CORM-2 on ANP secretion in normoxic and hypoxic conditions. In normoxic conditions, a high dose of CORM-2 suppressed ANP secretion ($63.8 \pm 12.1\%$ vs. $124.9 \pm 17.6\%$) (Fig. 5A), but a low dose did not. In hypoxic conditions, change in high stretched-ANP secretion was significantly higher ($363.7 \pm 8.5\%$ vs. $124.9 \pm 17.6\%$, $p < 0.01$) (Fig. 5A) and change




Fig. 3. Effects of wide range of doses of CORM-2 (A) and hemin (B) on ANP secretion and atrial contractility. Values are expressed by the relative percent change from the mean of the first five control values (fraction no. 1–5) and the last five peak values (fraction no. 21–25). CORM, carbon monoxide releasing molecule; ANP, atrial natriuretic peptide; APP, atrial pulse pressure. Legends are the same as in Fig. 1. Values are expressed as the means \pm SEM.




Fig. 4. Modulation of CORM-2-induced suppression of ANP secretion (A), APP (B), ECF translocation (C) and ANP concentration (D) by several modulators. Twenty minutes before sample collection, atria were pretreated with K_{ATP} channel activator (pinacidil, 50 μ M), mitochondrial K_{ATP} channel blocker (5-hydroxydecanoic acid; 5-HD, 100 μ M), L-type Ca^{2+} channel blocker (diltiazem; Dilt, 50 μ M), high-conductance Ca^{2+} -activated K^+ channel blocker (paxillin, 25 μ M), nitric oxide synthase inhibitor (L-NAME; 100 μ M), PI3K inhibitor (wortmannin; Wort, 10 μ M) (LY-294002; LY, 10 μ M), and soluble guanylate cyclase inhibitor (ODQ, 100 μ M) or vehicle was perfused into atria after 10-min control period. Relative percent changes in atrial parameters by CORM-2 and vehicle in the presence of modulators are shown. CORM, carbon monoxide releasing molecule; ANP, atrial natriuretic peptide; APP, atrial pulse pressure; ECF, extracellular fluid. Values are expressed as the means \pm SEM. * vs. vehicle group, $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$.




Fig. 5. Comparison of responsiveness of ANP secretion (A) and atrial contractility (B) by CORM-2 (10 nM, 50 μ M) in normoxic and hypoxic conditions. CORM, carbon monoxide releasing molecule; ANP, atrial natriuretic peptide; APP, atrial pulse pressure. Values are expressed as the means \pm SEM. * vs. vehicle group, $p < 0.05$, ** $p < 0.01$.

in APP was lower ($-34.3 \pm 7.5\%$ vs. $19.6 \pm 2.9\%$, $p < 0.01$) than in normoxic conditions (Fig. 5B). During hypoxia, CORM-2 50 μ M suppressed ANP secretion ($283.7 \pm 18.4\%$ vs. $363.7 \pm 8.5\%$,

$p < 0.05$; Fig. 4A). CORM-2 suppressed ANP secretion in both normoxic (by approximately 50%) and hypoxic conditions, (by approximately 20%). No significant differences in APP were ob-

served after administration of CORM-2 (Fig. 5B).

DISCUSSION

Our experiment showed that high doses of CORM-2 and hemin suppressed high stretch-induced ANP secretion, while CORM-3 and ZnPP did not suppress ANP secretion. The order of potency for the suppression of ANP secretion was hemin > CORM-2 >> CORM-3. The suppression of ANP secretion by CORM-2 was blocked by pretreatment with 5-HD, paxilline, and ODQ, but not by pretreatment with diltiazem, wortmannin, LY, and L-NAME. The effect of CORM-2 on ANP secretion appeared to have been attenuated in hypoxic conditions. These results suggest that CORM-2 suppresses ANP secretion via a mitochondrial K_{ATP} channel, a BK_{Ca} channel, and sGC pathway.

The beneficial effects of gasotransmitters generally manifest with lower doses, while the effects diminish at higher doses. In some cases, a higher dose may result in harmful levels of toxicity. Several recent studies have suggested that administration of a low level of CO either by CO inhalation or CORMs can protect the heart, liver, and kidneys against I/R [12,32,33]. In addition, low concentrations of CO also exhibit pleiotropic biological functions, including anti-apoptotic, cytoprotective, and anti-inflammatory effects [11-13]. We hypothesized that these protective effects may partly relate to the cardiac hormone ANP. Our aim, therefore, was to determine any direct effect of CO on ANP secretion, as well as the associated atrial dynamics, in normal isolated rat atria. Our experiments employed CORM-2 and HO activator (hemin) across a wide range of concentrations. At low concentrations (between 0.1 and 100 nM) CORM-2 and hemin had no significant effect on ANP secretion. At higher doses, stretch-activated ANP secretion was suppressed by CORM-2 and hemin. The dose-response curve of CORM-2 revealed no significant change in ANP secretion prior to 10 μM concentration. ANP secretion levels thereafter abruptly decreased with the administration of 50 μM and 100 μM of CORM-2. The dose-response curve of hemin was very similar to that of CORM-2, excepting a notable difference in potency. Our results are in broad agreement with a previous report showing that a characteristic gasotransmitter pharmacological response assumes the form of a bell-shaped dose-response curve [4,26]. A number of prior studies testing the beneficial effects of the administration of endogenous and exogenous CO in the case of I/R injury have been performed, with rat, mouse, and cardiac cell cultures exposed variously to CO [32], CORM-2 [33], and CORM-3 [12,34]. We compared the suppressive effect of CORM-2 on ANP secretion to that of CORM-3 and HO inhibitor. Notably, neither CORM-3 nor HO inhibitor (ZnPP) showed any significant effect on ANP secretion. While we are unable to explain why the effect of CORM-3 on ANP secretion is different from that of CORM-2, we hypothesize that it may be due to different rates of CO release.

CO can cause vaso-relaxation by activating sGC, increasing cGMP, and activating the BK_{Ca} channel [26,35-37]. K⁺ channel activation leads to membrane hyperpolarization followed by an inhibition of calcium entry via voltage operated Ca²⁺ channels. However, both cGMP-dependent and -independent signaling mechanisms are involved in the vascular action of CO [4,26]. It has also been reported that CORM-2-mediated cardioprotection may be a result of the activation of the p38 MAPK and protein kinase C pathways prior to ischemia, and the PI3K pathway during reperfusion [33]. Therefore, to elucidate the underlying signaling mechanisms involved in CORM-2-induced suppression of ANP secretion, atria were pretreated with several modulators. The suppression of ANP secretion by CORM-2 was completely blocked by pretreatment with the BK_{Ca} channel inhibitor (paxilline) and the mitochondrial K_{ATP} channel blocker (5-HD), and partially reduced by the sGC inhibitor. These results are consistent with a prior report showing complete blockage of a CORM-2 induced electrical current by a BK_{Ca} channel inhibitor [37] and a decrease in vascular resistance by CO through sGC activation and K_{Ca} channel opening in a mouse kidney [38]. However, diltiazem and wortmannin (including LY) did not block the effects of CORM-2. Another previous study has demonstrated that, with respect to CORM-2 mediated cardioprotection, PI3K is important during reperfusion rather than ischemia [33]. It follows that the differences in the signaling pathway of CORM-2 may be due to exposure time and duration to modulator and experimental tissues.

We also compared the effect of low and high doses of CORM-2 on ANP secretion in hypoxic conditions, which HO-1 can be induced. In the normoxic condition, CORM-2 (50 μM) suppressed ANP secretion by 50% compared to the control group (65% vs. 125%) whereas in a hypoxic condition, CORM-2 suppressed ANP secretion by 20% (280% vs. 360%). No change in the expression of HO-1 protein was observed in atria exposed to a hypoxic condition (data not shown). The attenuation of CORM-2's suppressive effect may be due to modulation of the K_{ATP} and BK_{Ca} channels during hypoxia.

In summary, these results suggest that CORM-2 suppresses ANP secretion via the mitochondrial K_{ATP} channel, BK_{Ca} channel, and sGC pathway.

FUNDING

This research was supported by a grant from the National Research Foundation of Korea (NRF) funded by the Korean government (MSIP) (NO 2020 R11A 3068178), as well as with funds provided by the Biomedical Research Institute of Jeonbuk National University Hospital.

ACKNOWLEDGEMENTS

We would like to express our gratitude to the editors of the Writing Center at Jeonbuk National University for their skilled English-language assistance.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

REFERENCES

1. Tenhunen R, Marver HS, Schmid R. The enzymatic conversion of heme to bilirubin by microsomal heme oxygenase. *Proc Natl Acad Sci U S A*. 1968;61:748-755.
2. Szabo C. Gaseotransmitters: new frontiers for translational science. *Sci Transl Med*. 2010;2:59ps54.
3. Caliendo G, Cirino G, Santagada V, Wallace JL. Synthesis and biological effects of hydrogen sulfide (H₂S): development of H₂S-releasing drugs as pharmaceuticals. *J Med Chem*. 2010;53:6275-6286.
4. Peers C, Steele DS. Carbon monoxide: a vital signalling molecule and potent toxin in the myocardium. *J Mol Cell Cardiol*. 2012;52:359-365.
5. Kim HP, Ryter SW, Choi AM. CO as a cellular signaling molecule. *Annu Rev Pharmacol Toxicol*. 2006;46:411-449.
6. Andreadou I, Iliodromitis EK, Rassaf T, Schulz R, Papapetropoulos A, Ferdinandy P. The role of gasotransmitters NO, H₂S and CO in myocardial ischaemia/reperfusion injury and cardioprotection by preconditioning, postconditioning and remote conditioning. *Br J Pharmacol*. 2015;172:1587-1606.
7. Haldane J. The relation of the action of carbonic oxide to oxygen tension. *J Physiol*. 1895;18:201-217.
8. André L, Gouzi F, Thireau J, Meyer G, Boissiere J, Delage M, Abdellaoui A, Feillet-Coudray C, Fouret G, Cristol JP, Lacampagne A, Obert P, Reboul C, Fauconnier J, Hayot M, Richard S, Cazorla O. Carbon monoxide exposure enhances arrhythmia after cardiac stress: involvement of oxidative stress. *Basic Res Cardiol*. 2011;106:1235-1246.
9. Dallas ML, Yang Z, Boyle JP, Boycott HE, Scragg JL, Milligan CJ, Elies J, Duke A, Thireau J, Reboul C, Richard S, Bernus O, Steele DS, Peers C. Carbon monoxide induces cardiac arrhythmia via induction of the late Na⁺ current. *Am J Respir Crit Care Med*. 2012;186:648-656.
10. Gandini C, Castoldi AF, Candura SM, Locatelli C, Butera R, Priori S, Manzo L. Carbon monoxide cardiotoxicity. *J Toxicol Clin Toxicol*. 2001;39:35-44.
11. Ryter SW, Otterbein LE. Carbon monoxide in biology and medicine. *Bioessays*. 2004;26:270-280.
12. Stein AB, Guo Y, Tan W, Wu WJ, Zhu X, Li Q, Luo C, Dawn B, Johnson TR, Motterlini R, Bolli R. Administration of a CO-releasing molecule induces late preconditioning against myocardial infarction. *J Mol Cell Cardiol*. 2005;38:127-134.
13. Muchova L, Wong RJ, Hsu M, Morioka I, Vitek L, Zelenka J, Schröder H, Stevenson DK. Statin treatment increases formation of carbon monoxide and bilirubin in mice: a novel mechanism of in vivo antioxidant protection. *Can J Physiol Pharmacol*. 2007;85:800-810.
14. Lakkisto P, Siren JM, Kytö V, Forsten H, Laine M, Pulkki K, Tikkanen I. Heme oxygenase-1 induction protects the heart and modulates cellular and extracellular remodelling after myocardial infarction in rats. *Exp Biol Med (Maywood)*. 2011;236:1437-1448.
15. Ewing JF, Raju VS, Maines MD. Induction of heart heme oxygenase-1 (HSP32) by hyperthermia: possible role in stress-mediated elevation of cyclic 3':5'-guanosine monophosphate. *J Pharmacol Exp Ther*. 1994;271:408-414.
16. Foresti R, Goatly H, Green CJ, Motterlini R. Role of heme oxygenase-1 in hypoxia-reoxygenation: requirement of substrate heme to promote cardioprotection. *Am J Physiol Heart Circ Physiol*. 2001;281:H1976-H1984.
17. Stec DE, Drummond HA, Vera T. Role of carbon monoxide in blood pressure regulation. *Hypertension*. 2008;51:597-604.
18. de Bold AJ, Borenstein HB, Veress AT, Sonnenberg H. A rapid and potent natriuretic response to intravenous injection of atrial myocardial extract in rats. *Life Sci*. 1981;28:89-94.
19. de Bold AJ. Thirty years of research on atrial natriuretic factor: historical background and emerging concepts. *Can J Physiol Pharmacol*. 2011;89:527-531.
20. Goetze JP, Bruneau BG, Ramos HR, Ogawa T, de Bold MK, de Bold AJ. Cardiac natriuretic peptides. *Nat Rev Cardiol*. 2020;17:698-717.
21. Cho KW, Seul KH, Ryu H, Kim SH, Koh GY. Characteristics of distension-induced release of immunoreactive atrial natriuretic peptide in isolated perfused rabbit atria. *Regul Pept*. 1988;22:333-345.
22. Zimmerman RS, Ryan J, Edwards BS, Klee G, Zimmerman D, Scott N, Burnett JC Jr. Cardiorenal endocrine dynamics during volume expansion in hypothyroid dogs. *Am J Physiol*. 1988;255(1 Pt 2):R61-R66.
23. Skvorak JP, Nazian SJ, Dietz JR. Endothelin acts as a paracrine regulator of stretch-induced atrial natriuretic peptide release. *Am J Physiol*. 1995;269(5 Pt 2):R1093-R1098.
24. Dietz JR. Mechanisms of atrial natriuretic peptide secretion from the atrium. *Cardiovasc Res*. 2005;68:8-17.
25. Varadi J, Lekli I, Juhasz B, Bacskay I, Szabo G, Gesztelyi R, Szendrei L, Varga E, Bak I, Foresti R, Motterlini R, Tosaki A. Beneficial effects of carbon monoxide-releasing molecules on post-ischemic myocardial recovery. *Life Sci*. 2007;80:1619-1626.
26. Motterlini R. Carbon monoxide-releasing molecules (CO-RMs): vasodilatory, anti-ischaemic and anti-inflammatory activities. *Biochem Soc Trans*. 2007;35(Pt 5):1142-1146.
27. Yuan K, Cao C, Han JH, Kim SZ, Kim SH. Adenosine-stimulated atrial natriuretic peptide release through A1 receptor subtype. *Hypertension*. 2005;46:1381-1387.
28. Han JH, Cao C, Kim SZ, Cho KW, Kim SH. Decreases in ANP secretion by lysophosphatidylcholine through protein kinase C. *Hypertension*. 2003;41:1380-1385.
29. Cho KW, Kim SH, Koh GY, Seul KH, Huh KS, Chu D, Rapp NS, Moon HB, Kim KK, Kook YJ. Plasma concentration of atrial natriuretic peptide in different phases of Korean hemorrhagic fever. *Nephron*. 1989;51:215-219.
30. Cho KW, Kim SH, Kim CH, Seul KH. Mechanical basis of ANP secretion in beating atria: atrial stroke volume and ECF translocation.

- Am J Physiol.* 1995;268(5 Pt 2):R1129-R1136.
31. Cho KW, Kim SH, Hwang YH, Seul KH. Extracellular fluid translocation in perfused rabbit atria: implication in control of atrial natriuretic peptide secretion. *J Physiol.* 1993;468:591-607.
 32. Rochetaing A, Barbé C, Kreher P. Acute ischemic preconditioning and high subchronic CO exposure independently increase myocardial tolerance to ischemia. *Inhal Toxicol.* 2001;13:1015-1032.
 33. Soni HM, Jain MR, Mehta AA. Mechanism(s) involved in carbon monoxide-releasing molecule-2-mediated cardioprotection during ischaemia-reperfusion injury in isolated rat heart. *Indian J Pharm Sci.* 2012;74:281-291.
 34. Clark JE, Naughton P, Shurey S, Green CJ, Johnson TR, Mann BE, Foresti R, Motterlini R. Cardioprotective actions by a water-soluble carbon monoxide-releasing molecule. *Circ Res.* 2003;93:e2-e8.
 35. Williams SE, Wootton P, Mason HS, Bould J, Iles DE, Riccardi D, Peers C, Kemp PJ. Hemoxygenase-2 is an oxygen sensor for a calcium-sensitive potassium channel. *Science.* 2004;306:2093-2097.
 36. Riesco-Fagundo AM, Pérez-García MT, González C, López-López JR. O₂ modulates large-conductance Ca²⁺-dependent K⁺ channels of rat chemoreceptor cells by a membrane-restricted and CO-sensitive mechanism. *Circ Res.* 2001;89:430-436.
 37. Wang R, Wu L, Wang Z. The direct effect of carbon monoxide on K_{Ca} channels in vascular smooth muscle cells. *Pflugers Arch.* 1997;434:285-291.
 38. Ryan MJ, Jernigan NL, Drummond HA, McLemore GR Jr, Rimoldi JM, Poreddy SR, Gadepalli RS, Stec DE. Renal vascular responses to CORM-A1 in the mouse. *Pharmacol Res.* 2006;54:24-29.