

## Supplementary References

1. Kawase T, Bishnoi I, Tanaka R, Dash C, Kato Y, Yamada Y : Study of incidence and factors: risk and preventive, of chronic subdural hematoma/hygroma in clipped patients of unruptured intracranial aneurysms - an institutional experience. **Asian J Neurosurg** **13** : 707-713, 2018
2. Kim JH, Kim CH, Lee CY : Efficacy of arachnoid-plasty on chronic subdural hematoma following surgical clipping of unruptured intracranial aneurysms. **World neurosurgery** **104** : 303-310, 2017
3. Lee WJ, Jo KI, Yeon JY, Hong SC, Kim JS : Incidence and risk factors of chronic subdural hematoma after surgical clipping for unruptured anterior circulation aneurysms. **J Korean Neurosurg Soc** **57** : 271-275, 2015
4. Lee WJ, Nam TM, Jo KI, Yeon JY, Hong SC, Kim JS : Modified arachnoid plasty reduces chronic subdural hematoma after unruptured aneurysm clipping : technical note. **J Korean Neurosurg Soc** **61** : 761-766, 2018
5. Murakami T, Nakagawa I, Park HS, Kotsugi M, Takamura Y, Takeshima Y, et al. : Extensive postoperative subdural fluid volume affects the onset of chronic subdural hematoma after unruptured aneurysmal clipping surgery. **Clin Neurol Neurosurg** **187** : 105533, 2019
6. Yagi K, Irie S, Inagaki T, Ishii Y, Saito O, Lee T, et al. : Intraoperative arachnoid plasty has possibility to prevent chronic subdural hematoma after surgery for unruptured cerebral aneurysms. **Neurol Med Chir (Tokyo)** **55** : 493-497, 2015