

Supplementary Table 3. Biomarker Values in Patients Who Proceeded to Colectomy Compared to Patients Who Avoided Colectomy by 12 Months, Taken at Baseline, Day 1 Post Infliximab Salvage and Day 3 Post Infliximab Salvage

Biomarker	Colectomy	No colectomy	P-value
Baseline (admission)			
CRP, mg/L (n = 94)	65 (39–177)	46 (20–100)	0.140
Albumin, g/L (n = 94)	27 (22–32)	32 (27–35)	0.025
Neutrophils, 10 ⁹ /L (n = 94)	7.4 (4.7–10.7)	7.9 (6.0–11.3)	0.560
Lymphocytes, 10 ⁹ /L (n = 94)	1.0 (0.8–1.7)	1.5 (1.0–2.1)	0.060
Platelets, 10 ⁹ /L (n = 94)	392 (324–496)	392 (298–449)	0.660
Day 1 post infliximab			
CRP, mg/L (n = 93)	32 (16–89)	11 (4–26)	<0.001
Albumin, g/L (n = 91)	21 (18–27)	27 (24–31)	0.003
Neutrophils, 10 ⁹ /L (n = 93)	7.0 (4.5–11.4)	8.5 (6.6–11.4)	0.240
Lymphocytes, 10 ⁹ /L (n = 93)	1.3 (0.7–2.4)	1.8 (1.3–2.7)	0.090
Platelets, 10 ⁹ /L (n = 93)	362 (259–481)	376 (306–459)	0.600
Day 3 post infliximab			
CRP, mg/L (n = 84)	27 (11–56)	5 (2–12)	<0.001
Albumin, g/L (n = 81)	23 (18–27)	28 (24–31)	<0.001
Neutrophils, 10 ⁹ /L (n = 84)	6.3 (5.9–8.8)	8.7 (6.2–12.0)	0.060
Lymphocytes, 10 ⁹ /L (n = 84)	1.3 (0.7–2.0)	2.8 (1.8–4.2)	0.001
Platelets, 10 ⁹ /L (n = 84)	380 (271–500)	382 (318–466)	0.640

Values are presented as median (interquartile range).
CRP, C-reactive protein.