

**Supplementary Table 5.** Diagnostic Properties of CAR and CLR on Day 3 Post Salvage, Using Outcomes of 12-Month Colectomy and 1-Month Colectomy

Cutoff	Predicting colectomy by 12 months				Predicting colectomy by 1 month				Proportion classified as low risk (%)
	Sens.	Spec.	PPV	NPV	Sens.	Spec.	PPV	NPV	
<b>CAR (mg/g)</b>									
≥ 0.14	1.00	0.44	0.31	1.00	1.00	0.39	0.16	1.00	35
≥ 0.28	0.90	0.59	0.35	0.96	1.00	0.55	0.21	1.00	50
≥ 0.47 <sup>a</sup>	0.79	0.80	0.50	0.94	0.90	0.75	0.30	0.98	68
≥ 1.1	0.53	0.93	0.67	0.89	0.70	0.91	0.47	0.96	85
≥ 1.7	0.42	0.99	0.89	0.87	0.50	0.95	0.56	0.94	90
<b>CLR (mg/10<sup>9</sup>)</b>									
≥ 1.5	0.95	0.41	0.29	0.97	1.00	0.38	0.16	1.00	35
≥ 2.6	0.84	0.60	0.35	0.94	0.90	0.56	0.20	0.98	50
≥ 6.0 <sup>a</sup>	0.84	0.84	0.57	0.96	0.90	0.77	0.32	0.98	70
≥ 16	0.63	0.96	0.80	0.91	0.80	0.92	0.53	0.97	85
≥ 24	0.42	0.99	0.89	0.87	0.50	0.95	0.56	0.94	90

<sup>a</sup>Optimal cutoff defined as maximum Youden's statistic.

CAR, C-reactive protein (CRP)-albumin ratio; CLR, CRP-lymphocyte rate; Sens., sensitivity; Spec., specificity; PPV, positive predictive value; NPV, negative predictive value.