

# Clinical Significance of p53 Protein Expression, Beta-catenin Expression and HER2 Expression for Epstein-Barr Virus-associated Gastric Cancer

Dong Won Baek<sup>1</sup>, Byung Woog Kang<sup>1,\*</sup>, Soyeon Hwang<sup>1</sup>, Jong Gwang Kim<sup>1,\*</sup>, An Na Seo<sup>2</sup>, Han Ik Bae<sup>2</sup>, Oh Kyoung Kwon<sup>3</sup>, Seung Soo Lee<sup>3</sup>, Ho Young Chung<sup>3</sup>, and Wansik Yu<sup>3</sup>

<sup>1</sup>Department of Oncology/Hematology, Kyungpook National University Medical Center, Kyungpook National University Cancer Research Institute, Kyungpook National University School of Medicine, Departments of <sup>2</sup>Pathology, <sup>3</sup>Surgery, Kyungpook National University Hospital, Kyungpook National University School of Medicine, Daegu, Korea

This study assessed the expression of the p53 protein, beta-catenin, and HER2 and their prognostic implications in patients with EBV-associated gastric cancer (EBVaGC). After reviewing 1318 consecutive cases of surgically resected or endoscopic submucosal dissected gastric cancers, 117 patients were identified as EBV-positive using EBV-encoded RNA in-situ hybridization. The immunohistochemistry results were interpreted as follows: strong p53 nuclear expression in at least 50% of tumor nuclei was interpreted as a positive result, strong beta-catenin expression in at least 10% of cytoplasmic nuclei was interpreted as a positive result, and moderate or strong complete or basolateral membrane staining in 10% of tumor cells was interpreted as a positive result for HER2. Immunohistochemical staining for p53 was performed on tumor tissue from 105 patients, among whom 25 (23.8%) tested positive. Meanwhile, beta-catenin expression was positive in 10 patients (17.5%) and HER2 expression was positive in 8 patients (6.8%). The positive expression of p53 was significantly associated with a high T stage ( $p=0.006$ ). More patients with lymph node metastasis were p53-positive ( $p=0.013$ ). In the univariate analysis, the p53-positive patients showed significantly decreased disease-free survival (DFS) when compared with the p53-negative patients ( $p=0.022$ ), although the p53 status was only marginally associated with overall survival (OS) ( $p=0.080$ ). However, p53 expression showed no prognostic significance on DFS in the multivariate analysis. Moreover, beta-catenin and HER2 showed no association with DFS and OS in the survival analysis. The current study found a significant correlation between p53 expression and tumor progression and lymph node metastases in patients with EBVaGC.

**Key Words:** Epstein-Barr Virus Infections; Stomach Neoplasms; Tumor Suppressor Protein p53; Beta Catenin; Human epidermal growth factor receptor-2

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Recent data from the The Cancer Genome Atlas (TCGA) data revealed that Epstein-Barr virus (EBV)-associated tumors showed a *PIK3CA* mutation, *PD-L1/2* overexpression, EBV-CIMP, *CDKN2A* silencing, and immune cell signaling.<sup>1</sup> Moreover, EBV infection can cause tumorigenesis and malignant transformation of host cells

through the activation of oncogenic pathways.<sup>2,3</sup> Various evidence sources have also indicated that gene alterations may play an important role in the pathogenesis of EBV infection, including the promotion of pathological changes seen in gastric cancer.<sup>4,5</sup> Some representative examples are p53, beta-catenin, and human epidermal growth factor receptor-2 (HER2), which are already known to be associated with the pathogenesis of EBVaGC. Therefore, studies of

### Article History:

Received January 25, 2017

Revised March 10, 2017

Accepted March 16, 2017

### Corresponding Author:

Jong Gwang Kim  
Department of Oncology/Hematology,  
Kyungpook National University  
Medical Center, Kyungpook National  
University School of Medicine,  
Kyungpook National University  
Cancer Research Institute, 807  
Hoguk-ro, Buk-gu, Daegu 41404,  
Korea  
Tel: +82-53-200-2623  
Fax: +82-53-200-3399  
E-mail: jkk21c@knu.ac.kr

Byung Woog Kang  
Department of Oncology/Hematology,  
Kyungpook National University  
Medical Center, Kyungpook National  
University School of Medicine, 807  
Hoguk-ro, Buk-gu, Daegu 41404,  
Korea  
Tel: +82-53-200-2622  
Fax: +82-53-200-3399  
E-mail: bwkang@knu.ac.kr

molecular alterations could provide potential biomarkers as therapeutic targets for EBV-associated gastric cancer (EBVaGC).

The *TP53* gene is a tumor suppressor gene that is composed of 11 exons and encodes a nuclear p53 protein, which acts as a potent transcription factor with a key role in the maintenance of genetic stability.<sup>6</sup> Thus, genetic alterations of the *TP53* gene could be involved in gastric carcinogenesis. Previous studies have already demonstrated that a *TP53* mutation is one of the most prevalent genetic alterations in gastric cancer.<sup>3,7</sup> Plus, the association of p53 expression with tumor biological behavior and prognosis has also been reported in the gastric cancer patients.<sup>8</sup> Interestingly, EBV infection has been associated with abnormal p53 expression, which may play an important role in inhibiting cell apoptosis and the tumorigenesis of EBVaGC.<sup>4,7,9</sup> Meanwhile, beta-catenin is an important mediator of the Wnt signaling pathway, and mutation of beta-catenin has been identified in various human malignancies, including gastric cancer.<sup>10</sup> Furthermore, Wnt signaling, induced by activated beta-catenin, regulates T cell responses.<sup>11</sup> Therefore, these findings point to the possibility of beta-catenin as a prognostic and predictive marker in gastric cancer patients with EBV tumors, suggesting a pivotal role for the immune mechanism in these subsets of gastric cancer. In the ToGA trial, human epidermal growth factor receptor-2 (HER2)-positive gastric cancer patients who received first-line treatment with trastuzumab, an antibody targeting HER2, showed an improved overall survival.<sup>12</sup> Several recent clinical studies have also investigated the link between HER2 and EBVaGC, where Zhang et al. reported that HER2 expression was inversely correlated with EBV infection.<sup>13,14</sup> Notwithstanding, the relationship between EBVaGC and HER2 remains controversial.<sup>15,16</sup>

Accordingly, given that p53 protein, beta-catenin, and HER2 may be potential biomarkers as therapeutic targets for EBVaGC, this study assessed the expression of the p53 protein, beta-catenin, and HER2 and their prognostic implication in patients with EBVaGC.

## PATIENTS AND METHODS

### 1. Study population

All the tissue samples used in this study were obtained from patients included in a previous study of EBVaGC, the inclusion criteria and results for which have already been reported.<sup>5</sup> In brief, the study included 117 patients with histologically confirmed adenocarcinomas of the stomach that were identified as EBV-positive using EBV-encoded RNA *in situ* hybridization. Among the 117 patients, p53 expression was examined in 105 tumor tissues and beta-catenin in 57 tumor tissues, respectively. The baseline characteristics, including age, gender, TNM stage according to the American Joint Committee on Cancer (AJCC) staging 7th edition, and tumor histologic differentiation were all collected from the patients' medical records and surgical

pathologic reports. This study was approved by the institutional review board at Kyungpook National University Medical Center (KNUMC) and informed consent was obtained from all the patients included in this study.

### 2. Immunohistochemistry and scoring

Immunohistochemical staining was performed using a Ventana Benchmark XT autoimmunostainer (Ventana Medical Systems, Tucson, AZ, USA) according to the manufacturer's instructions. Antibodies for the following molecules were used in this study: p53 (1:800; DAKO, Denmark), beta-catenin (1:200; DAKO, Denmark), and HER2 (pre-dilution; Ventana Medical Systems, Tucson, AZ, USA). The immunohistochemistry results were interpreted as follows: strong p53 nuclear expression in at least 50% of the tumor nuclei was interpreted as a positive result, strong beta-catenin expression in at least 10% of the cytoplasmic nuclei was interpreted as a positive result, and moderate or strong complete or basolateral membrane staining in 10% of the tumor cells was interpreted as a pos-

TABLE 1. Baseline patient characteristics

Characteristic	Total (N= 117)
	N (%)
Age, years	
Median (Range)	62.0 (32.0-80.0)
Gender	
Male	93 (79.5)
Female	24 (20.5)
Tumor location	
Gastroesophageal junction, cardia and fundus	11 (9.4)
Body	82 (70.1)
Antrum	10 (8.5)
Others	14 (12.0)
Tumor depth (pathologic T category)	
T1	67 (57.3)
T2	11 (9.4)
T3	19 (16.2)
T4	20 (17.1)
Lymph node metastasis (pathologic N category)	
N0	86 (73.5)
N1	12 (10.3)
N2	8 (6.8)
N3	11 (9.4)
pTNM stage	
I	74 (63.2)
II	25 (21.4)
III	18 (15.4)
Lauren classification	
Intestinal	25 (21.4)
Diffuse	81 (69.2)
Mixed	11 (9.4)
WHO classification	
Gastric carcinoma with lymphoid stroma (GCLS)	57 (48.7)
Non-GCLS	60 (51.3)

itive result for HER2. The histopathologic and immunohistochemistry analyses were conducted by two pathologists (A.N.S and H.I.B) at KNUMC.

### 3. Statistical analysis

The descriptive statistics are reported as proportions and medians. The categorical variables were evaluated using a  $\chi^2$  test and Fisher's exact test, as appropriate. Disease-free survival (DFS) was defined as the time between diagnosis to tumor recurrence or death from any cause. Overall survival (OS) was calculated from the date of diagnosis to death from any cause. Data was censored if patients were free of recurrence or alive at the last follow-up. The Kaplan-Meier method was used to estimate the DFS and OS. The survival curves were compared using a log rank test according to the expression differences. Multivariate survival analyses were carried out using the Cox proportional hazard regression model. A p-value < 0.05 was considered statistically significant. The statistical analyses were performed using SPSS for Windows (version 19.0, SPSS Inc., Chicago, Ill., USA).

## RESULTS

### 1. Patient characteristics

The patient characteristics are shown in Table 1. Most

of the patients were male (79.5%) and the median age was 62 years (range 32-80). Most of the tumors (70.1%) were located in the body of the stomach. At the initial resection, 74 (63.2%) tumors were classified as stage I, 25 (21.4%) as stage II, 18 (15.4%) as stage III. Plus, according to the Lauren classification, 25 (21.4%) patients' tumors were intestinal, 81 (69.2%) were diffuse, and 11 (9.4%) were mixed type. Among the 117 patients, 57 (48.7%) were classified as having gastric carcinoma with lymphoid stroma (GCLC).

### 2. Immunohistochemistry analysis for p53, beta-catenin and HER2

Immunohistochemical staining for p53 was performed on 105 tumor tissues, among which 25 tissues (23.8%) tested positive (Table 2). Positive expression of p53 was significantly associated with a high T stage (p=0.006). More tumors with lymph node metastasis were p53-positive (p=0.013). Beta-catenin expression was examined in 57 tumor tissues (Table 3). No difference in characteristics was noted according to the positive or negative expression of beta-catenin. Plus, no significant difference in tumor-infiltrating lymphocytes (TILs) was found according to the positive or negative expression of beta-catenin (p=0.735). Positive HER2 expression was only observed in 10 tumor tissues.

**TABLE 2.** Association between clinicopathologic features and p53 expression


	p53			p
	< 50 Negative	≥ 50 Positive	Total (N=105)	
	N (%)	N (%)	N (%)	
Gender				1.000
Male	64 (80.0)	20 (80.0)	84 (80.0)	
Female	16 (20.0)	5 (20.0)	21 (20.0)	
Age				0.492
< 62	40 (50.0)	10 (40.0)	50 (47.6)	
≥ 62	40 (50.0)	15 (60.0)	55 (52.4)	
T stage				0.006
T1/2	61 (76.2)	11 (44.0)	72 (68.6)	
T3/4	19 (23.8)	14 (56.0)	33 (31.4)	
LN metastasis				0.013
No	66 (82.5)	14 (56.0)	80 (76.2)	
Yes	14 (17.5)	11 (44.0)	25 (23.8)	
Stage				0.007
I	58 (72.5)	11 (44.0)	69 (66.0)	
II	15 (18.8)	6 (24.0)	21 (20.0)	
III	7 (8.7)	8 (32.0)	15 (14.0)	
WHO classification				0.360
GCLS	42 (52.5)	10 (40.0)	52 (49.5)	
Non-GCLS	38 (47.5)	15 (60.0)	53 (50.5)	
Lauren classification				0.867
Intestinal	17 (21.25)	7 (28.0)	24 (22.9)	
Diffuse	56 (70.0)	16 (64.0)	72 (68.5)	
Mixed	7 (8.75)	2 (8.0)	9 (8.6)	

GCLS: Gastric carcinoma with lymphoid stroma.

**TABLE 3.** Association between clinicopathologic features and beta-catenin expression

	Beta-catenin			p
	Negative	Positive	Total (N=57)	
	N (%)	N (%)	N (%)	
Gender				1.000
Male	39 (83.0)	9 (90.0)	48 (84.2)	
Female	8 (17.0)	1 (10.0)	9 (15.8)	
Age				0.467
< 62	18 (32.3)	2 (20.0)	20 (35.1)	
≥ 62	29 (61.7)	8 (80.0)	37 (64.9)	
T stage				0.708
T1/2	33 (70.2)	8 (80.0)	41 (71.9)	
T3/4	14 (29.8)	2 (20.0)	16 (28.1)	
LN metastasis				1.000
No	34 (72.3)	7 (70.0)	41 (71.9)	
Yes	13 (27.7)	3 (30.0)	16 (28.1)	
WHO classification				0.166
GCLS	22 (46.8)	2 (20.0)	24 (42.1)	
Non-GCLS	25 (53.2)	8 (80.0)	33 (57.9)	
Lauren classification				0.184
Intestinal	11 (23.4)	5 (50.0)	16 (28.1)	
Diffuse	33 (70.2)	4 (40.0)	37 (64.9)	
Mixed	3 (6.4)	1 (10.0)	4 (7.0)	

GCLS: Gastric carcinoma with lymphoid stroma.

**FIG. 1.** Kaplan-Meier survival curves for disease-free survival (A) and overall survival (B) according to p53 status.

### 3. Survival according to p53 expression and beta-catenin

With a median follow-up duration of 36.8 months (range 6.5-60.6), the estimated 5-year DFS and OS rates were 83.9% and 81.0%, respectively. In the univariate analysis, the p53-positive EBVaGC patients showed an reduced DFS when compared with p53-negative EBVaGC patients ( $p=0.022$ ), although the p53 status was only marginally associated with OS ( $p=0.080$ ; Fig. 1). However, p53 expression showed no prognostic significance on DFS in the multivariate analysis (Table 4). Moreover, beta-catenin expression was not significantly associated with DFS ( $p=0.341$ ) or OS ( $p=0.472$ ; Fig. 2). Similarly, HER2 ex-

pression was not found to be associated with DFS or OS in the survival analysis (data not shown).


### DISCUSSION

This study investigated the clinical significance of the p53 protein, beta-catenin, and HER2 expression in 117 patients with curatively resected EBVaGC. As a result, the overexpression of p53 was found to be associated with a significantly higher T stage and increased lymph node metastasis, suggesting that impaired p53 activity may be involved in the tumor progression of EBVaGC. Regarding the

**TABLE 4.** Multivariate analysis of disease-free survival for patients with EBVaGC

Variables	Category	Disease-free survival in all patients					
		Uivariate analysis			Multivariate analysis		
		HR	95% CI	p	HR	95% CI	p
Age, years	< 62 vs. ≥62	1.26	0.33-4.73	0.724	1.16	0.29-4.50	0.830
pTNM Stage	I & II vs. III	9.78	2.59-36.85	0.001	5.87	1.17-29.44	0.031
P53	< 50 vs. ≥50	4.12	1.10-15.38	0.022	1.87	0.42-8.33	0.407
Gender	Male vs. Female	0.42	0.05-3.38	0.402	0.52	0.06-4.42	0.555
Sub-classification	Non-GCLC vs. GCLC	0.22	0.04-1.10	0.066	0.57	0.08-3.80	0.570

GCLS: Gastric carcinoma with lymphoid stroma.

**FIG. 2.** Kaplan-Meier survival curves for disease-free survival (A) and overall survival (B) according to beta-catenin status.

relationship between p53 overexpression and prognosis, while a trend was indicated, this needs to be elucidated in further studies.

EBV is a gammaherpesvirus whose latent infection is reported to be associated with approximately 10% of all gastric cancers.<sup>17</sup> EBV can infect cells in a latent or lytic manner.<sup>18</sup> EBVaGC also belongs to latency type I, which can include the expression of the viral genes EBV nuclear antigen 1 (EBNA1), latent membrane protein 2A (LMP2A), Bam HI-A rightward transcripts (BARTs), and EBV-encoded small RNA (EBER1/2).<sup>19</sup> In particular, the expression of latency genes fulfils a relevant task in the initiation and neoplastic progression of EBV-associated epithelial cancers, including EBVaGC.<sup>20</sup> Interestingly, these viral factors are known to interact with various cellular proteins, such as p53, which maintains genomic integrity in multicellular organisms by activating DNA repair, arresting cells at the G1/S checkpoint, and inducing apoptosis.<sup>21</sup> For instance, EBNA1 expression decreases p53 activation and apoptosis in response to DNA damage, resulting in increased cell survival in gastric cancer.<sup>22</sup> Plus, EBNA3C can down regulate p53 transcriptional activity in EBV-infected cells.<sup>23</sup> Moreover, cumulative studies reported that the p53 protein is accumulated and phosphorylated in nasopharyngeal cancer, mediated by LMP1.<sup>24-26</sup> The present study found an association between p53 over-

expression and tumor progression. Similar to the current results, Chen et al. reported that p53 mutation was associated with the initiation stages of gastric cancer.<sup>27</sup> Although the exact mechanism remains unclear, the presence of clonal EBV in every tumor cell suggests that EBV infection precedes the final transforming event giving rise to a tumor.<sup>28</sup> Yet, despite the association between advanced stage and p53 overexpression in EBVaGC tissue, no prognostic significance of p53 for DFS and OS was demonstrated. To date, the prognostic role of p53 expression detected by immunohistochemistry in gastric cancer remains controversial.<sup>7,29</sup> Some studies have linked the absence of p53 expression with longer survival, implying that p53 is a negative prognostic factor.<sup>30-32</sup> Conversely, other studies have found no relation between p53 expression and survival.<sup>33-35</sup> Therefore, its prognostic relevance should be evaluated in further studies aimed at elucidating the biological significance of p53 expression in EBVaGC.

Beta-catenin has already been shown to be an important mediator of the Wnt signaling cascades.<sup>36</sup> As a central molecule in the Wnt signaling pathway, beta-catenin expression is localized in the membrane, cytoplasm and nucleus. Cytoplasmic and nuclear beta-catenin are mainly involved in regulation of the Wnt signaling pathway.<sup>37</sup> Beta-catenin accumulates in the nucleus and activates a

target gene expression program with loss of E-cadherin, linking Epithelial-mesenchymal transition to Wnt signaling.<sup>38</sup> Recent studies demonstrated that aberrant activation of the Wnt/beta-catenin signaling pathway is involved in the development and progression of a significant proportion of gastric cancer cases.<sup>10,39,40</sup> In particular, activation of the Wnt/beta-catenin signaling pathway is involved in the immune system, including lymphocytes and immunosuppressive cytokines.<sup>41</sup> Therefore, beta-catenin could be an important biomarker in EBVaGC, as EBV-positive tumors often show immune signaling activation.<sup>1</sup>

However, the present study found no statistically significant prognostic impact of beta-catenin in EBVaGC. Plus, no significant correlations were observed between beta-catenin and TILs. There are several possible reasons for this finding. First, the sample size was too small to make conclusions about the prognostic significance of beta-catenin. Second, the methodology used to interpret the expression of beta-catenin and the cutoff values need to be standardized. In this study, strong staining in 10% or more of the cytoplasmic-nuclei was interpreted as positive expression of beta-catenin. However, a recent study by Retterspitz et al. regarded that widespread staining of at least 35% of tumor nuclei indicated a positive expression.<sup>40</sup> Thus, different sensitivities have been used to interpret the expression degree of beta-catenin.

HER2, a known proto-oncogene located on the long arm of human chromosome 17 (17q12), is a member of the epidermal growth factor receptor (EGFR) superfamily associated with tumor cell proliferation, apoptosis, adhesion, migration, and differentiation.<sup>42,43</sup> In gastric cancer, HER2 overexpression is associated with poor outcomes and aggressive disease.<sup>44</sup> Thus, due to its pivotal role in tumor growth and relationship with therapeutic efficacy in gastric cancer, HER2 expression in EBVaGC was also evaluated in the current study. Clinical studies already reported that HER2 overexpression was only found in 1.6-5.6% of EBVaGC cases, which is consistent with the present result in terms of the frequency of HER2.<sup>16,45</sup> Interestingly, Zhang et al. reported a significantly reduced expression of HER2 in EBVaGC, possibly induced by LMP2A, a EBV latent protein.<sup>14</sup> EBVaGC has distinct pathological features, presenting a generally diffuse-type histology.<sup>9,46</sup> Thus, HER2 overexpression is relatively rare, making it difficult to analyze the relationship between HER2 and survival in EBVaGC.

In conclusion, the current study found a significant correlation between p53 expression and tumor progression in patients with EBVaGC. However, tissue expression of p53, beta-catenin, and HER2 was not identified as an independent prognostic marker in EBVaGC. Therefore, further investigation is needed to clarify the precise biologic significance of p53 in EBVaGC.

## ACKNOWLEDGEMENTS

This work was supported by a National Research

Foundation of Korea (NRF) grant funded by the Korea government. (2014R1A5A2009242)

## CONFLICT OF INTEREST STATEMENT

None declared.

## REFERENCES

1. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature* 2014; 513:202-9.
2. Tsao SW, Tsang CM, To KF, Lo KW. The role of Epstein-Barr virus in epithelial malignancies. *J Pathol* 2015;235:323-33.
3. Gonzalez RS, Messing S, Tu X, McMahon LA, Whitney-Miller CL. Immunohistochemistry as a surrogate for molecular subtyping of gastric adenocarcinoma. *Hum Pathol* 2016;56:16-21.
4. Szkaradkiewicz A, Karpiński TM, Majewski J, Malinowska K, Goślińska-Kuźniarek O, Linke K. The participation of p53 and bcl-2 proteins in gastric carcinomas associated with Helicobacter pylori and/or Epstein-Barr Virus (EBV). *Pol J Microbiol* 2015;64: 211-6.
5. Kang BW, Seo AN, Yoon S, Bae HI, Jeon SW, Kwon OK, et al. Prognostic value of tumor-infiltrating lymphocytes in Epstein-Barr virus-associated gastric cancer. *Ann Oncol* 2016;27:494-501.
6. Belyi VA, Ak P, Markert E, Wang H, Hu W, Puzio-Kuter A, et al. The origins and evolution of the p53 family of genes. *Cold Spring Harb Perspect Biol* 2010;2:a001198.
7. Bellini MF, Cadamuro AC, Succi M, Proenca MA, Silva AE. Alterations of the TP53 gene in gastric and esophageal carcinogenesis. *J Biomed Biotechnol* 2012;2012:891961.
8. Tahara T, Shibata T, Okamoto Y, Yamazaki J, Kawamura T, Horiguchi N, et al. Mutation spectrum of TP53 gene predicts clinicopathological features and survival of gastric cancer. *Oncotarget* 2016;7:42252-60.
9. Lee JH, Kim SH, Han SH, An JS, Lee ES, Kim YS. Clinicopathological and molecular characteristics of Epstein-Barr virus-associated gastric carcinoma: a meta-analysis. *J Gastroenterol Hepatol* 2009;24:354-65.
10. Chiurillo MA. Role of the Wnt/ $\beta$ -catenin pathway in gastric cancer: an in-depth literature review. *World J Exp Med* 2015;5: 84-102.
11. Staal FJ, Arens R. Wnt signaling as master regulator of T lymphocyte responses: implications for transplant therapy. *Transplantation* 2016. doi: 10.1097/TP.0000000000001393. [Epub ahead of print]
12. Bang YJ, Van Cutsem E, Feyereislova A, Chung HC, Shen L, Sawaki A, et al. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet* 2010;376:687-97.
13. Li Z, Lai Y, Sun L, Zhang X, Liu R, Feng G, et al. PD-L1 expression is associated with massive lymphocyte infiltration and histology in gastric cancer. *Hum Pathol* 2016;55:182-9.
14. Zhang YW, Zhao XX, Tan C, Zhang ZG, Jiang Y, Chen JN, et al. Epstein-Barr virus latent membrane protein 2A suppresses the

- expression of HER2 via a pathway involving TWIST and YB-1 in Epstein-Barr virus-associated gastric carcinomas. *Oncotarget* 2015;6:207-20.
15. Irkkan C, Balci S, Guler Tezel G, Akinci B, Yalcin B, Guler G. Comparison of clinicopathologic parameters and survivals between Epstein-Barr Virus-positive and Her2-positive gastric cancers. *Appl Immunohistochem Mol Morphol* 2016. doi: 10.1097/PAI.0000000000000353. [Epub ahead of print]
  16. Sukawa Y, Yamamoto H, Noshio K, Kunimoto H, Suzuki H, Adachi Y, et al. Alterations in the human epidermal growth factor receptor 2-phosphatidylinositol 3-kinase-v-Akt pathway in gastric cancer. *World J Gastroenterol* 2012;18:6577-86.
  17. Takada K. Epstein-Barr virus and gastric carcinoma. *Mol Pathol* 2000;53:255-61.
  18. Young LS, Rickinson AB. Epstein-Barr virus: 40 years on. *Nat Rev Cancer* 2004;4:757-68.
  19. Yau TO, Tang CM, Yu J. Epigenetic dysregulation in Epstein-Barr virus-associated gastric carcinoma: disease and treatments. *World J Gastroenterol* 2014;20:6448-56.
  20. Giudice A, D'Arena G, Crispo A, Tecce MF, Nocerino F, Grimaldi M, et al. Role of viral miRNAs and epigenetic modifications in Epstein-Barr Virus-associated gastric carcinogenesis. *Oxid Med Cell Longev* 2016;2016:6021934.
  21. Liu Y, Kulesz-Martin M. p53 protein at the hub of cellular DNA damage response pathways through sequence-specific and non-sequence-specific DNA binding. *Carcinogenesis* 2001;22:851-60.
  22. Sivachandran N, Dawson CW, Young LS, Liu FF, Middeldorp J, Frappier L. Contributions of the Epstein-Barr virus EBNA1 protein to gastric carcinoma. *J Virol* 2012;86:60-8.
  23. Shukla SK, Jha HC, El-Naccache DW, Robertson ES. An EBV recombinant deleted for residues 130-159 in EBNA3C can deregulate p53/Mdm2 and Cyclin D1/CDK6 which results in apoptosis and reduced cell proliferation. *Oncotarget* 2016;7:18116-34.
  24. Li L, Li W, Xiao L, Xu J, Chen X, Tang M, et al. Viral oncoprotein LMP1 disrupts p53-induced cell cycle arrest and apoptosis through modulating K63-linked ubiquitination of p53. *Cell Cycle* 2012;11:2327-36.
  25. Guo L, Tang M, Yang L, Xiao L, Bode AM, Li L, et al. Epstein-Barr virus oncoprotein LMP1 mediates survivin upregulation by p53 contributing to G1/S cell cycle progression in nasopharyngeal carcinoma. *Int J Mol Med* 2012;29:574-80.
  26. Yang CF, Peng LX, Huang TJ, Yang GD, Chu QQ, Liang YY, et al. Cancer stem-like cell characteristics induced by EB virus-encoded LMP1 contribute to radioresistance in nasopharyngeal carcinoma by suppressing the p53-mediated apoptosis pathway. *Cancer Lett* 2014;344:260-71.
  27. Chen HC, Chen HJ, Khan MA, Rao ZZ, Wan XX, Tan B, et al. Genetic mutations of p53 and k-ras in gastric carcinoma patients from Hunan, China. *Tumour Biol* 2011;32:367-73.
  28. Imai S, Koizumi S, Sugiura M, Tokunaga M, Uemura Y, Yamamoto N, et al. Gastric carcinoma: monoclonal epithelial malignant cells expressing Epstein-Barr virus latent infection protein. *Proc Natl Acad Sci U S A* 1994;91:9131-5.
  29. Yildirim M, Kaya V, Demirpence O, Gunduz S, Bozcuk H. Prognostic significance of p53 in gastric cancer: a meta-analysis. *Asian Pac J Cancer Prev* 2015;16:327-32.
  30. Begnami MD, Fregnani JH, Nonogaki S, Soares FA. Evaluation of cell cycle protein expression in gastric cancer: cyclin B1 expression and its prognostic implication. *Hum Pathol* 2010;41:1120-7.
  31. Goncalves AR, Carneiro AJ, Martins I, de Faria PA, Ferreira MA, de Mello EL, et al. Prognostic significance of p53 protein expression in early gastric cancer. *Pathol Oncol Res* 2011;17:349-55.
  32. Ye YW, Zhang X, Zhou Y, Wu J, Zhao C, Yuan L, et al. The correlations between the expression of FGFR4 protein and clinicopathological parameters as well as prognosis of gastric cancer patients. *J Surg Oncol* 2012;106:872-9.
  33. Lazăr D, Tăban S, Sporea I, Dema A, Cornianu M, Lazăr E, et al. The immunohistochemical expression of the p53-protein in gastric carcinomas. Correlation with clinicopathological factors and survival of patients. *Rom J Morphol Embryol* 2010;51:249-57.
  34. Murakami D, Tsujitani S, Osaki T, Saito H, Katano K, Tatebe S, et al. Expression of phosphorylated Akt (pAkt) in gastric carcinoma predicts prognosis and efficacy of chemotherapy. *Gastric Cancer* 2007;10:45-51.
  35. Tsujitani S, Saito H, Wakatsuki T, Ikeguchi M, Shirabe K, Morita M, et al. Relationship between expression of apoptosis-related proteins and the efficacy of postoperative chemotherapy in patients with T3 gastric cancer. *Surg Today* 2012;42:225-32.
  36. Scanlon CS, Van Tubergen EA, Inglehart RC, D'Silva NJ. Biomarkers of epithelial-mesenchymal transition in squamous cell carcinoma. *J Dent Res* 2013;92:114-21.
  37. Schmalhofer O, Brabletz S, Brabletz T. E-cadherin, beta-catenin, and ZEB1 in malignant progression of cancer. *Cancer Metastasis Rev* 2009;28:151-66.
  38. Behrens J. Cadherins and catenins: role in signal transduction and tumor progression. *Cancer Metastasis Rev* 1999;18:15-30.
  39. Li LF, Wei ZJ, Sun H, Jiang B. Abnormal  $\beta$ -catenin immunohistochemical expression as a prognostic factor in gastric cancer: a meta-analysis. *World J Gastroenterol* 2014;20:12313-21.
  40. Retterspitz MF, Monig SP, Schreckenber S, Schneider PM, Holscher AH, Dienes HP, et al. Expression of (beta)-catenin, MUC1 and c-met in diffuse-type gastric carcinomas: correlations with tumour progression and prognosis. *Anticancer Res* 2010;30:4635-41.
  41. Yaguchi T, Goto Y, Kido K, Mochimaru H, Sakurai T, Tsukamoto N, et al. Immune suppression and resistance mediated by constitutive activation of Wnt/ $\beta$ -catenin signaling in human melanoma cells. *J Immunol* 2012;189:2110-7.
  42. Gravalos C, Jimeno A. HER2 in gastric cancer: a new prognostic factor and a novel therapeutic target. *Ann Oncol* 2008;19:1523-9.
  43. Hynes NE, Stern DF. The biology of erbB-2/neu/HER-2 and its role in cancer. *Biochim Biophys Acta* 1994;1198:165-84.
  44. Liang JW, Zhang JJ, Zhang T, Zheng ZC. Clinicopathological and prognostic significance of HER2 overexpression in gastric cancer: a meta-analysis of the literature. *Tumour Biol* 2014;35:4849-58.
  45. Lee HS, Chang MS, Yang HK, Lee BL, Kim WH. Epstein-barr virus-positive gastric carcinoma has a distinct protein expression profile in comparison with epstein-barr virus-negative carcinoma. *Clin Cancer Res* 2004;10:1698-705.
  46. Moritani S, Kushima R, Sugihara H, Hattori T. Phenotypic characteristics of Epstein-Barr-virus-associated gastric carcinomas. *J Cancer Res Clin Oncol* 1996;122:750-6.