



Comparison of Nutrient Intake and Diet Quality Between Hyperuricemia Subjects and Controls in Korea

Kyoung A Ryu¹, Hyun Hee Kang¹, So Young Kim¹, Min Kyong Yoo¹,
Jeong Seon Kim², Chan Haw Lee³, Gyung Ah Wie^{1*}

¹Department of Clinical Nutrition, National Cancer Center, Goyang 410-769, South Korea

²Division of Cancer Epidemiology & Prevention, Research Institute, National Cancer Center, Goyang 410-769, South Korea

³Center for Cancer Prevention and Detection, National Cancer Center, Goyang 410-769, South Korea

Hyperuricemia is associated with metabolic syndrome as well as gout, and the prevalence of hyperuricemia is increasing in Korea. This study aimed to compare the nutrient intake and diet quality between hyperuricemia subjects and controls. Of the 28,589 people who participated in a health examination between 2008 and 2011, 9,010 subjects were selected whose 3-day food records were available. Clinical and laboratory data were collected from electronic medical records. Diet quality was evaluated using the food habit score (FHS), nutrient adequacy ratio (NAR), and mean adequacy ratio (MAR). The prevalence of hyperuricemia was 13.8% (27.1%, men; 5.2%, women). Body mass index, waist circumference, triglycerides, total cholesterol, and low-density lipoprotein cholesterol were significantly higher ($p < 0.0001$), while high-density cholesterol ($p < 0.001$) was significantly lower in the hyperuricemia subjects than in the controls. The hyperuricemia subjects had a lower intake of vitamin A ($p < 0.004$), vitamin C, folate, fiber, and calcium than the controls ($p < 0.0001$). Intake of vegetables and dairy products was significantly lower, whereas alcohol intake was significantly higher in the hyperuricemia subjects than in the controls ($p < 0.0001$). The FHS ($p < 0.0001$), MAR ($p < 0.0001$), and NARs for vitamin A ($p = 0.01$), vitamin B₂, vitamin C, folate, and calcium ($p < 0.0001$) were significantly lower in the hyperuricemia subjects than in the controls. In conclusion, the hyperuricemia subjects reported poorer diet quality than the controls, including higher alcohol intake and lower vegetable and dairy product intake.

Key Words: Hyperuricemia, Uric acid, Diet records, Nutritive Value, Food Habits

*Corresponding author Gyung Ah Wie

Address Department of Clinical Nutrition, National Cancer Center,
111 Jungbalsan-ro, Ilsandong-gu, Goyang 410-769, South Korea

Tel +82-31-920-0273 **Fax** +82-31-920-0269

E-mail gawie@ncc.re.kr

Received December 3, 2013

Revised January 7, 2014

Accepted January 9, 2014

© 2014 The Korean Society of Clinical Nutrition

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

The global prevalence of hyperuricemia is increasing [1], and this may be related to recent changes in diet, lifestyle and increasing prevalence of obesity. There appears to be a recent increase in the prevalence of hyperuricemia owing to the rapid economic development and the effect of westernized dietary habits in Korea [2].

Hyperuricemia is a precursor to gout and is associated with metabolic syndrome and cardiovascular disease. Many studies have reported that elevated uric acid is one of the most important risk factors for cardiovascular disease and that it plays a significant role in the development of metabolic syndrome [3-7]. Hyperuricemia can induce endothelial dysfunction in addition to inflammatory and oxidative changes, which

Nutrient Intake and Diet Quality in Hyperuricemia

may explain the association with metabolic syndrome [8]. Uric acid is an organic compound that is endogenously produced as a purine metabolite. It is formed by the liver and primarily excreted by the kidneys (65–75%) and intestines (25–35%) [9]. Hyperuricemia occurs due to a decrease in uric acid excretion, an overproduction of uric acid, or an excessive intake of purine. But the primary cause is known to be a decrease in the excretion by the kidneys [10].

Although a purine-rich diet is responsible for increasing uric acid by only 1–2 mg/dL [11], the management of hyperuricemia has been focused on dietary recommendations to avoid purine-rich foods such as meats and meat products and alcoholic beverage [12]. To our knowledge, a few studies have reported the prevalence and clinical features of hyperuricemia in Korea [13–15], but there has been no study on the associations between diet and hyperuricemia among Korean adults. Therefore, the objective of our study was to investigate and compare nutrient intake and diet quality between hyperuricemia subjects and controls in Korea.

Materials and Methods

Subjects

This cross-sectional study compared nutritional intake and quality between hyperuricemia subjects and controls. Of the 28,589 subjects who underwent health examination between January 2008 and December 2011, those that completed a self-administered questionnaire, laboratory tests, and a complete 3-day food record were included. Subjects were excluded who reported an unrealistic daily energy intake (<500 kcal or $\geq 3,500$ kcal). This resulted in 9,010 subjects in the final analysis (Figure 1).

Anthropometric and biochemical measurements

Weight (kg) and height (m) were measured using an automatic body composition analyzer (X-scan Plus II, Jawon Medical, Gyeongsan, South Korea) in light clothing and with shoes off, and body mass index (BMI) was calculated (kg/m^2). The waist circumference was taken at the narrowest point between the lower costal margin and the superior iliac crest.

Biochemical data, including uric acid, glucose, and serum lipid such as total cholesterol, serum triglycerides (TG), high-density lipoprotein (HDL) cholesterol, and low-density lipoprotein (LDL) cholesterol levels, were collected from the medical records relating to the health examination. During the screen-




Figure 1. Inclusion and exclusion criteria for the study subjects. *Visitors for health examination survey from January 1, 2008 to December 31, 2011; †Hyperuricemia was defined as serum uric acid concentration ≥ 7 mg/dL in men, and ≥ 6 mg/dL in women [16].

ing, venous blood samples were collected in the morning following a 10–12 h overnight fast.

Nutrient intake assessment

Usual dietary intake was assessed using 3-day food records (1 weekend and 2 week days). A self-administered 3-day food record and written instructions were sent via post to the subjects 2 weeks before the scheduled visit.

During the scheduled visit, the food records were reviewed by trained nutritionists using aids such as food models to improve the reporting accuracy. The daily energy, macro-, and micro-nutrient intakes were calculated using the Korean Nutrition Society nutrient database (Can-Pro 3.0, The Korean Nutrition Society, Seoul, South Korea) [17]. Nutrient density (ND) was calculated by standardizing the nutrient intake per 1,000 kcal based on the individual's intake.

Dietary habits and diet quality assessment

Dietary habits were evaluated using a self-reported 10-item questionnaire focusing on skipped meals, meal speed, meal regularity, and preference for salty foods. For each question, subjects selected Yes (1 point) or No (0 points), and the total Food Habit Score (FHS) ranged from 0–10 points. A higher score indicated better food habits.

The nutrient adequacy ratio (NAR; nutrient intake of an individual/recommended nutrient intake for that nutrient) and mean adequacy ratio (MAR; sum of the NAR for all of the nutrients/number of nutrients) were calculated to assess the

appropriateness of nutrient intake. In addition, the index of nutritional quality (INQ; nutrient intake per 1,000 kcal/recommended nutrient intake per 1,000 kcal), a comparison of the nutrient intake against the 12 nutrients for which the recommended nutrient intake (RNI) is defined by the Korean Dietary Reference Intake (KDRIs), was analyzed.

Statistical analysis

Data were expressed as frequencies or mean \pm standard deviation and were analyzed using SAS ver.9.1 (SAS Institute, Cary, NC, USA). A generalized linear model (GLM) was used to compare the distribution of subjects and the mean differences between the hyperuricemia subjects and controls. The significance level was set at $p < 0.05$.

Results

The overall prevalence of hyperuricemia was 13.8% (27.1%, men; 5.2%, women), and the proportion of men was significantly higher in the hyperuricemia subjects than in the controls (83.7% vs 49.3%, $p < 0.0001$). Table 1 provided the demographic and clinical characteristics of the subjects, compared between hyperuricemia subjects and controls. BMI and energy intake were also significantly higher in the hyperuricemia

subjects than in the controls (both $p < 0.0001$).

The age-, sex-, and BMI-adjusted value for waist circumference, triglycerides, total cholesterol, and LDL cholesterol were all significantly higher ($p < 0.0001$), and HDL cholesterol ($p < 0.001$) was significantly lower in the hyperuricemia subjects than in controls.

The differences in ND between the hyperuricemia subjects and controls were provided in Table 2. Hyperuricemia subjects had significantly lower intakes of carbohydrate ($p < 0.01$), fiber ($p < 0.001$), calcium ($p < 0.001$), phosphorus ($p < 0.001$), iron ($p < 0.001$), sodium ($p < 0.05$), potassium ($p < 0.001$), vitamin A ($p < 0.01$), vitamin B₂ ($p < 0.01$), vitamin C ($p < 0.001$), and folate ($p < 0.001$).

Table 3 provided the comparison of daily food group intake between the groups. The intake of vegetables ($p < 0.001$), seaweed ($p < 0.05$), and dairy products ($p < 0.001$) was significantly lower in the hyperuricemia subjects than in controls, and alcohol intake was significantly higher with the hyperuricemia consuming approximately twice the amount of alcohol than the controls (137.1 g vs 69.2 g, $p < 0.001$). Although the consumption of meat, eggs, fish, and shellfish was higher in the hyperuricemia, the differences were not significant.

Table 4 listed the total and component FHS compared between the hyperuricemia subjects and controls. The total FHS

Table 1. Demographic and clinical characteristics of the study subjects

Variables	Hyperuricemia subjects (n = 1242)	Controls (n = 7768)	p value
Age, yrs	50.5 \pm 10.0	50.9 \pm 9.6	NS
Male : Female, %	1039 (83.7) : 203 (16.3)	3830 (49.3) : 3938 (50.7)	$p < 0.0001$
Body Mass Index, kg/m ²	25.5 \pm 2.9	23.4 \pm 2.9	$p < 0.0001$
Total energy intake, kcal	1970.1 \pm 508.9	1799.9 \pm 471.5	$p < 0.0001$
Waist Circumference, cm*	89.8 \pm 7.5	83.6 \pm 8.5	$p < 0.0001$
Uric acid, mg/dL*	7.6 \pm 0.9	4.9 \pm 1.0	$p < 0.0001$
SBP, mmHg*	125.7 \pm 13.5	121.0 \pm 14.2	NS
DBP, mmHg*	77.5 \pm 10.0	73.8 \pm 10.2	NS
FBS, mg/dL*	91.8 \pm 18.2	89.3 \pm 18.7	$p < 0.01$
Triglyceride, mg/dL*	161.9 \pm 116.8	107.9 \pm 71.0	$p < 0.0001$
Total-cholesterol, mg/dL*	207.3 \pm 37.5	196.4 \pm 34.1	$p < 0.0001$
HDL-cholesterol, mg/dL*	52.5 \pm 12.4	59.2 \pm 14.1	$p < 0.001$
LDL-cholesterol, mg/dL*	131.3 \pm 32.7	120.8 \pm 30.3	$p < 0.0001$

SBP: systolic blood pressure, DBP: diastolic blood pressure, FBS: fasting blood sugar, HDL-cholesterol: high-density lipoprotein cholesterol, LDL-cholesterol: low-density lipoprotein cholesterol, NS: not significant

Data are presented as mean \pm standard deviation; Significance as determined by the GLM test.

*Adjusted for age, sex, body mass index.

Table 2. Comparisons of the nutrient density (ND; intake/1,000 kcal) between hyperuricemia subjects and controls

Variables	Hyperuricemia subjects (n = 1242)	Controls (n = 7768)	p value*
Carbohydrate, g	148.7 ± 25.3	154.2 ± 22.2	p < 0.001
Protein, g	40.0 ± 7.1	40.1 ± 7.0	NS
Fat, g	23.1 ± 6.9	23.3 ± 6.8	NS
Fiber, g	10.5 ± 3.1	11.8 ± 3.4	p < 0.0001
Cholesterol, mg	161.4 ± 72.9	161.0 ± 75.5	NS
Calcium, mg	268.7 ± 100.4	300.3 ± 108.4	p < 0.0001
Phosphorus, mg	572.3 ± 104.1	595.8 ± 105.3	p < 0.0001
Iron, mg	7.6 ± 2.7	8.0 ± 2.6	p < 0.0001
Sodium, mg	2318.1 ± 749.0	2407.1 ± 789.2	p < 0.01
Potassium, mg	1524.9 ± 403.4	1655.1 ± 435.9	p < 0.0001
Zinc, mg	5.7 ± 3.6	5.7 ± 3.0	NS
Vitamin A, µg	391.6 ± 198.1	437.9 ± 247.5	p < 0.001
Vitamin E, mg	7.0 ± 2.4	7.3 ± 2.5	NS
Vitamin B ₁ , mg	0.7 ± 0.3	0.7 ± 0.7	NS
Vitamin B ₂ , mg	0.6 ± 0.2	0.6 ± 0.2	p < 0.001
Vitamin B ₆ , mg	1.0 ± 0.3	1.1 ± 0.8	NS
Niacin, mg	9.5 ± 2.4	9.4 ± 2.5	NS
Vitamin C, mg	57.5 ± 33.5	69.7 ± 39.8	p < 0.0001
Folate, µg	187.1 ± 97.9	208.3 ± 105.8	p < 0.0001

NS: not significant.

Data are presented as mean ± standard deviation; Significance as determined by the GLM test.

*Adjusted for age, sex, body mass index.

was significantly lower in the hyperuricemia subjects than in the controls (p < 0.001). The items that contributed to the lower score in the hyperuricemia were meal regularity, vegetable consumption, fruit consumption, and dairy product consumption.

The NAR and MAR values for the hyperuricemia subjects and controls were provided in Table 5. The NARs for calcium (p < 0.001), vitamin A (p < 0.05), vitamin B₂ (p < 0.001), vitamin C (p < 0.001), and folate (p < 0.001) were significantly lower in the hyperuricemia subjects than in the controls. The MAR was also significantly lower in the hyperuricemia subjects than in the controls (p < 0.001).

The comparison of the INQ values was shown in Table 6. The INQs for calcium (p < 0.001), vitamin A (p < 0.01), vitamin B₂ (p < 0.01), vitamin C (p < 0.001), and folate (p < 0.001) were significantly lower, while the phosphorus (p < 0.01) and iron (p < 0.01) values were significantly higher in the hyperuricemia subjects than in the controls.

Discussion

To our knowledge, this study was the first to compare nutrient intake and diet quality between hyperuricemia subjects and controls in Korea. A larger portion of the hyperuricemia subjects was men, and they also had higher BMIs, poorer lipid profiles, and poorer diet quality than the controls.

Previous studies have also reported that the prevalence of hyperuricemia is higher in men than in women. Such a difference between genders is probably due to the estrogen which purportedly increases urinary excretion of uric acid [11,18]. Poletto et al. [18] reported that the prevalence of hyperuricemia, in 1,330 Japanese-Brazilian adults was lower in younger women than in older women (women <55 years, 24%; women ≥55 years, 76%; p < 0.05). Similarly, in the current study, the prevalence of hyperuricemia was significantly higher in men than in women.

Obesity is another strong risk factor for the development of hyperuricemia. A large prospective study indicated that weight gain was the risk factor for hyperuricemia in men, while

Table 3. Comparisons of daily food group intake between hyperuricemia subjects and controls

Variables	Hyperuricemia subjects (n = 1242)	Controls (n = 7768)	p value*
Total, g	1446.4 ± 482.1	1374.0 ± 434.2	NS
Cereals, g	289.2 ± 88.4	267.8 ± 92.8	NS
Potato and starches, g	32.5 ± 58.2	37.7 ± 67.4	NS
Sugars and sweeteners, g	7.1 ± 8.5	6.8 ± 8.6	NS
Pulses, g	46.9 ± 51.5	47.2 ± 54.2	NS
Nuts and seeds, g	5.7 ± 15.5	6.5 ± 15.0	NS
Vegetables, g	323.0 ± 142.8	326.0 ± 154.5	p < 0.0001
Fungi and mushrooms, g	4.7 ± 8.8	5.4 ± 15.3	NS
Fruits, g	189.1 ± 203.3	213.6 ± 200.0	NS
Meats, g	93.4 ± 74.0	77.2 ± 65.9	NS
Eggs, g	26.5 ± 28.2	24.6 ± 24.4	NS
Fish and shellfish, g	94.1 ± 70.3	81.3 ± 64.3	NS
Seaweeds, g	2.8 ± 5.5	3.5 ± 9.0	p < 0.01
Dairy products, g	66.6 ± 98.9	85.6 ± 106.0	p < 0.0001
Oils and fat, g	8.4 ± 5.5	7.5 ± 5.1	NS
Beverages, g	84.9 ± 163.8	81.4 ± 149.4	NS
Seasoning, g	33.4 ± 19.8	31.3 ± 17.7	NS
Alcohols, g	137.1 ± 277.5	69.2 ± 182.3	p < 0.0001
Others, g	0.8 ± 3.1	0.9 ± 5.3	NS

NS: not significant.

Data are presented as mean ± standard deviation; Significance as determined by the GLM test.

*Adjusted for age, sex, body mass index.

Table 4. Comparisons of the Food Habit Score (FHS) between hyperuricemia subjects and controls

Variables	Hyperuricemia subjects (n = 1242)	Controls (n = 7768)	p value*
Total score (0-10)	5.50 ± 1.92	5.83 ± 1.89	p < 0.0001
Meal regularity	0.80 ± 0.40	0.83 ± 0.38	p < 0.001
Meal speed	0.78 ± 0.42	0.83 ± 0.38	NS
Meat/egg consumption	0.40 ± 0.49	0.37 ± 0.48	NS
Seafood consumption	0.55 ± 0.50	0.53 ± 0.50	NS
Tofu, soy, soymilk consumption	0.50 ± 0.50	0.53 ± 0.50	NS
Vegetables, seaweeds, mushrooms consumption	0.51 ± 0.50	0.55 ± 0.50	p < 0.01
Fruits consumption	0.54 ± 0.50	0.65 ± 0.48	p < 0.001
Dairy products consumption	0.36 ± 0.48	0.43 ± 0.49	p < 0.01
Variety of diet	0.87 ± 0.34	0.86 ± 0.34	p < 0.01
Salty taste	0.50 ± 0.50	0.55 ± 0.50	NS

NS: not significant.

Data are presented as mean ± standard deviation; Significance as determined by the GLM test.

*Adjusted for age, sex, body mass index.

Nutrient Intake and Diet Quality in Hyperuricemia

Table 5. Comparison of the Nutrient Adequacy Ratio (NAR) and Mean Adequacy Ratio (MAR) between hyperuricemia subjects and controls

Variables	Hyperuricemia subjects (n = 1242)	Controls (n = 7768)	p value*
Nutrient Adequacy Ratio (NAR)			
Protein	0.99 ± 0.05	0.98 ± 0.06	NS
Calcium	0.69 ± 0.20	0.72 ± 0.21	p < 0.0001
Phosphorus	0.99 ± 0.05	0.99 ± 0.06	NS
Iron	0.97 ± 0.09	0.94 ± 0.13	NS
Zinc	0.95 ± 0.10	0.94 ± 0.13	NS
Vitamin A	0.84 ± 0.21	0.85 ± 0.20	p < 0.01
Vitamin B ₁	0.91 ± 0.14	0.90 ± 0.14	p < 0.001
Vitamin B ₂	0.76 ± 0.19	0.78 ± 0.19	p < 0.0001
Vitamin B ₆	0.96 ± 0.10	0.95 ± 0.10	NS
Niacin	0.93 ± 0.13	0.91 ± 0.14	NS
Vitamin C	0.83 ± 0.21	0.86 ± 0.20	p < 0.0001
Folate	0.74 ± 0.24	0.75 ± 0.24	p < 0.0001
Mean Adequacy Ratio (MAR)	0.88 ± 0.09	0.88 ± 0.10	p < 0.0001

NS: not significant.

Data are presented as mean ± standard deviation; Significance as determined by the GLM test.

*Adjusted for age, sex, body mass index.

Table 6. Comparison of the Index of Nutritional Quality (INQ) between hyperuricemia subjects and controls

Variables	Hyperuricemia subjects (n = 1242)	Controls (n = 7768)	p value*
Protein	1.72 ± 0.32	1.69 ± 0.31	NS
Calcium	0.82 ± 0.30	0.87 ± 0.31	p < 0.0001
Phosphorus	1.80 ± 0.35	1.73 ± 0.33	p < 0.001
Iron	1.76 ± 0.61	1.69 ± 0.66	p < 0.001
Zinc	1.42 ± 0.91	1.38 ± 0.73	NS
Vitamin A	1.22 ± 0.61	1.33 ± 0.74	p < 0.001
Vitamin B ₁	1.28 ± 0.69	1.29 ± 1.22	NS
Vitamin B ₂	0.90 ± 0.25	0.96 ± 0.30	p < 0.001
Vitamin B ₆	1.52 ± 0.44	1.51 ± 1.04	NS
Niacin	1.34 ± 0.35	1.28 ± 0.35	NS
Vitamin C	1.25 ± 0.71	1.40 ± 0.77	p < 0.0001
Folate	1.03 ± 0.53	1.06 ± 0.53	p < 0.0001

NS: not significant.

Data are presented as mean ± standard deviation; Significance as determined by the GLM test.

*Adjusted for age, sex, body mass index.

weight loss reduced the risk [19]. Our study also showed that BMI was significantly higher in the hyperuricemia subjects than in the controls.

A number of previous studies have reported that hyperuri-

cemia is associated with various chronic diseases such as arterial hypertension, cardiovascular disease, dyslipidemia, diabetes mellitus, and gout, all of which are associated with an increase in mortality risk [3-7, 20-24]. Furthermore, animal studies indi-

cate that a decrease in uric acid levels can prevent or reverse the features of metabolic syndrome [25-26]. We also found significantly higher waist circumference, triglyceride, total cholesterol, and LDL cholesterol and lower HDL cholesterol in the hyperuricemia subjects than in the controls after adjusting for age, sex, and BMI.

Relationships between the consumption of animal protein and hyperuricemia have been reported in the literature [27]. In this study, the intake of the purine-rich foods such as meat, fish, and shellfish tended to be higher in the hyperuricemia subjects than in the controls. The intake of vegetables, seaweed, and dairy products was significantly lower, while alcohol intake was higher in the hyperuricemia subjects than in the controls. Consequently, the intake of fiber, potassium, vitamin C, and folate was lower and resulted in lower NARs for calcium, vitamin A, vitamin B₂, vitamin C, and folate in the hyperuricemia subjects than in the controls.

Previous studies indicate that a greater vitamin C intake is independently associated with lower serum uric acid levels via its uricosuric effects [28-29]. The presence of milk-forming proteins such as lactalbumin and casein, which have uricosuric effects [30], may explain the inverse relationship observed between the consumption of dairy products and uric acid levels [31]. However, there does not appear to be an association between calcium supplementation and uric acid levels [2].

Alcohol abuse can increase the production of uric acid by promoting the metabolism of adenine nucleotide to uric acid [32]. In addition, ethanol can decrease renal uric acid excretion through an increase in lactic acid concentrations, which then competitively suppress the excretion of acid in the renal tubules by activating the ion exchange function of human urate anion exchanger (URAT1) to stimulate the reabsorption of uric acid in the proximal tubule [33-34]. We also found that alcohol intake in the hyperuricemia subjects were significantly higher than in the controls.

In summary, the results of the dietary quality indices indicated that the hyperuricemia subjects had poorer diet quality than the controls, especially relating to meal regularity and the intake of vegetables, fruit, and dairy products. Few studies have examined the associations between diet quality and hyperuricemia. However, a number of studies have reported that poor diet quality is likely to result in obesity, abdominal obesity, elevated blood pressure, hyperlipidemia, and other risk factors associated with metabolic syndrome [35-37].

Our study had certain limitations. First, the dietary information might not reflect an individual's usual intake owing to

the use of 3-day food records. Second, it was not possible to establish causal associations between hyperuricemia and diet, because of the cross-sectional study design. Therefore, prospective or longitudinal studies are necessary to further elucidate the dietary factors related to hyperuricemia among Koreans. However, the strengths of the current study include the large sample size and the related statistical power. Furthermore, this study was the first to investigate nutrient intake and diet quality in people with hyperuricemia in Korea.

In conclusion, the hyperuricemia subjects reported poorer diet quality than the controls, including higher alcohol intake and lower vegetable and dairy product intake. Our findings suggest that a balanced diet that includes consumption of dairy products and plenty of vegetables, limited alcohol intake, and maintenance of a healthy weight can assist with the management of hyperuricemia. However, further research is needed to clarify the role of dietary factors in hyperuricemia and to establish evidence-based dietary guidelines for the management of hyperuricemia.

Conflict of interest

We declare that we have no conflict of interest.

References

1. Puig JG, Martínez MA, Mora M, Fraile JM, Montoya F, Torres RJ. Serum urate, metabolic syndrome, and cardiovascular risk factors. A population-based study. *Nucleosides Nucleotides Nucleic Acids* 2008;27:620-3.
2. Zgaga L, Theodoratou E, Kyle J, Farrington SM, Agakov F, Tenesa A, Walker M, McNeill G, Wright AF, Rudan I, Dunlop MG, Campbell H. The association of dietary intake of purine-rich vegetables, sugar-sweetened beverages and dairy with plasma urate, in a cross-sectional study. *PLoS One* 2012;7:e38123.
3. Fang J, Alderman MH. Serum uric acid and cardiovascular mortality the NHANES I epidemiologic follow-up study, 1971-1992. *National Health and Nutrition Examination Survey. JAMA* 2000;283:2404-10.
4. Fessel WJ. High uric acid as an indicator of cardiovascular disease. Independence from obesity. *Am J Med* 1980;68:401-4.
5. Chien KL, Chen MF, Hsu HC, Chang WT, Su TC, Lee YT, Hu FB. Plasma uric acid and the risk of type 2 diabetes in a Chinese community. *Clin Chem* 2008;54:310-6.
6. Heinig M, Johnson RJ. Role of uric acid in hypertension, renal disease, and metabolic syndrome. *Cleve Clin J Med* 2006;73:1059-64.
7. Burack RC, Keller JB, Higgins MW. Cardiovascular risk factors and obesity: are baseline levels of blood pressure, glucose, cholesterol and uric acid elevated prior to weight gain? *J Chronic Dis* 1985;38:865-72.
8. Feig DI, Kang DH, Johnson RJ. Uric acid and cardiovascular risk. *N Engl J Med* 2008;359:1811-21.
9. de Oliveira EP, Burini RC. High plasma uric acid concentration: causes and consequences. *Diabetol Metab Syndr* 2012;4:12.
10. Lee KH, Lee SK. Pathophysiology of gout. *Korean J Med* 2011;80:251-4.
11. Emmerson BT. The management of gout. *N Engl J Med* 1996;334:445-51.
12. Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G. Purine-rich foods, dairy and protein intake, and the risk of gout in men. *N Engl J*

- Med 2004;350:1093-103.
13. Hong SJ, Kim YS, Kim HS. Prevalence and clinical features of hyperuricemia in Gwangju and Jeonnam territories. *J Rheum Dis* 2012;19:138-46.
 14. Kim YB, Jhang WG, Hwangbo Y, Kim HS, Ahn KD, Lee BK, Lee YJ, Lee JE, Lee JS, Lss CG, Lee HJ, Lee SS. Risk factors of hyperuricemia in the citizens of Asan. *J Soonchunhyang Med Coll* 2006;12:41-9.
 15. Kim EH, Jeon K, Park KW, Kim HJ, Ahn JK, Jeon CH, Cha HS, Koh EM. The prevalence of gout among hyperuricemic population. *J Korean Rheum Assoc* 2004;11:7-13.
 16. Hochberg MC, Silman AJ, Smolen JS, Weinblatt ME, Weisman MH. *Rheumatology*. 3th ed. St. Louis (MO): Mosby; 2003.
 17. The Korean Nutrition Society. *Dietary reference intakes for Koreans 2010*. Seoul: The Korean Nutrition Society; 2010.
 18. Poletto J, Harima HA, Ferreira SR, Gimeno SG. Hyperuricemia and associated factors: a cross-sectional study of Japanese-Brazilians. *Cad Saude Publica* 2011;27:369-78.
 19. Choi HK, Atkinson K, Karlson EW, Curhan G. Obesity, weight change, hypertension, diuretic use, and risk of gout in men: the health professionals follow-up study. *Arch Intern Med* 2005;165:742-8.
 20. Lottmann K, Chen X, Schädlich PK. Association between gout and all-cause as well as cardiovascular mortality: a systematic review. *Curr Rheumatol Rep* 2012;14:195-203.
 21. Vučak J, Katić M, Bielen I, Vrdoljak D, Lalić DI, Kranjčević K, Marković BB. Association between hyperuricemia, prediabetes, and prehypertension in the Croatian adult population--a cross-sectional study. *BMC Cardiovasc Disord* 2012;12:117.
 22. Torralba KD, De Jesus E, Rachabattula S. The interplay between diet, urate transporters and the risk for gout and hyperuricemia: current and future directions. *Int J Rheum Dis* 2012;15:499-506.
 23. Kang DH. Does hyperuricemia play a causative role in the development and/or aggravation of renal, cardiovascular and metabolic disease? *Korean J Med* 2011;80:524-8.
 24. Chin HJ, Na KY, Kim Y, Chae DW, Kim S. The impact of uric acid and metabolic syndrome on the incidence of hypertension in a Korean population. *Korean J Med* 2007;73:58-66.
 25. Nakagawa T, Hu H, Zharikov S, Tuttle KR, Short RA, Glushakova O, Ouyang X, Feig DI, Block ER, Herrera-Acosta J, Patel JM, Johnson RJ. A causal role for uric acid in fructose-induced metabolic syndrome. *Am J Physiol Renal Physiol* 2006;290:F625-31.
 26. Sánchez-Lozada LG, Tapia E, Bautista-García P, Soto V, Avila-Casado C, Vega-Campos IP, Nakagawa T, Zhao L, Franco M, Johnson RJ. Effects of febuxostat on metabolic and renal alterations in rats with fructose-induced metabolic syndrome. *Am J Physiol Renal Physiol* 2008;294:F710-8.
 27. Schlesinger N. Dietary factors and hyperuricaemia. *Curr Pharm Des* 2005;11:4133-8.
 28. Choi HK, Gao X, Curhan G. Vitamin C intake and the risk of gout in men: a prospective study. *Arch Intern Med* 2009;169:502-7.
 29. Sutton JL, Basu TK, Dickerson JW. Effect of large doses of ascorbic acid in man on some nitrogenous components of urine. *Hum Nutr Appl Nutr* 1983;37:136-40.
 30. Ghadirian P, Shatenstein B, Verdy M, Hamet P. The influence of dairy products on plasma uric acid in women. *Eur J Epidemiol* 1995;11:275-81.
 31. Choi HK, Liu S, Curhan G. Intake of purine-rich foods, protein, and dairy products and relationship to serum levels of uric acid: the Third National Health and Nutrition Examination Survey. *Arthritis Rheum* 2005;52:283-9.
 32. Wang Y, Yan S, Li C, Zhao S, Lv J, Wang F, Meng D, Han L, Wang Y, Miao Z. Risk factors for gout developed from hyperuricemia in China: a five-year prospective cohort study. *Rheumatol Int* 2013;33:705-10.
 33. Luk AJ, Simkin PA. Epidemiology of hyperuricemia and gout. *Am J Manag Care* 2005;11:S435-42.
 34. Choi HK, Mount DB, Reginato AM; American College of Physicians; American Physiological Society. Pathogenesis of gout. *Ann Intern Med* 2005;143:499-516.
 35. Nicklas TA, O'Neil CE, Fulgoni VL 3rd. Diet quality is inversely related to cardiovascular risk factors in adults. *J Nutr* 2012;142:2112-8.
 36. Lim SY, Yoo HJ, Kim AL, Oh JA, Kim HS, Choi YH, Cho JH, Lee JH, Yoon KH. Nutritional intake of pregnant women with gestational diabetes or type 2 diabetes mellitus. *Clin Nutr Res* 2013;2:81-90.
 37. Drewnowski A, Fiddler EC, Dauchet L, Galan P, Hercberg S. Diet quality measures and cardiovascular risk factors in France: applying the Healthy Eating Index to the SU.VI.MAX study. *J Am Coll Nutr* 2009;28:22-9.