

# Transantral Orbital Floor Fracture Repair Using a Folded Silastic Tube

Joo Yeon Kim · Gwan Choi · Jae Hwan Kwon

*Department of Otolaryngology-Head and Neck Surgery, Kosin University College of Medicine, Busan, Korea*

**Objectives.** The purpose of this study was to evaluate the advantages and limitations of using a silicon tube to support the fractured orbital floor by a transantral approach.

**Methods.** A retrospective study was conducted from January 2000 to December, 2011 in 51 patients with pure orbital floor fractures. The patients underwent reduction surgery via a transantral approach for inserting a folded silastic tube to support the fractured orbital floor in the maxillary sinus. A chart review of preoperative and postoperative ocular symptoms, operation records, and complications was maintained.

**Results.** In 18 out of 25 patients with diplopia, postoperative improvement was seen. In 13 out of 15 patients with extraocular muscle limitation, postoperative improvement was seen. Enophthalmos resolved postoperatively in four of five patients. Postsurgical complications occurred in three patients: an overcorrection, an infection in the maxillary sinus, and an implant extrusion, all of which were resolved by revision surgeries.

**Conclusion.** During the course of the study, we sensed reduction using a folded silastic tube via a transantral approach as an easy and effective technique with good postoperative results, and minimal implant related complications. This novel procedure is recommended as a surgical option for the reduction of orbital floor fractures.

**Keywords.** *Orbital Fractures; Silastic Tube; Transantral Approach*

## INTRODUCTION

Orbital floor fracture is one of the most common facial skeleton fractures after mid facial trauma, accounting for up to 40% of craniofacial injuries [1,2]. Patients may complain of ocular symptoms, aberration of aesthetic appearance, and dysesthesia around the damaged cheek. The goal of treatment is restoration of the anatomic and functional defects of the globe. Recently, numerous approaches for reduction of orbital floor fractures have been proposed including transorbital, transantral, and transnasal using endoscopy [3-6]. Various materials have been introduced to cover the floor defect, or support the prolapsed orbital contents

from the maxillary sinus [5,7,8]. Several alternate materials used to support the orbital floor in the maxillary sinus via a transantral approach, include hydroxyapatite block, silicone block, and balloon and gauze soaked with iodine formaldehyde. Potential problems with these methods are instability, implant displacement, inadequate reduction, infection, and cost [9-11]. We propose the use of a silicon tube for the reduction of blowout fractures of the inferior orbital wall via a transantral approach. The aim of this study was to evaluate the usefulness, advantages and limitations of the proposed technique.

## MATERIALS AND METHODS

### Patients

Fifty-one patients with pure blowout fracture of the inferior orbital wall from January 2000 to December 2011, at the Department of Otolaryngology-Head and Neck Surgery of Maryknoll Medical Center were included in the study. All enrolled patients

• Received August 10, 2014  
 Revised September 19, 2014  
 Accepted September 21, 2014

• Corresponding author: **Jae Hwan Kwon**  
 Department of Otolaryngology-Head and Neck Surgery, Kosin University College of Medicine, 262 Gamcheon-ro, Seo-gu, Busan 602-702, Korea  
 Tel: +82-51-990-6247, Fax: +82-51-245-8539  
 E-mail: entkwon@hanmail.net

Copyright © 2015 by Korean Society of Otorhinolaryngology-Head and Neck Surgery.

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

underwent reduction surgery using folded silicon tube placement via the transantral approach. Clinical indications for fracture repair included persistent diplopia (>2 weeks), extraocular muscle entrapment on face computed tomography (CT) scan, a large posterior fracture (>50% of the orbital floor), or early enophthalmos [5,8,12,13]. Exclusion criteria were as follows: (1) patients with orbital floor fractures, who did not undergo reduction surgery; (2) patients with accompanying ethmoidal, maxillary, or zygomatic bone fractures; (3) insufficient medical data (preoperative and postoperative CT scans, operation, and medical records); (4) patients, who underwent surgery with other implanted materials; (5) patients, who did not follow up for more than three months despite persistent orbital symptoms; (6) patients, who underwent reduction surgery via both transantral and transorbital approaches; (7) patients, who underwent simultaneous reduction surgery for both orbital floor and medial wall fractures. All protocols were approved by the Kosin University Institutional Review Board (IRB No. 12-104).

### Analysis

All patients received a complete, preoperative and postoperative ophthalmic examination for visual acuity, diplopia, extraocular muscle limitation, and enophthalmos. Diplopia in the cardinal fields of gaze was considered significant. Enophthalmos was regarded as significant if a 2-mm discrepancy between eyeballs existed. A Hertel exophthalmometer was used for measure-

ments a day presurgery and 4 weeks postsurgery. Facial CT was performed in all patients, preoperatively and postoperatively. The mean displacement between fractured bone chip and intact orbital floor was based on anatomic positioning on CT scans. Displacement was classified as: grade I, an exact anatomic position of reduced orbital floor or minimal displacement; grade II,




Fig. 1. Illustrations showing the repair of the orbital floor fracture. (A) Note that the fractured fragments of the orbital floor and orbital fat are returned to their original place before the insertion of implants. (B) A silastic sheet (1 mm; arrowhead) is placed under the orbital floor and a bent silicon tube (arrow) is inserted into the maxillary sinus under the silastic sheet via a transantral approach.




Fig. 2. Silastic sheet and tubes used in this technique. (A) Silastic sheet (1 mm in thickness) placed under the fractured orbital floor in the maxillary sinus through the antral window. (B) Silicon tubes (0.5 mm for children, 1 mm for adults) for upholding the fractured orbital floor. (C) Various strategies for bending the silicon tubes.

discrepancy of displacement below 2 mm; and grade III, a discrepancy of displacement above 2 mm. Complications and management strategies were evaluated by chart review.

### Operation technique

Surgical procedures were performed under general anesthesia. Typically, 1% lidocaine with 1:100,000 dilution of epinephrine was injected into the upper gingivobuccal sulcus. A 3- to 4-cm horizontal incision was made just superior to the sulcus. The periosteum and overlying soft tissue were gently elevated from the underlying maxillary bone to the infraorbital foramen using a periosteal elevator. An antral window (2 cm×1.5 cm) was made into the maxillary sinus with an electric saw. The bone fragment was preserved in saline solution for repositioning at the end of surgery. The maxillary sinus and prolapsed orbital contents were visualized employing a 0-degree endoscope, through the antral window. The herniated orbital contents were reduced by digital manipulation or by using surgical instruments like an elevator, without removing the mucosa and fractured bone chip. A silastic sheet (1 mm), slightly larger than the fractured area,

was inserted into the maxillary sinus and placed below the orbital floor. A silicon tubing (1 mm in adult and 0.5 mm in pediatric cases) was bent one or two times, and inserted into the maxillary sinus under the silastic sheet (Fig. 1). When placed into the maxillary sinus, the tube recoiled automatically and adjusted to the sinus wall to firmly support the reduced orbital floor with its elasticity (Fig. 2). Care was taken not to obstruct the natural ostium of the maxillary sinus (Fig. 3). A forced duction test was performed to confirm correct positioning of the orbital floor and to avoid entrapment of the orbital contents before closing. The antral wall fragment, stored in saline, was fixed in its original position using a plate and screws. The gingivobuccal incision was closed, and postoperative antibiotics were continued for 14 days. The silastic sheet and tube were removed 6 to 8 weeks postsurgery under local anesthesia through the same antral maxillary window, created during the first surgery (Fig. 4). The second surgery was performed under general anesthesia in all children and some adults.




Fig. 3. Intraoperative endoscopic views of the left maxillary sinus of an orbital floor fracture, through the antral window. (A) The orbital floor was fractured, and periorbital soft tissue was herniated into the maxillary sinus (arrow). (B) The fractured orbital contents were reduced to anatomic position and a silastic sheet was placed under the fractured bone. (C) Bent silastic tubing was inserted into the maxillary sinus to support the reduced orbital floor. (D) The silastic tube and sheet were removed, and the fractured orbital floor was well reduced (arrow, natural ostium).



Fig. 4. Coronal computed tomography (CT) scans of the blowout fracture of the inferior orbital wall. (A) Preoperative view. (B) Postoperative CT scan before silicone tube removal, the inferior orbital wall was supported with the silastic sheet and silicon tube. (C) Well reduced state of fractured bony wall after removal of silicon tube, two years postsurgery.

## RESULTS

Among the 51 patients included, 39 were male and 12 were female. The mean age of the patients was 31 years (range, 12 to 69 years). Trauma occurred on the right side in 24, the left side in 26, and on both sides in one patient. The patient with both orbital wall fractures had surgery performed through the more grossly damaged side. Forty-three patients had inferior orbital wall fractures. Eight patients with combined inferior and medial orbital wall fractures underwent reduction surgery for orbital floor fractures only, because of disregarded medial wall fractures. The most common causes of fracture were falls, assaults, sports injuries, and industrial accidents. The mean interval between injury and repair was 8.5 days (range, 2 to 35 days) and the mean length of follow-up was 87 days (range, 28 to 401 days).

### Ophthalmic symptoms

Twenty-five patients reported diplopia before surgery, of which postoperative improvement was seen in 18 patients. In seven patients, diplopia persisted at extreme upper gaze postsurgery, but disappeared within the 3-month follow-up period. Thirteen out of 15 patients had improved extraocular muscle limitation postoperatively. Enophthalmos resolved postoperatively in four of five patients (Table 1).

### Discrepancy between fractured bone chip and orbital floor based on CT scan

Preoperatively, there were three cases of grade II and 48 cases of grade III displacement. All three preoperative grade II cases were restored to grade I postoperatively. Among 48 patients with a preoperative status of grade III, 16, 25, and seven cases had grades I, II, III, respectively, postoperatively (Table 2).

Table 1. Ocular symptoms before and after surgery

Ocular symptom	Preoperative	Postoperative
Diplopia	25	7
EOM limitation	15	2
Enophthalmos	5	1

EOM, extraocular muscle movement.

Table 2. Mean discrepancy between fractured bone chip and intact orbital floor based on coronal computed tomography scans (n=51)

Grade	Preoperative	Postoperative
I: Exact anatomic position of reduced orbital floor or minimal displacement	0	16 (31.4)
II: Discrepancy of displacement <2 mm	3 (5.9)	28 (54.9)
III: Discrepancy of displacement >2 mm	48 (94.1)	7 (13.7)

Values are presented as number (%).

Table 3. Postoperative complications and management strategies

Complication	Management strategy
Overcorrection (n=1)	Reposition of silicone tube
Implant displacement (n=1)	Reposition of silicone tube
Sinusitis (n=1)	Silicone tube removal

### Complications and management

Inflammatory changes of maxillary sinus mucosa were observed in 21 patients upon silicone tube removal. The inflamed mucosa was either shaved using a microdebrider after removal of the implants, or left undisturbed. One patient complained of severe diplopia, ocular pain, and aggravated extraocular muscle limitation after surgery. The symptoms were attributed to overcorrection of the fractured bony fragments, and were alleviated after immediate revision surgery. A patient, 3 days postsurgery, presented extrusion of the silicon tube through the maxillary sinus window. Corrective implant repositioning was achieved by re-

sion surgery. Another patient developed maxillary sinusitis at 4 weeks postsurgery, with symptoms of cheek pain, swelling, and nasal discharge. The sinusitis symptoms resolved on removal of the implant. These results are summarized in Table 3.

## DISCUSSION

The orbital floor can be accessed through a transorbital, transantral, or endoscopic transnasal approach [3-6]. The transorbital approach has a limited ability to restore large floor defects and access the posterior orbital region; it also leads to unsightly facial scars or eyelid complications [14,15]. The transantral approach is clinically useful in cases involving severe fractures or posterior floor fractures; it has the advantage of providing a direct view of the defect size and configuration, and preventing injury to lower eyelid structures. However, the transantral approach can lead to entrapment of the extraocular muscle in trapdoor-type fractures, and lead to infraorbital nerve injury [10,16,17]. Recently, an endoscopic endonasal approach for orbital floor fracture was introduced [5]. However, it is only suitable to limited cases of medial or posterior orbital floor fractures.

Reports on the transantral approach, have discussed various materials for supporting the fractured orbital floor and filling up the maxillary sinus. These materials include iodophor gauze, hydroxyl appetite block scaffolding, silicone block, and sinus balloons [7,9-11,18]. However, the techniques of implant placement within the maxillary sinus to reduce fractured orbital floors are limited due to the difficulty in restoring orbital floor anatomic positioning. Furthermore, these techniques increase the risks of extrusion, migration, and infection. The procedure can be prohibitively expensive, and requires a second surgery for implant removal [9,10,19]. These limitations have restricted the widespread use of transantral approach.

In this study, we used folded silicon tubing to uphold the fractured orbital floor in the maxillary sinus. Implanting bent silicon tubes provided several benefits: (1) Silicon tubes were bent and easily fitted into the maxillary sinus, thus shortening operation times; (2) The elasticity of the silicon tube maintained the orbital floor in a stable anatomic location until implant removal; (3) A more physiological healing was attained because the sinus mucosa and fractured orbital bone chips were not removed; (4) The material is inexpensive; and (5) It is suitable for addressing relatively large and posterior orbital floor fractures. The shortcomings of this technique are as follows: (1) Secondary surgery is needed to remove the implant (note that the surgery can be performed easily under local anesthesia through the previously constructed maxillary window); and (2) Transient inflammatory changes of the maxillary sinus mucosa can occur.

Postoperatively, diplopia improved but persisted with maximum upward gaze in seven of the 25 patients. All diplopia symptoms resolved within the 3-month follow-up. Similarly, Folkestad

and Westin [9] reported 38% postoperative diplopia in patients, who underwent operative repair using balloon and iodine formaldehyde soaked gauze via a transantral approach. Four of the five patients with enophthalmos, experienced resolution postsurgery. Ikeda et al. [5] reported two cases of postoperative enophthalmos of the 11 surgically treated cases of orbital floor fractures. Even though this study included treatments of more challenging posterior orbital bone fractures and large fractures, our results showed acceptable outcomes.

This surgery has physiological advantages, with respect to that it not only reduces periorbital soft tissue but also restores fractured bony fragments and mucosa to their original anatomic positions. Thus, overall, there is more stable restoration and a decrease in ophthalmic complications.

Entrapment of periorbital soft tissue can occur, in trapdoor-type fractures, but can be resolved by employing a combination transorbital approach [20].

Twenty-one patients in the current study had mild polypoid changes in the maxillary sinus mucosa on implant removal. This postoperative inflammatory change was not significantly different from that which occurs with other filler materials (e.g., hydroxyapatite block, silicon block, or gauze soaked with iodine formaldehyde) implanted via a transantral approach [9-11], and the inflammation was supposed to be spontaneously resolved after implant removal.

There were three cases with complications that were successfully treated by revision surgery. One female patient (on no antibiotics) had acute sinusitis, which resolved after implant removal. The second case involved extrusion of implants. The third complication case was overcorrection, which occurred among the early surgeries, when surgeons were still perfecting this technique.

In summary, using folded silicon tubing to repair orbital floor fractures can yield exact positioning of fractured bone and promote physiological healing. It is an easy, economical, and safe procedure compared to previously used implant materials for transantral procedures. It is proposed that this procedure can be a novel surgical option for orbital floor fracture reduction.

## CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

## ACKNOWLEDGMENTS

This study was supported by a grant from Kosin University College of Medicine (2014).

## REFERENCES

1. Jank S, Schuchter B, Emshoff R, Strobl H, Koehler J, Nicasi A, et al. Clinical signs of orbital wall fractures as a function of anatomic location. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2003 Aug; 96(2):149-53.
2. Manolidis S, Weeks BH, Kirby M, Scarlett M, Hollier L. Classification and surgical management of orbital fractures: experience with 111 orbital reconstructions. *J Craniofac Surg.* 2002 Nov;13(6):726-37.
3. Lee HH, Alcaraz N, Reino A, Lawson W. Reconstruction of orbital floor fractures with maxillary bone. *Arch Otolaryngol Head Neck Surg.* 1998 Jan;124(1):56-9.
4. Lai A, Gliklich RE, Rubin PA. Repair of orbital blow-out fractures with nasoseptal cartilage. *Laryngoscope.* 1998 May;108(5):645-50.
5. Ikeda K, Suzuki H, Oshima T, Takasaka T. Endoscopic endonasal repair of orbital floor fracture. *Arch Otolaryngol Head Neck Surg.* 1999 Jan;125(1):59-63.
6. Walter WL. Early surgical repair of blowout fracture of the orbital floor by using the transantral approach. *South Med J.* 1972 Oct; 65(10):1229-43.
7. Gray LN, Kalimuthu R, Jayaram B, Lewis N, Sohaey M. A retrospective study of treatment of orbital floor fractures with the maxillary sinus approach. *Br J Plast Surg.* 1985 Jan;38(1):113-5.
8. Lemke BN, Kikkawa DO. Repair of orbital floor fractures with hydroxyapatite block scaffolding. *Ophthal Plast Reconstr Surg.* 1999 May;15(3):161-5.
9. Folkestad L, Westin T. Long-term sequelae after surgery for orbital floor fractures. *Otolaryngol Head Neck Surg.* 1999 Jun;120(6): 914-21.
10. Converse JM, Smith B, Obear MF, Wood-Smith D. Orbital blowout fractures: a ten-year survey. *Plast Reconstr Surg.* 1967 Jan;39(1): 20-36.
11. Courtney DJ, Thomas S, Whitfield PH. Isolated orbital blowout fractures: survey and review. *Br J Oral Maxillofac Surg.* 2000 Oct;38(5): 496-504.
12. Brady SM, McMann MA, Mazzoli RA, Bushley DM, Aimbinder DJ, Carroll RB. The diagnosis and management of orbital blowout fractures: update 2001. *Am J Emerg Med.* 2001 Mar;19(2):147-54.
13. Saunders CJ, Whetzel TP, Stokes RB, Wong GB, Stevenson TR. Transantral endoscopic orbital floor exploration: a cadaver and clinical study. *Plast Reconstr Surg.* 1997 Sep;100(3):575-81.
14. Whitehouse RW, Batterbury M, Jackson A, Noble JL. Prediction of enophthalmos by computed tomography after 'blow out' orbital fracture. *Br J Ophthalmol.* 1994 Aug;78(8):618-20.
15. Raskin EM, Millman AL, Lubkin V, della Rocca RC, Lisman RD, Mather EA. Prediction of late enophthalmos by volumetric analysis of orbital fractures. *Ophthal Plast Reconstr Surg.* 1998 Jan;14(1):19-26.
16. Otori N, Haruna S, Moriyama H. Endoscopic endonasal or transmaxillary repair of orbital floor fracture: a study of 88 patients treated in our department. *Acta Otolaryngol.* 2003 Aug;123(6):718-23.
17. Persons BL, Wong GB. Transantral endoscopic orbital floor repair using resorbable plate. *J Craniofac Surg.* 2002 May;13(3):483-8.
18. Ortiz Monasterio F, Rodriguez A, Benavides A. A simple method for the correction of enophthalmos. *Clin Plast Surg.* 1987 Jan;14(1): 169-75.
19. Nishiike S, Nagai M, Nakagawa A, Konishi M, Kato T, Sakata Y, et al. Endoscopic transantral orbital floor repair with antral bone grafts. *Arch Otolaryngol Head Neck Surg.* 2005 Oct;131(10):911-5.
20. Kwon JH, Kim JG, Moon JH, Cho JH. Clinical analysis of surgical approaches for orbital floor fractures. *Arch Facial Plast Surg.* 2008 Jan-Feb;10(1):21-4.