

Original Article



Altered Sphingolipid Metabolism Is Associated With Asthma Phenotype in House Dust Mite-Allergic Patients

Krzysztof Kowal,^{1,2*} Ewa Żebrowska,³ Adrian Chabowski³

¹Department of Experimental Allergology and Immunology, Medical University of Białystok, Białystok, Poland

²Department of Allergology and Internal Medicine, Medical University of Białystok, Białystok, Poland

³Department of Physiology, Medical University of Białystok, Białystok, Poland



Received: Sep 2, 2018

Revised: Nov 18, 2018

Accepted: Dec 11, 2018

Correspondence to

Krzysztof Kowal MD, PhD

Department of Experimental Allergology and Immunology, Medical University of Białystok, Skłodowskiej-Curie 24A, 15-276 Białystok, Poland.

Tel: +48-85-6865153

Fax: +48-85-6865152

E-mail: kowalkmd@umb.edu.pl

Copyright © 2019 The Korean Academy of Asthma, Allergy and Clinical Immunology · The Korean Academy of Pediatric Allergy and Respiratory Disease

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclosure

There are no financial or other issues that might lead to conflict of interest.

ABSTRACT

Purpose: Sphingolipids play an important role in cell growth, survival, inflammation and tissue remodeling. House dust mite (HDM) allergy is a major risk factor for asthma. The aim of the study was to evaluate if allergic asthma phenotype is associated with altered sphingolipid metabolism.

Methods: Twenty-two HDM-allergic asthmatic patients and 11 HDM-allergic rhinitis patients were challenged intrabronchially with biologically standardized *Dermaphagoides pteronyssinus* extract. Whole blood and platelet-poor plasma samples were collected before, during early asthmatic response (EAR), late asthmatic response (LAR) and 24 hours after the challenge. Concentrations of sphinganine (SFA), sphinganine-1-phosphate (SFA1P), ceramide, sphingosine (SFO) and sphingosine-1-phosphate (S1P) were measured using high performance liquid chromatography.

Results: In all house dust mite-allergic patients (HDM-APs), baseline lung function and severity of airway hyperreactivity (AHR) correlated significantly with plasma S1P and SFA1P concentrations. Exhaled nitric oxide concentration, however, correlated with SFA and ceramide, but not with S1P or SFA1P concentration. Allergen challenge increased plasma S1P concentration during EAR, but only in patients who developed both EAR and LAR. The magnitude of the increase determined during EAR correlated with the severity of subsequently developed LAR. Platelet and eosinophil counts were independent predictors of plasma S1P concentration. A significant increase in plasma SFA concentration in response to allergen challenge was seen only in patients who did not develop asthmatic response.

Conclusions: Altered sphingolipid metabolism, with augmented synthesis of S1P and impaired *de novo* sphingolipid synthesis in response to allergen challenge, may participate in the development of asthma phenotype in HDM-APs.

Keywords: Lipid mediators; asthma; inflammation

INTRODUCTION

Asthma is a heterogeneous disorder characterized by chronic airway inflammation, tissue remodeling and airway hyperreactivity (AHR).¹ Based on common clinical presentation as well as on epidemiology, genetics, biomarkers, lung physiology, histopathology and response to therapy, different asthma endotypes have been proposed.¹ Allergic asthma is the most

frequent asthma endotype and house dust mites (HDMs) allergy is the most frequent among asthmatic patients, which is also one of the greatest risk factors for asthma and AHR.²⁻⁴ Yet, a substantial number of house dust mite-allergic patients (HDM-APs) do not suffer from asthma and have normal bronchial reactivity.^{5,6}

Many mediators and metabolic pathways are potentially involved in the development of chronic airway inflammation, airway remodeling and AHR in allergic patients.^{7,8} Recently, sphingolipids and enzymes involved in sphingolipid metabolism have been proposed to play a role in the pathogenesis of allergic diseases, AHR and asthma.^{9,12} Sphingolipids play an important role in cell growth, survival, inflammation and tissue remodeling.^{13,14} Sphingosine-1-phosphate (S1P) is of particular interest due to its proinflammatory and anti-apoptotic effects.^{9,10} Experimental models of allergic asthma demonstrated that S1P played a crucial role in the development of asthmatic phenotype in different experimental models in mice.^{15,16} Two subcutaneous injections of S1P, 7 days apart resulted in progressively developed airway inflammation and AHR reaching maximum 2 weeks after the last dose.¹⁵ Moreover, local application of exogenous S1P aggravated allergen-induced airway inflammation in a mouse asthma model.¹⁶ On the contrary, inhibition of sphingosine kinase (SPHK) 1, which is crucial for synthesis of S1P, led to attenuated airway inflammation and AHR in a model of allergic asthma.¹⁷ Similarly, application of S1P analogue, which interfered with S1P-S1P receptor interaction, inhibited airway remodeling following repeated allergen exposure in rats.¹⁸

Human studies also provided some evidence for a potential involvement of S1P in the pathogenesis of asthma.^{19,20} An increase in concentration of S1P was demonstrated in bronchoalveolar lavage (BAL) fluid 24 hours after segmental allergen challenge.¹⁹ *In vitro*, S1P directly induced pro-remodeling response in human airway smooth muscle cells.²⁰ An S1P plasma-interstitial fluid gradient exists and inflammation-driven increased vessel permeability, as seen in the airways of asthmatic patients, may result in additional influx of S1P to tissue compartment.²¹

Other sphingolipids also affect processes involved in cell growth, survival and apoptosis.^{12,44} Interestingly, activation of *de novo* sphingolipid synthesis leads to production of sphinganine (SFA) and ceramide which exert opposite to S1P effects.^{12,44} Therefore, the balance between sphingolipid synthesis and degradation is important for regulation of cell growth, survival, inflammation and tissue remodeling.^{12,44}

In the current study, we evaluated the effect of bronchial allergen challenge on plasma concentration of selected sphingolipids in a well characterized group of HDM-APs.

MATERIALS AND METHODS

The study was performed on 33 HDM-APs. All patients reported rhinitis symptoms, while 22 patients reported also asthma symptoms upon exposure to house dust. Sensitization to the HDM components *Dermatophagoides pteronyssinus* (*Dp*) and *Dermatophagoides farinae* (*Df*) was confirmed by demonstration of positive skin prick tests and elevated (>0.7 kU/L) serum concentrations of anti-*Dp* specific immunoglobulin E (IgE). Before the initial visit, none of the patients received allergen immunotherapy or any anti-asthma medication, except sporadic application of short-acting-beta agonists.

The study was approved by the local Ethics Committee (R-I-003/131/2004). All participants provided written informed consent.

Pulmonary function tests

Histamine bronchial challenge was performed as previously described.²² All patients inhaled doubling concentrations of histamine starting from a concentration of 0.125 mg/mL. Forced expiratory maneuvers were performed 90 seconds after fifth inhalation of each histamine concentration. The procedure was continued until either at least a 20% reduction in forced expiratory volume during the first second of expiration (FEV1) or a histamine concentration of 32 mg/mL was reached. Nonspecific bronchial reactivity was expressed as histamine concentration causing 20% fall in FEV1 (PC20).

Bronchial provocation test with aqueous *Dp* extracts (Allergopharma, Germany) were performed as described before.²² Increasing doses of allergen (0.8, 4, 20, 100, 500 and 2,500 SBU) were administered using a De Vilbiss#646 nebulizer attached to a Rosenthal-French dosimeter. Forced expiratory maneuvers were performed 15 minutes after inhalation of each dose of the allergen extract. Allergen inhalations were continued until either at least a 20% reduction in FEV1 (PD20) or a cumulative dose of 5,000 SBU was reached. Subsequently, FEV1 was measured every 15 minutes during the first hour after challenge, every 60 minutes during the next 11 hours and after 24 hours. Specific bronchial reactivity was expressed as the allergen dose causing a PD20. Bronchial challenge with *Dp* allergen extract was performed on all patients sensitive to HDM allergens.

Exhaled nitric oxide (NO) measurements

Concentration of NO in the exhaled air was measured using a chemiluminescence analyzer NOA 280i (Sievers Instruments, Boulder, CO, USA) according to ATS recommendations as described elsewhere.²² Briefly, each patient exhaled against a fixed expiratory resistance of 16 cm H₂O resulting in a constant flow of 50 mL/s. A plateau of NO concentration in the exhaled air at the selected exhalation rate was automatically selected by the computer software. NO measurements were repeated 3 times and the mean value was used for analysis.

Blood samples

Plasma samples were obtained using citrate-theophylline-adenosine-dipyridamole (CTAD) anticoagulation as previously described.²² In addition, EDTA-anticoagulated samples were collected for assessment of complete blood count. The CTAD-anticoagulated blood samples were incubated on ice for 30 minutes and then plasma was separated by centrifugation at 4°C. The supernatants containing platelet poor plasma were aliquoted and stored at -80°C until tested. The samples were collected before bronchial allergen challenge (T_0), at 45 minutes (T_{EAR}), 6-8 hours (T_{LAR}) and 24 hours (T_{24}) after administration of the last allergen dose.

Biochemical and immunologic assays

Total IgE and specific IgE were measured in the serum samples using the UniCap system (Pharmacia, Uppsala, Sweden).

Complete blood count, including red blood cell (RBC) and platelet counts, as well as white blood cell (WBC) differential was measured using automated hematological analyzer ADVI-120 (Bayer, Leverkusen, Germany).

Total cholesterol (TC) and high-density lipoprotein cholesterol (HDL-C) were measured using Abbott Diagnostics (Wiesbaden, Germany) on the Architect c8000 analyzer (Abbott Laboratories, Abbott Park, IL, USA).

The levels of ceramide, sphingosine (SFO), SFA, S1P and sphinganine-1-phosphate (SFA1P) were determined as previously described.²³ Briefly, lipids were extracted from 250 µL of plasma in the presence of internal standards (10 pmol C17-sphingosine and 30 pmol C17-S1P, Avanti Polar Lipids). An aliquot of the lipid extract was transferred to a fresh tube with preadded 40 pmol N-palmitoyl-Derythro-sphingosine (C17 base; a kind gift of Dr. Z. Szulc, Medical University of South Carolina) as an internal standard, and then subjected to alkaline hydrolysis to deacylate ceramide to SFO. The amounts of S1P and SFA1P were determined indirectly after dephosphorylation to SFO and SFA, respectively, with the use of alkaline phosphatase (bovine intestinal mucosa, Fluka). Free SFO and SFA, dephosphorylated sphingoid bases and SFO released from ceramide were then converted to their o-phthalaldehyde derivatives and analyzed using a HPLC system equipped with a fluorescence detector and C18 reversed-phase column (Varian Inc., OmniSpher 5; 4.6 × 150 mm). The isocratic eluent composition of acetonitrile (Merck, Darmstadt, Germany): water (9:1, v/v) and a flow rate of 1 ml/min were used. All measurements were taken in a blind fashion.

Statistical analysis

Data were analyzed by Student's *t* test for independent means. Repeated measurements were analyzed using Student's *t* test for dependent means. For multiple comparisons the Bonferroni correction was applied. Values of $P < 0.05$ were considered significant. All results are expressed as the means with 95% confidence intervals (CIs). Linear correlation by Pearson was used to estimate correlations between studied parameters. Multiple regression analysis was used to estimation of predictors of individual sphingolipid plasma concentration.

RESULTS

All patients ($N = 33$) who were allergic to HDM allergens reported rhinitis triggered by exposure to house dust. Twenty-two patients also reported asthma symptoms upon exposure to house dust. The patients developed asthmatic response after bronchial challenge with *Dp* extract (responders; Rs). The patients who did not complain of asthma symptoms upon house dust exposure ($n = 11$) also did not develop asthmatic response after bronchial challenge with *Dp* extract (non-responders; NRs). There was no difference in the demographic characteristics between the 2 groups (**Table 1**). However, Rs were characterized by greater AHR, peripheral blood eosinophilia and specific IgE concentration. Eleven of the Rs (single responders; SRs) developed isolated early asthmatic response (EAR), while the remaining 11 patients (dual responders; DRs) developed both EAR and LAR after the challenge with *Dp* extract. The level of sphingolipids examined in our study was in the range previously demonstrated among healthy subjects.²⁴

Elevated plasma S1P and SFA1P concentrations are associated with AHR

There was no significant difference between the mean concentration of TC or HDL-C between Rs and NRs (**Table 1**). In Rs, the mean baseline concentrations of S1P (179 pmol/mL) and SFA (5.9 pmol/mL) were significantly greater than those in NRs (129 and 3.4 pmol/mL, $P = 0.0068$ and $P = 0.0005$, respectively) (**Table 1**). No significant difference in the concentration of ceramide ($P = 0.509$), SFO ($P = 1.0$) or SFA1P ($P = 0.07$) between Rs and NRs was demonstrated

Table 1. Patient characteristics

Characteristics	NRs	Rs	P value
Sex (M/F)	5/6	10/12	1.0
Age (years)	28.3 (21.4–35.2)	25.1 (21.9–28.2)	0.567
BMI (kg/m ²)	23.2 (21.3–25.2)	23.8 (22.3–25.2)	0.89
TC (mg/dL)	171 (153–189)	173 (163–184)	0.791
HDL-C (mg/dL)	48 (43–52)	51 (46–56)	0.208
Total IgE (kU/L)	180 (114–284)	349 (238–510)	0.039
Dp IgE (kU/L)	5.5 (2.9–10.2)	21 (12.3–35.9)	0.006
FEV1 (% predicted)	111.4 (99.8–123.1)	104.8 (100.2–109.3)	0.252
MEF (% predicted)	100.3 (81.9–118.7)	85.6 (76.1–95.1)	0.136
PC20 (mg/mL)	16.3 (4.98–53)	2.1 (1.32–3.37)	0.0039
PD20 (SBU)	>5,000	1,222 (837–1,783)	NA
FeNO (ppb)	47 (30–75)	100 (75–134)	0.0056
Eosinophils (cells/ μ L)	181 (116–246)	366 (269–463)	0.012
Neutrophils (cells/ μ L)	3,150 (2,402–3,918)	3,576 (3,131–4,020)	0.29
Monocytes (cells/ μ L)	379 (324–434)	424 (373–475)	0.257
Lymphocytes (cells/ μ L)	1,995 (1,731–2,258)	1,900 (1,763–2,218)	0.98
Platelets ($\times 10^3$ cells/ μ L)	210 (175–245)	238 (213–262)	0.18
RBC ($\times 10^6$ cells/ μ L)	4,848 (4,596–5,101)	4,873 (4,714–5,032)	0.856
Ceramide (pmol/mL)	3,388 (2,933–3,843)	3,643 (3,126–4,160)	0.509
S1P (pmol/mL)	129 (107–151)	179 (156–202)	0.0068
SFO (pmol/mL)	26.9 (21.9–31.7)	26.9 (22.9–30.7)	1.0
SFA (pmol/mL)	3.4 (2.47–4.35)	5.9 (5.08–6.73)	0.0005
SFA1P (pmol/mL)	16.1 (12.7–19.4)	23.4 (17.9–28.98)	0.07

The results are presented as means with 95% CI. Geometric means are presented for total IgE, DpIgE, PC20, PD20 and FeNO.

NRs, non-responders; Rs, responders; BMI, body mass index; TC, total cholesterol; HDL-C, high-density lipoprotein cholesterol; IgE, immunoglobulin E; Dp, *Dermatophagoides pteronyssinus*; FEV1, forced expiratory volume in 1 second; MEF, maximal expiratory flow; PC20, histamine concentration causing 20% fall in FEV1; PD20, allergen dose causing a 20% reduction in FEV1; FeNO, nitric oxide fraction in exhaled air; RBC, red blood cell; S1P, sphingosine-1-phosphate; SFO, sphingosine; SFA, sphinganine; SFA1P, sphinganine-1-phosphate; CI, confidence interval.

(**Table 1**). Analysis of individual clinical and functional parameters revealed significant associations between plasma concentration of some sphingolipids and lung function parameters. When HDM-APs were divided into 2 groups according to the AHR, significant differences in plasma S1P and SFA1P concentrations were demonstrated. Those with histamine PC20 \leq 4 mg/mL had significantly greater baseline plasma concentrations of S1P (192 pmol/mL) and SFA1P (27.3 pmol/mL) than those without (126 and 13.4 pmol/mL, $P < 0.001$ and $P < 0.001$, respectively) (**Fig. 1**). In those groups, however, no difference in the mean baseline plasma concentration of ceramide ($P = 0.217$), SFA ($P = 0.118$) or SFO ($P = 0.979$) was demonstrated.

The baseline plasma concentration of S1P correlated with the baseline lung function expressed as FEV1 ($r = -0.563$; $P = 0.001$) or MEF ($r = -0.622$; $P < 0.001$) (**Table 2**). Similarly, a significant correlation was demonstrated between the baseline SFA1P and FEV1 ($r = -0.477$; $P = 0.005$) or MEF ($r = -0.485$; $P = 0.004$). Also, AHR significantly correlated with plasma S1P ($r = -0.7451$; $P < 0.001$) and SFA1P ($r = -0.5782$; $P < 0.001$) concentration (**Table 2**). No correlation of the baseline lung function and plasma concentration of any other sphingolipid could be demonstrated.

Interestingly, the baseline FeNO correlated neither with S1P ($r = 0.307$; $P = 0.0824$) nor with SFA1P ($r = 0.242$; $P = 0.176$) plasma concentration (**Table 2**). However, significant correlations between FeNO and the baseline plasma ceramide ($r = 0.505$; $P = 0.0027$) and SFA ($r = 0.605$; $P < 0.001$) concentrations were demonstrated. In contrast, peripheral blood eosinophilia correlated with baseline plasma S1P ($r = 0.592$; $P < 0.001$) and SFA1P ($r = 0.518$; $P = 0.002$), but not with plasma ceramide ($r = 0.278$; $P = 0.117$) or SFA ($r = 0.338$; $P = 0.08$) concentration (**Table 2**).




Fig. 1. Plasma concentrations of selected sphingolipids in relation to the severity of AHR in HDM-APs. PC20 ≤ 4; patients with histamine PC20 equal to or less than 4 mg/mL; PC20 > 4; patients with histamine PC20 greater than 4 mg/mL. Only significant differences were indicated. AHR, airway hyperreactivity; HDM-AP, house dust mite-allergic patient; PC20, histamine concentration causing 20% fall in forced expiratory volume in 1 second.

Table 2. Correlations between individual plasma sphingolipid concentrations and selected functional and biochemical parameters at T0

Parameters	S1P	SFA1P	Ceramide	SFA	SFO
FEV1	-0.563 (-0.76 to -0.273) P = 0.001	-0.477 (-0.705 to -0.159) P = 0.005	-0.113 (-0.44 to 0.239)	-0.332 (-0.628 to 0.022)	-0.286 (-0.573 to 0.064)
MEF	-0.622 (-0.796 to -0.355) P < 0.001	-0.485 (-0.71 to -0.17) P = 0.004	-0.096 (-0.422 to 0.26)	-0.337 (-0.631 to 0.027)	-0.214 (-0.519 to 0.139)
Log _{FeNO}	0.307 (-0.041 to 0.588)	0.242 (-0.111 to 0.54)	0.505 (0.196 to 0.723) P = 0.003	0.605 (0.33 to 0.785) P < 0.001	0.229 (-0.125 to 0.53)
Log _{PC20}	-0.745 (-0.877 to -0.509) P < 0.001	-0.578 (-0.769 to -0.293) P < 0.001	0.012 (-0.698 to 0.39)	-0.348 (-0.643 to 0.038)	-0.304 (-0.613 to 0.086)
Eosinophils	0.592 (0.312 to 0.777) P < 0.001	0.518 (0.212 to 0.731) P = 0.002	0.278 (-0.072 to 0.567)	0.338 (-0.006 to 0.611)	0.294 (-0.055 to 0.579)
Platelets	0.543 (0.246 to 0.747) P < 0.001	0.493 (0.181 to 0.715) P = 0.004	0.019 (-0.326 to 0.36)	0.113 (-0.24 to 0.439)	0.338 (-0.06 to 0.618)
RBC	-0.052 (-0.389 to 0.296)	-0.133 (-0.455 to 0.221)	0.195 (-0.159 to 0.505)	0.324 (-0.11 to 0.608)	0.133 (-0.22 to 0.456)
Neutrophils	0.053 (-0.296 to 0.389)	0.014 (-0.33 to 0.356)	0.219 (-0.135 to 0.523)	0.132 (-0.222 to 0.454)	0.148 (-0.206 to 0.468)
Lymphocytes	0.13 (-0.224 to 0.453)	0.05 (-0.298 to 0.387)	0.275 (-0.075 to 0.565)	-0.159 (-0.476 to 0.195)	0.135 (-0.219 to 0.457)
Monocytes	0.123 (-0.23 to 0.448)	-0.077 (-0.41 to 0.273)	0.089 (-0.262 to 0.42)	0.052 (-0.296 to 0.389)	0.069 (-0.281 to 0.403)

Bold-faced values are considered significant correlations.

S1P, sphingosine-1-phosphate; SFA1P, sphinganine-1-phosphate; SFA, sphinganine; SFO, sphingosine; FEV1, forced expiratory volume in 1 second; MEF, maximal expiratory flow; FeNO, nitric oxide fraction in exhaled air; PC20, histamine concentration causing 20% fall in FEV1; RBC, red blood cell.

Since ceramide and S1P exert opposite effects on regulation of cell growth, survival, inflammation and tissue remodeling analysis of associations of plasma ceramide/S1P ratio with clinical, functional and immunological parameters were also performed (data not shown). The plasma ceramide/S1P ratio was significantly lower in HDM-APs with histamine PC20 ≤ 4 mg/mL (18.2) than in those without ($r = 31.5$, $P < 0.001$). Moreover, the baseline plasma ceramide/S1P ratio significantly correlated with histamine PC20 ($r = 0.449$, $P = 0.019$) and baseline lung function expressed as FEV1 ($r = 0.351$, $P = 0.045$) or MEF ($r = 0.437$, $P = 0.011$) (data not shown).

Peripheral blood eosinophil and platelet counts are independent predictors of S1P plasma concentration

Analysis of the cellular composition of blood samples revealed that the baseline concentration of both S1P and SFA1P correlated with eosinophil ($P = 0.0003$ and $P = 0.002$, respectively) and platelet ($P = 0.0035$ and $P = 0.0036$, respectively) count, but not with RBC, neutrophil, lymphocyte or monocyte count (**Table 2**). No significant correlation between the baseline plasma concentration of SFA/SFO or ceramide and any of the cellular blood component was demonstrated (**Table 2**).

Multiple regression analysis of individual cellular blood components as possible determinants of plasma sphingolipids indicated that peripheral blood eosinophilia ($P = 0.003$) and platelet count ($P = 0.010$) were independent predictors of the baseline plasma S1P concentration. Similarly, concentration of SFA1P was determined independently by peripheral blood eosinophilia ($P = 0.008$) and platelet count ($P = 0.009$). Concentration of other sphingolipids, including ceramide ($P = 0.22$), SFO ($P = 0.29$) and SFA ($P = 0.19$), could not be predicted based on cellular blood components in the population of HDM-APs (data not shown).

Allergen challenge affects plasma S1P concentration

In DRs, the appearance of EAR was associated with an increase in plasma S1P concentration (from 168 to 215 pmol/mL, $P = 0.04$), but a decrease in SRs (from 190 to 132 pmol/mL; $P = 0.004$) (**Fig. 2**). Moreover, at T_{EAR} an increase in plasma S1P concentration was seen in each DR (100%), while in each SR (100%) a decrease in plasma S1P concentration was demonstrated (not shown). Subsequently, at T_{LAR} , plasma S1P concentration decreased in DRs, while in SRs remained at a decreased level seen at T_{EAR} (**Fig. 2**). At T_{24} , plasma S1P concentration in SRs but not in DRs was still lower than the baseline value (**Fig. 2**). No effect of allergen challenge on plasma S1P concentration was demonstrated in NRs (**Fig. 2**).

Plasma concentration of SFA, SFA1P, SFO or ceramide did not change significantly during EAR or LAR. However, at T_{24} plasma concentration of SFA increased in NRs (4.54 pmol/mL, $P = 0.04$), but did not change significantly in SRs (5.4 pmol/mL; $P = 0.55$) or DRs (5.7 pmol/mL, $P = 0.5$) (**Fig. 2**). Interestingly, the changes in S1P plasma concentration during EAR correlated with the severity of subsequently developed LAR ($r = 0.596$, $P = 0.0071$) (**Fig. 3**).

DISCUSSION

Our study for the first time demonstrated that in HDM-APs, altered intravascular sphingolipid metabolisms, including their phosphorylation and *de novo* synthesis, are associated with impaired lung function and AHR. In addition, we show that circulating platelets and eosinophils are major determinants of plasma S1P and that SFA1P concentration and allergen-induced asthmatic responses lead to further alterations in intravascular sphingolipid metabolism. An association between asthma and altered sphingolipid metabolism has already been reported.^{19,25} Greater serum baseline concentrations of S1P and SFO were demonstrated in aspirin-exacerbated respiratory disease (AERD) in comparison to aspirin tolerating asthmatics.²⁵ However, the former group did not precisely match the latter in terms of age and severity of AHR.²⁵ The mechanisms regulating intravascular sphingolipid levels are tightly controlled by *de novo* synthesis and recycling processes.²¹ The processes effectively operate in both healthy subjects and HDM-APs, so that the baseline




Fig. 2. Changes in plasma S1P (A, B and C) and SFA (D, E and F) concentrations during allergen challenge in dual responders (A and D), single responders (B and E) and non-responders (C and F).

S1P, sphingosine-1-phosphate; SFO, sphingosine; SFA, sphinganine; SFA1P, sphinganine-1-phosphate.

*Significantly greater than at baseline; †Significantly less than at baseline.




Fig. 3. Correlations between changes in plasma S1P concentration during EAR and the severity of subsequent LAR. S1P, sphingosine-1-phosphate; EAR, early asthmatic response; LAR, late asthmatic response.

plasma concentration of individual sphingolipids in the studied patients remained within the broad range determined previously in a population of healthy subjects.²⁴ Yet, even in healthy subjects, sphingolipid metabolism is affected by age, sex, weight, exercise and blood morphology resulting in significant differences in plasma concentration of individual sphingolipids.²¹ In contrast to the study by Trinh *et al.*²⁵ who found a weak correlation between AHR and serum sphingomyelin concentration, we were able to demonstrate a significant association of AHR with plasma concentrations of S1P and SFA1P. Sphingomyelin

represents a reservoir of sphingolipids, but its effects on cell growth and apoptosis are mainly mediated by its metabolite ceramide.^{12,14} It is difficult to speculate if the differences are related to blood processing techniques or truly represent characteristic features of allergic *vs* aspirin intolerant asthmatic patients. In the blood, platelets constitute a very rich source of S1P which is released upon their activation.²¹ Therefore, assessment of sphingolipid concentration in plasma, not in serum, seems to better reflect their *in vivo* metabolism.²¹ Moreover, it cannot be unequivocally inferred from the study by Trinh *et al.*

²⁵ whether greater AHR seen in AERD asthmatics is independently linked to altered sphingolipid metabolism as the latter is affected by age and weight of the studied subjects. In our patients, the associations of plasma S1P and SFA1P concentration with lung function and AHR remained significant after controlling for age, sex, body mass index (BMI), TC and HDL-C concentration. Moreover, our study was performed on a homogenous population as all patients were allergic to HDMs. The association between plasma concentration of both phosphorylated sphingolipids with AHR indicates a possible role of enhanced SPHKs activity in the pathogenesis of allergic asthma. In fact, experimental studies demonstrated a crucial role of SPHKs in the development of asthma phenotype in mice.^{17,26} The study with pharmacological inhibition of SPHKs also indicated that the enzymes were crucial for developing AHR, but not airway inflammation in allergic asthma model.²⁶ This is consistent with our findings which indicate that baseline plasma concentration of phosphorylated sphingolipid derivatives, S1P and SFA1P, correlated with AHR, but not with FeNO. On the contrary, elevated plasma concentration of SFA did not correlate with lung function or AHR, but correlated with FeNO. This suggests that lung function impairment and chronic airway inflammation affect different pathways of sphingolipid metabolism. Furthermore, the correlation of plasma SFA with FeNO, but not with AHR, indicates that *de novo* sphingolipid synthesis may represent a counterregulatory mechanism which is triggered by chronic airway inflammation yet inhibits the development of asthma phenotype in HDM-APs.

Analysis of the dynamic changes in plasma sphingolipid concentration after allergen challenge reinforces the concept of altered sphingolipid metabolisms in the pathogenesis of asthma in HDM-APs. Among all sphingolipids studied, only plasma concentration of S1P increased significantly during EAR in DRs, but not in SRs or NRs. Previously, increased S1P concentrations were demonstrated in BAL fluid 24 hours after segmental allergen challenge in asthmatic, but not non-asthmatic allergic patients.¹⁹ Interestingly, the major increase in plasma S1P concentration in the current study was demonstrated much earlier than that reported in BAL fluid.¹⁹ An influx of S1P from plasma into bronchial compartment in response to allergen challenge may represent one of the mechanisms responsible for a subsequent increase in its local concentration in the airways. The reported mean concentration of S1P in BAL fluid was more than 20-fold lower than that seen in plasma in our study, which supports a possible S1P influx from plasma to the airway compartment during LAR.¹⁹ Interestingly, a systemic administration of S1P was sufficient for developing AHR and asthma phenotype in mice,¹⁵ which supports our observations that changes in intravascular sphingolipid metabolism correlated with lung function and AHR. Activated platelets and/or eosinophils, which are major determinants of plasma S1P concentration, may also participate in local increases in S1P concentration in the airways. It was shown that eosinophil number in BAL fluid correlated with S1P level 24 hours after the challenge.¹⁹ However, eosinophils were not sufficient to up-regulate S1P synthesis because persistent eosinophilia in BAL 7 days after the challenge was not associated with elevated S1P concentration.¹⁹ Platelets are a very rich source of S1P, which can be released from alpha-granules upon platelet activation.²¹ Allergen challenge leads to intravascular platelet

activation in asthmatic patients.²² Moreover, allergen challenge leads to a decrease in the number of circulating platelets and eosinophils possibly due to their sequestration into bronchial tissues.²² It seems therefore that differences in sphingolipid metabolisms between DRs and SRs at T_{EAR} may explain participation of S1P in enhanced AHR associated with the development of LAR in HDM-APs.^{5,7}

Other experimental studies provide further support for a possible role of S1P in the development of chronic airway inflammation and AHR. In an experimental allergic asthma model, application of FTY720, which down-regulates the activity of S1P receptors S1P₁, S1P₂ and S1P₅, significantly attenuated influx of Th2 cells and eosinophils into the airways of mice sensitized and challenged with ovalbumin.²⁷ In a rat model of repeated allergen exposure, application of FTY720 attenuated airway remodeling.¹⁸ S1P could also affect AHR acting directly on smooth muscle cells by inducing a broad range of genes coding for proteins involved in cell proliferation which led to tissue remodeling.^{20,28} Moreover, S1P plays an important role in inflammatory cell chemotaxis, including that of eosinophils, and interacts with its endothelial receptors promoting P-selectin-dependent role of leukocytes.^{29,30} Therefore, it is likely that an increase in plasma S1P at T_{EAR} in DRs participates in influx of inflammatory cells, which in turn is responsible for the development of LAR. This concept is supported by the correlation between changes in plasma S1P concentration at T_{EAR} and the magnitude of the subsequently developed LAR. It may also indicate stronger activation of circulating platelets and eosinophils which are major determinants of plasma S1P concentration. In fact, stronger activation of circulating platelets in response to allergen challenge in DRs in comparison to SRs has already been demonstrated.²² Finally, S1P can directly activate mast cells leading to enhanced bronchoconstriction.^{31,32} We were able to demonstrate the correlation between the severity of allergen-induced bronchoconstriction and changes in plasma S1P, supporting a potential role of S1P in amplification of EAR. Interestingly, we were not able to demonstrate a direct correlation between plasma S1P concentration and FeNO, indicating a possible indirect interaction between chronic airway inflammation and intravascular S1P metabolism. However, the plasma levels of SFA and ceramide correlated with FeNO, which indicates activation of *de novo* sphingolipid synthesis in response to chronic airway inflammation. Allergen challenge triggered *de novo* sphingolipid synthesis, but the effect was seen only in NRs at T₂₄; on the contrary, in Rs a tendency for a decrease in plasma SFA was demonstrated. This argues for a protective rather than pro-asthmatic effects of *de novo* sphingolipid synthesis. Failure to up-regulate *de novo* sphingolipid synthesis in Rs within 24 hours after the challenge may reflect impaired or delayed compensatory mechanisms in the patients. Orosomucoid like 3 (*ORMDL3*) which regulates serine palmitoyltransferase activity has recently been implicated in asthma pathogenesis.³³ The presence of functional single nucleotide mutations in the *ORMDL3* gene which increase its expression is highly correlated with the risk of childhood asthma.³³ It seems that elevated *ORMDL3* synthesis increases the risk for asthma by depressing *de novo* sphingolipid biosynthesis.³⁴ If *de novo* ceramide synthesis in response to allergen challenge were attenuated, the effect of S1P would be enhanced. This concept is supported by detection of a lower ceramide/S1P ratio in the plasma of HDM-APs with greater AHR. It is also consistent with a recently published paper which demonstrates that in mice which overexpress *ORMDL3* exogenous S1P dramatically increases AHR and that the effect is much greater than that seen in wild type mice.^{35,36} The small number of patients included in this study did not allow us to perform genetic studies regarding mutations in the *ORMDL3* gene. However, the detailed assessment of the patients included as well as multiple longitudinal sampling seems to at least partially offsets the limitations associated with the small number of patients studied.

In addition, consistent changes in plasma S1P concentration in DRs and SRs allowed for powerful demonstration of altered sphingolipid metabolism in relation to the development of prolonged airway inflammation associated with late phase asthmatic response.

Increasing body of evidence indicates that asthma is heterogenous and that at least several endotypes based on pathogenetic mechanisms may represent targets for different therapeutic options.^{37,38} Further studies are warranted to demonstrate whether altered sphingolipid metabolisms represent distinct endotypes of allergic asthma or represent characteristic features of allergic asthmatic patients.

In summary, our study provides evidence for altered intravascular sphingolipid metabolisms in HDM-allergic asthmatic patients. Both enhanced synthesis of S1P possibly derived from platelets and/or eosinophils as well as impaired *de novo* sphingolipid synthesis may participate in the development of asthma phenotype in HDM-APs. Pharmacological modulation of sphingolipid metabolisms seems to be an attractive option for the management of allergic asthma.

ACKNOWLEDGMENTS

The study was funded by a grant N/ST/ZB/17/007/1118 from the Medical University of Białystok.

REFERENCES

1. Lötvald J, Akdis CA, Bacharier LB, Bjermer L, Casale TB, Custovic A, et al. Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *J Allergy Clin Immunol* 2011;127:355-60.
[PUBMED](#) | [CROSSREF](#)
2. Nelson RP Jr, DiNicolo R, Fernández-Caldas E, Seleznick MJ, Lockey RF, Good RA. Allergen-specific IgE levels and mite allergen exposure in children with acute asthma first seen in an emergency department and in nonasthmatic control subjects. *J Allergy Clin Immunol* 1996;98:258-63.
[PUBMED](#) | [CROSSREF](#)
3. Wong GW, Li ST, Hui DS, Fok TF, Zhong NS, Chen YZ, et al. Individual allergens as risk factors for asthma and bronchial hyperresponsiveness in Chinese children. *Eur Respir J* 2002;19:288-93.
[PUBMED](#) | [CROSSREF](#)
4. Obase Y, Shimoda T, Mitsuta K, Matsuo N, Matsuse H, Kohno S. Sensitivity to the house dust mite and airway hyperresponsiveness in a young adult population. *Ann Allergy Asthma Immunol* 1999;83:305-10.
[PUBMED](#) | [CROSSREF](#)
5. Kowal K, Møller HJ, Dubuske LM, Moestrup SK, Bodzenta-Lukaszyk A. Differential expression of monocyte CD163 in single- and dual-asthmatic responders during allergen-induced bronchoconstriction. *Clin Exp Allergy* 2006;36:1584-91.
[PUBMED](#) | [CROSSREF](#)
6. Kowal K, Bodzenta-Lukaszyk A, Zukowski S. Exhaled nitric oxide in evaluation of young adults with chronic cough. *J Asthma* 2009;46:692-8.
[PUBMED](#) | [CROSSREF](#)
7. Hall S, Agrawal DK. Key mediators in the immunopathogenesis of allergic asthma. *Int Immunopharmacol* 2014;23:316-29.
[PUBMED](#) | [CROSSREF](#)
8. Chapman DG, Irvin CG. Mechanisms of airway hyper-responsiveness in asthma: the past, present and yet to come. *Clin Exp Allergy* 2015;45:706-19.
[PUBMED](#) | [CROSSREF](#)
9. Jolly PS, Rosenfeldt HM, Milstien S, Spiegel S. The roles of sphingosine-1-phosphate in asthma. *Mol Immunol* 2002;38:1239-45.
[PUBMED](#) | [CROSSREF](#)

10. Lai WQ, Wong WS, Leung BP. Sphingosine kinase and sphingosine 1-phosphate in asthma. *Biosci Rep* 2011;31:145-50.
[PUBMED](#) | [CROSSREF](#)
11. Schaubberger E, Peinhaupt M, Cazares T, Lindsley AW. Lipid mediators of allergic disease: pathways, treatments, and emerging therapeutic targets. *Curr Allergy Asthma Rep* 2016;16:48.
[PUBMED](#) | [CROSSREF](#)
12. Ono JG, Worgall TS, Worgall S. Airway reactivity and sphingolipids-implications for childhood asthma. *Mol Cell Pediatr* 2015;2:13.
[PUBMED](#) | [CROSSREF](#)
13. Yang Y, Uhlig S. The role of sphingolipids in respiratory disease. *Ther Adv Respir Dis* 2011;5:325-44.
[PUBMED](#) | [CROSSREF](#)
14. Nixon GF. Sphingolipids in inflammation: pathological implications and potential therapeutic targets. *Br J Pharmacol* 2009;158:982-93.
[PUBMED](#) | [CROSSREF](#)
15. Roviezzo F, Sorrentino R, Bertolino A, De Gruttola L, Terlizzi M, Pinto A, et al. S1P-induced airway smooth muscle hyperresponsiveness and lung inflammation *in vivo*: molecular and cellular mechanisms. *Br J Pharmacol* 2015;172:1882-93.
[PUBMED](#) | [CROSSREF](#)
16. Chiba Y, Suzuki K, Kurihara E, Uechi M, Sakai H, Misawa M. Sphingosine-1-phosphate aggravates antigen-induced airway inflammation in mice. *Open Respir Med J* 2010;4:82-5.
[PUBMED](#) | [CROSSREF](#)
17. Price MM, Oskeritzian CA, Falanga YT, Harikumar KB, Allegood JC, Alvarez SE, et al. A specific sphingosine kinase 1 inhibitor attenuates airway hyperresponsiveness and inflammation in a mast cell-dependent murine model of allergic asthma. *J Allergy Clin Immunol* 2013;131:501-511.e1.
[PUBMED](#) | [CROSSREF](#)
18. Karmouty-Quintana H, Siddiqui S, Hassan M, Tsuchiya K, Risse PA, Xicota-Vila L, et al. Treatment with a sphingosine-1-phosphate analog inhibits airway remodeling following repeated allergen exposure. *Am J Physiol Lung Cell Mol Physiol* 2012;302:L736-45.
[PUBMED](#) | [CROSSREF](#)
19. Ammit AJ, Hastie AT, Edsall LC, Hoffman RK, Amrani Y, Krymskaya VP, et al. Sphingosine 1-phosphate modulates human airway smooth muscle cell functions that promote inflammation and airway remodeling in asthma. *FASEB J* 2001;15:1212-4.
[PUBMED](#) | [CROSSREF](#)
20. Fuerst E, Foster HR, Ward JP, Corrigan CJ, Cousins DJ, Woszczek G. Sphingosine-1-phosphate induces pro-remodeling response in airway smooth muscle cells. *Allergy* 2014;69:1531-9.
[PUBMED](#) | [CROSSREF](#)
21. Książek M, Chacińska M, Chabowski A, Baranowski M. Sources, metabolism, and regulation of circulating sphingosine-1-phosphate. *J Lipid Res* 2015;56:1271-81.
[PUBMED](#) | [CROSSREF](#)
22. Kowal K, Pampuch A, Kowal-Bielecka O, DuBuske LM, Bodzenta-Lukaszyk A. Platelet activation in allergic asthma patients during allergen challenge with *Dermatophagoides pteronyssinus*. *Clin Exp Allergy* 2006;36:426-32.
[PUBMED](#) | [CROSSREF](#)
23. Baranowski M, Charnas M, Długolecka B, Górski J. Exercise increases plasma levels of sphingoid base-1 phosphates in humans. *Acta Physiol (Oxf)* 2011;203:373-80.
[PUBMED](#) | [CROSSREF](#)
24. Knapp M, Lisowska A, Knapp P, Baranowski M. Dose-dependent effect of aspirin on the level of sphingolipids in human blood. *Adv Med Sci* 2013;58:274-81.
[PUBMED](#) | [CROSSREF](#)
25. Trinh HK, Kim SC, Cho K, Kim SJ, Ban GY, Yoo HJ, et al. Exploration of the sphingolipid metabolite, sphingosine-1-phosphate and sphingosine, as novel biomarkers for aspirin-exacerbated respiratory disease. *Sci Rep* 2016;6:36599.
[PUBMED](#) | [CROSSREF](#)
26. Chiba Y, Takeuchi H, Sakai H, Misawa M. SKI-II, an inhibitor of sphingosine kinase, ameliorates antigen-induced bronchial smooth muscle hyperresponsiveness, but not airway inflammation, in mice. *J Pharmacol Sci* 2010;114:304-10.
[PUBMED](#) | [CROSSREF](#)

27. Sawicka E, Zuany-Amorim C, Manlius C, Trifileff A, Brinkmann V, Kemeny DM, et al. Inhibition of Th1- and Th2-mediated airway inflammation by the sphingosine 1-phosphate receptor agonist FTY720. *J Immunol* 2003;171:6206-14.
[PUBMED](#) | [CROSSREF](#)
28. Rumzhum NN, Rahman MM, Oliver BG, Ammit AJ. Effect of sphingosine 1-phosphate on cyclooxygenase-2 expression, prostaglandin E2 secretion, and β 2-adrenergic receptor desensitization. *Am J Respir Cell Mol Biol* 2016;54:128-35.
[PUBMED](#) | [CROSSREF](#)
29. Roviezzo F, Del Galdo F, Abbate G, Bucci M, D'Agostino B, Antunes E, et al. Human eosinophil chemotaxis and selective *in vivo* recruitment by sphingosine 1-phosphate. *Proc Natl Acad Sci U S A* 2004;101:11170-5.
[PUBMED](#) | [CROSSREF](#)
30. Nussbaum C, Bannenberg S, Keul P, Gräler MH, Gonçalves-de-Albuquerque CF, Korhonen H, et al. Sphingosine-1-phosphate receptor 3 promotes leukocyte rolling by mobilizing endothelial P-selectin. *Nat Commun* 2015;6:6416.
[PUBMED](#) | [CROSSREF](#)
31. Olivera A, Urtz N, Mizugishi K, Yamashita Y, Gilfillan AM, Furumoto Y, et al. IgE-dependent activation of sphingosine kinases 1 and 2 and secretion of sphingosine 1-phosphate requires Fyn kinase and contributes to mast cell responses. *J Biol Chem* 2006;281:2515-25.
[PUBMED](#) | [CROSSREF](#)
32. Roviezzo F, Sorrentino R, Iacono VM, Brancaleone V, Terlizzi M, Riemma MA, et al. Disodium cromoglycate inhibits asthma-like features induced by sphingosine-1-phosphate. *Pharmacol Res* 2016;113:626-35.
[PUBMED](#) | [CROSSREF](#)
33. Moffatt MF, Kabesch M, Liang L, Dixon AL, Strachan D, Heath S, et al. Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature* 2007;448:470-3.
[PUBMED](#) | [CROSSREF](#)
34. Levy BD. Sphingolipids and susceptibility to asthma. *N Engl J Med* 2013;369:976-8.
[PUBMED](#) | [CROSSREF](#)
35. Miller M, Rosenthal P, Beppu A, Gordillo R, Broide DH. Oroscomucoid like protein 3 (ORMDL3) transgenic mice have reduced levels of sphingolipids including sphingosine-1-phosphate and ceramide. *J Allergy Clin Immunol* 2017;139:1373-1376.e4.
[PUBMED](#) | [CROSSREF](#)
36. Miller M, Tam AB, Mueller JL, Rosenthal P, Beppu A, Gordillo R, et al. Cutting edge: targeting epithelial ORMDL3 increases, rather than reduces, airway responsiveness and is associated with increased sphingosine-1-phosphate. *J Immunol* 2017;198:3017-22.
[PUBMED](#) | [CROSSREF](#)
37. Ferrando M, Bagnasco D, Varricchi G, Bernardi S, Bragantini A, Passalacqua G, et al. Personalized medicine in allergy. *Allergy Asthma Immunol Res* 2017;9:15-24.
[PUBMED](#) | [CROSSREF](#)
38. Braido F, Tiotiu A, Kowal K, Mihaicuta S, Novakova P, Oguzulgen IK. Phenotypes/endotypes-driven treatment in asthma. *Curr Opin Allergy Clin Immunol* 2018;18:184-9.
[PUBMED](#) | [CROSSREF](#)