



pISSN 2586-3290 · eISSN 2586-3533  
Arch Hand Microsurg 2022;27(4):279-284  
<https://doi.org/10.12790/ahm.22.0054>

**Received:** October 6, 2022  
**Revised:** October 26, 2022  
**Accepted:** October 27, 2022

### Corresponding author:

Jong Pil Kim  
Department of Orthopaedic Surgery,  
Naeunpil Hospital, 64 Mannam-ro,  
Dongnam-gu, Cheonan 31124, Korea  
Tel: +82-41-417-0001  
Fax: +82-41-417-0001  
E-mail: scanlip@naver.com  
ORCID:  
<https://orcid.org/0000-0002-1913-9163>

# Comparison of ulnar shortening osteotomy for idiopathic ulnar impaction syndrome using conventional or ulnar osteotomy plates and with or without interfragmentary screw fixation

Kyung Wook Kim<sup>1</sup>, Ju Hyun Kim<sup>1</sup>, Jeong Sang Kim<sup>1</sup>, Bum Jin Shin<sup>2</sup>,  
Jong Pil Kim<sup>3</sup>

<sup>1</sup>Department of Orthopaedic Surgery, Dankook University College of Medicine, Cheonan, Korea

<sup>2</sup>Department of Orthopaedic Surgery, Kyungpook National University Chilgok Hospital, Daegu, Korea

<sup>3</sup>Department of Orthopaedic Surgery, Naeunpil Hospital, Cheonan, Korea

**Purpose:** This study investigated the impact of plate type on the clinical and radiological outcomes of ulnar shortening osteotomy (USO) by comparing conventional and ulnar osteotomy plates. The effect of interfragmentary screw fixation (ISF) during USO was also assessed.

**Methods:** Seventy-eight patients were divided into three groups according to the type of plate: 3.5-mm dynamic compression plate (DCP), 3.5-mm limited contact DCP, and 2.7-mm locking compression plate ulna osteotomy system (all from Depuy-Synthes). The patients were also divided into two groups according to whether ISF was performed. Clinical and radiological outcomes, including time to bone union, presence of delayed union, and refracture after hardware removal, were analyzed. Other factors that might affect bone union, such as smoking and underlying diseases, were also evaluated.

**Results:** No significant differences were found in clinical and radiological outcomes according to the type of plate. Eight of 51 patients (15.7%) in the without-ISF group showed delayed bone union. Forty-three patients in the without-ISF group underwent hardware removal, and refracture due to low-energy trauma after hardware removal was observed in five of those 43 patients (11.6%). Bone union time was significantly shorter in the with-ISF group ( $7.6 \pm 2.7$  weeks vs.  $9.8 \pm 6.6$  weeks). Diabetes mellitus and ISF were associated with the delayed bone union.

**Conclusion:** The plate type had no influence on the clinical and radiological outcomes of USO in patients with idiopathic ulnar impaction syndrome. However, ISF during USO has several advantages, such as early bony union and prevention of refracture after hardware removal.

**Keywords:** Ulna, Shortening, Osteotomy, Re-fracture, Bone union

## Introduction

Ulnar impaction syndrome (UIS) can cause medial wrist pain related to excessive weight bearing across the ulnar aspect of the wrist. It occurs when the ulnar head abuts the triangular fibrocartilage complex and carpus, resulting in ulnar-sided wrist pain [1]. Moreover, idiopathic UIS is generally seen in patients with static or dynamic positive ulnar variance with wrist pronation and forceful grip [1,2]. Ulnar shortening osteotomy (USO) was first described as an operative

intervention to decompress the ulnocarpal articulation by Milch [3] in 1941. Since then, it has been widely used in clinical practice and various modified surgical methods have been introduced. Although USO has been successfully used for the treatment of idiopathic UIS [4–6], the surgery carries the risk of hardware irritation, delayed union, and nonunion [7,8].

The purpose of this study was to compare the clinical and radiological outcomes of conventional and ulnar osteotomy plates for USO in patients with idiopathic UIS. In addition, the effect of interfragmentary screw fixation (ISF) in USO was assessed.

## Methods

**Ethics statement:** This study design was approved by the Institutional Review Board of Dankook University Hospital (No. 2022-09-013). The study was performed in accordance with the Declaration of Helsinki, and written informed consent was obtained for publication of this article and accompanying images from the patients.

From March 2008 to May 2016, 78 patients who underwent USO using plate fixation were retrospectively reviewed. Among the patients who were diagnosed with UIS and took the USO, anyone who was diagnosed with secondary UIS and had less than 1 year of the follow-up period was excluded. All patients were evaluated and treated by a single specialized hand surgeon. Three types of plates were consecutively used: 3.5-mm dynamic compression plate (DCP; Depuy-Synthes, West Chester, PA, USA), group I (n = 31); 3.5-mm limited contact DCP (LC-DCP, Depuy-Synthes), group II (n = 19); and 2.7-mm locking compression plate (LCP) ulna osteotomy system (Depuy-Synthes), group III (n = 28). And also, patients were divided into two groups based on whether they underwent ISF: with-ISF group (n = 27) and without-ISF group (n = 51).

Clinical outcomes were respectively evaluated by DASH (Disability of Arm, Shoulder and Hand) and PRWE (Patient Related Wrist Evaluation) scores preoperatively and 1 year after surgery. In addition, radiological outcomes, including time to bone union, presence of delayed union, and refracture after hardware removal were assessed. Bone union was defined radiologically when callus formation was observed on three of the four cortices on anteroposterior and lateral plain radiographs. Furthermore, the delayed union was defined when the union was not observed until 12 weeks. Other possible factors that might affect bone union, such as smoking and underlying diseases (that is, diabetes mellitus) were also assessed.

## 1. Surgical procedures

The USO was performed by a single specialized hand surgeon. Before performing USO, the author planned the necessary amount of resection to restore the ulnar variance between 0 and –1 cm. On the distal ulnar-sided forearm, 7 to 8 cm of skin incision was placed. The ulna was approached by dissecting between the flexor carpi ulnaris and extensor carpi ulnaris, while preserving the dorsal sensory branch of the ulnar nerve. After that, plate was temporarily applied using guide pins to predrill the two distal holes and one proximal hole. After removing the plate, an oblique-cut osteotomy was performed considering the edges and rotation without any cutting guide or additional apparatus. The plate was placed using a reduction clamp to achieve anatomical reduction, and then screws were firmly secured. All patients who had undergone USO were applied protective long arm splint for 2 weeks, short arm cast for 4 weeks, and wrist brace for 6 weeks, postoperatively.

## 2. Statistical analysis

Categorical variables were compared using the chi-square test and continuous variables were compared using the independent t-test and one-way ANOVA. Factors potentially associated with bone union time were analyzed using a logistic regression analysis. Statistical analyses were performed using IBM SPSS Statistics (IBM Corp., Armonk, NY, USA). Results with p-values that were less than 0.05 were considered for statistical significance.

## Results

The average follow-up time was 15.8 months (range, 12–27 months). There were 43 male and 35 female patients with a mean age of 42 years (range, 22–69 years). Out of 78 patients, there were no significant differences in clinical and radiological outcomes according to the types of plates (Table 1). When comparing groups according to whether ISF was performed, there was no significant difference in demographic factors and clinical outcomes between the two groups. However, bone union time was significantly shortened from  $9.8 \pm 6.6$  to  $7.6 \pm 2.7$  weeks in the with-ISF group. Delayed bone union was confirmed only in the without-ISF group. Eight of the 51 patients (15.7%) who were treated without interfragmentary fixation showed delayed bone union. Unlike the with-ISF group, refracture by low-energy trauma after hardware removal was observed in five of 43 patients (11.6%) in the without-ISF group. However, this difference was not significant (Table 2). Moreover, to identify the risk factor related to delayed bone union, factors potentially associ-

**Table 1.** Clinical and radiological outcomes according to the type of plate

Variable	Group I	Group II	Group III	p-value
No of patients	31	19	28	
Age (yr)	38.8 ± 11.9	40.0 ± 13.5	45.9 ± 11.8	0.109
Smoking	8 (25.8)	9 (47.4)	4 (14.3)	0.192
Diabetes mellitus	1 (3.2)	1 (5.3)	3 (10.7)	0.537
DASH score	52.1 ± 20.1	52.6 ± 23.4	56.1 ± 23.3	0.799
PRWE score	40.4 ± 31.7	38.6 ± 41.2	36.9 ± 32.4	0.929
Interfragmentary screw fixation	11 (35.5)	1 (5.3)	15 (53.6)	0.333
Time to bone union (wk)	7.4 ± 4.1	10.1 ± 7.3	10.1 ± 5.5	0.192
Refracture after metal removal	1 (3.2)	2 (10.5)	22 (78.6)	0.555
Delayed bone union	2 (6.5)	3 (15.8)	3 (10.7)	0.580

Values are presented as number only, mean ± standard deviation, or number (%).

Group I, using 3.5-mm dynamic compression plate (DCP; Depuy-Synthes, West Chester, PA, USA); group II, using 3.5-mm limited contact DCP (Depuy-Synthes); group III, using 2.7-mm locking compression plate ulna osteotomy system (Depuy-Synthes).

DASH, Disability of Arm, Shoulder and Hand; PRWE, Patient Related Wrist Evaluation.

**Table 2.** Clinical and radiological outcomes according to interfragmentary screw fixation

Variable	With-ISF group	Without-ISF group	p-value
No. of patients	27	51	
Age (yr)	25.3 ± 12.3	39.7 ± 12.4	0.065
DASH score	42.9 ± 19.3	56.7 ± 22.8	0.079
PRWE score	28.9 ± 24.8	43.9 ± 37.2	0.064
Time to bone union (wk)	7.6 ± 2.6	9.8 ± 6.6	0.038
Refracture after metal removal	0/18 (0)	5/43 (11.6)	0.157
Delayed bone union	0 (0)	8 (15.7)	0.045

Values are presented as number only, mean ± standard deviation, or number (%).

ISF, interfragmentary screw fixation; DASH, Disability of Arm, Shoulder and Hand; PRWE, Patient Related Wrist Evaluation.

ated with bone union time, such as sex, smoking, diabetes mellitus, ISF, and the type of plates were analyzed using logistic regression analyses. Among the risk factors which are related to bone union, diabetes mellitus (odds ratio [OR], 4.86;  $p=0.046$ ) and ISF (OR, 2.44;  $p=0.042$ ) were associated with the delayed bone union.

All patients showed bone union at the final follow-up. Eight patients who showed delayed bone union were treated conservatively; thereafter, they showed radiological union at a mean period of 22.2 weeks (Fig. 1). Five patients who showed refracture after hardware removal underwent revision open reduction and internal fixation ( $n=4$ ) or closed reduction and intramedullary nail fixation ( $n=1$ ), and bone union was achieved at the final follow-up (Fig. 2).

## Discussion

As a result of positive ulnar variance, overloading between the ulnar carpus and the distal ulna occurs. Hence, patients

usually report ulnar-side wrist pain, diminished grip strength, swelling, and limitation of wrist motion. USO is the preferred treatment option for correcting positive ulnar variance, which relieves pain by restoring neutral or slightly negative ulnar variance. Jungwirth-Weinberger et al. [9] reported that using 2.7-mm LCP ulnar osteotomy system plate for USO in patients with UIS displayed shorter bone healing time than that resulting from the use of 3.5-mm LC-DCP. However, in our study, there were no significant differences in either clinical or radiological outcomes according to the types of plate.

Hulsizer et al. [10] reported that 12 out of 13 patients achieved complete pain relief at a mean of 2.3 years with freehand transverse osteotomies and four out of 13 had postoperative complications. On the other hand, Chun and Palmer described a series of 30 cases of an oblique freehand osteotomy technique with good results, although there were few complications, including no nonunion [5]. Theoretically, when the interfragmentary screw crosses the fracture line, the threads engage the fragment and the lag effect of the screw reduces the fracture gap. Further



**Fig. 1.** (A) A 62-year-old man was treated with a 2.7-mm locking compression plate (LCP) ulna osteotomy system (Depuy-Synthes, West Chester, PA, USA), and interfragmentary screw fixation was not performed. Immediately postoperatively, a neutral ulnar variance was achieved and the ulnar osteotomy plate settled well. (B) Twelve weeks later, the gap at the osteotomy site had increased, and delayed bone union occurred. (C) Twenty-four weeks later, radiological bone union was spontaneously achieved. (D) A 59-year-old man was treated with a 2.7-mm LCP ulna osteotomy system (Depuy-Synthes), and interfragmentary screw fixation was not performed. (E) Thirteen weeks later, callus formation was seen around the osteotomy site. (F) Twenty-six weeks later, radiological bone union was achieved with excessive callus formation.

tightening of the lag screw pulls the distal fragment in a proximal direction, compressing the fracture site and enhancing fracture stability, which results in improved fracture healing [11]. Hence, in this study, USO was performed using the oblique osteotomy technique in all patients. Previous studies reported 0% to 18% of nonunion [12-16]. However, the incidence of the nonunion and delayed union in this present study was 0% and 10.2%, respectively, which are lower rates than most reported in the literature. A study by Chen et al. [17] demonstrated that 30% of smokers compared to 0% of non-smokers developed nonunions. Moreover, Gaspar et al. [18] revealed that smoking and diabetes mellitus were significantly elevated risks for both delayed union and nonunion in USO. However, in this present study, only diabetes mellitus and ISF

were significantly associated with the delayed bone union.

Pomerance [19] insisted that local pain or irritation was the main cause of plate removal, and demonstrated a 2.5% incidence of refracture after hardware removal (mean time to hardware removal, 6.6 months) after USO. Furthermore, according to Mih et al. [20], 11.3% of patients experienced refracture after hardware removal (mean time to hardware removal, 19 months) from various forearm pathologies. We demonstrated a 78.2% (61 of 78) incidence of hardware removal in this study, which is comparable to the results of other studies. Among them, only five patients (8.2%) experienced a refracture after hardware removal. All patients underwent revision surgery, and the bone union was achieved at the final follow-up.

This study had several limitations. First, the retrospective na-





**Fig. 2.** (A) A 49-year-old woman underwent ulnar shortening osteotomy without interfragmentary screw fixation. Eleven months later, solid bone union was achieved and hardware removal was done. (B) Refracture occurred by low-energy trauma after hardware removal. (C) A revisional operation was done with ulnar osteotomy plate and interfragmentary screw fixation was performed. Seven weeks later, solid bone union was achieved. (D) A 29-year-old man underwent ulnar shortening osteotomy without interfragmentary screw fixation. Twelve months later, solid bone union was achieved and hardware removal was performed. (E) After hardware removal, refracture by low-energy trauma occurred. (F) Revisional closed reduction and intramedullary nail fixation were performed. Nine weeks later, solid bone union was achieved.

ture of this study could have introduced selection bias. The author suggests that a randomized comparative study would help elucidate the most important factors. Second, the degree of diabetes mellitus control has not been investigated in detail. These

details may have been partly responsible for the differences in the results. Third, in this study, not like the category of delayed bone union, refracture after metal removal showed no significant difference between the with-ISF group and the with-

out-ISF group. As refracture after metal removal could be considered a serious complication, this could undermine claims about the effectiveness of ISF. However, unlike the without-ISF group, which showed five cases of refracture (11.6%) after metal removal, none (0%) occurred in the with-ISF group. Hence, although a significant difference was not achieved, it was assumed that it could sufficiently support the effectiveness of ISF.

## Conclusion

Plate types did not affect the clinical and radiological outcomes of USO in patients with idiopathic UIS. However, ISF during USO has several advantages, such as early bony union and prevention of refracture after hardware removal. In conclusion, ISF during USO can be a good surgical option for patients with UIS.

## ORCID

Kyung Wook Kim, <https://orcid.org/0000-0003-3879-0768>

Ju Hyun Kim, <https://orcid.org/0000-0001-6454-3563>

Jeong Sang Kim, <https://orcid.org/0000-0003-3617-833X>

Bum Jin Shin, <https://orcid.org/0000-0002-3751-2304>

Jong Pil Kim, <https://orcid.org/0000-0002-1913-9163>

## Conflicts of interest

The authors have nothing to disclose.

## Funding

None.

## References

1. Friedman SL, Palmer AK, Short WH, Levinsohn EM, Halperin LS. The change in ulnar variance with grip. *J Hand Surg Am.* 1993;18:713-6.
2. Palmer AK, Glisson RR, Werner FW. Ulnar variance determination. *J Hand Surg Am.* 1982;7:376-9.
3. Milch H. Cuff resection of the ulna for malunited Colles' fracture. *J Bone Joint Surg.* 1941;23:311-3.
4. Baek GH, Chung MS, Lee YH, Gong HS, Lee S, Kim HH. Ulnar shortening osteotomy in idiopathic ulnar impaction syndrome. *J Bone Joint Surg Am.* 2005;87:2649-54.
5. Chun S, Palmer AK. The ulnar impaction syndrome: follow-up of ulnar shortening osteotomy. *J Hand Surg Am.* 1993;18:46-53.
6. Minami A, Kato H. Ulnar shortening for triangular fibrocartilage complex tears associated with ulnar positive variance. *J Hand Surg Am.* 1998;23:904-8.
7. Constantine KJ, Tomaino MM, Herndon JH, Sotereanos DG. Comparison of ulnar shortening osteotomy and the wafer resection procedure as treatment for ulnar impaction syndrome. *J Hand Surg Am.* 2000;25:55-60.
8. Vandenberghe L, Degreef I, Didden K, Moermans A, Kooreneef P, De Smet L. Ulnar shortening or arthroscopic wafer resection for ulnar impaction syndrome. *Acta Orthop Belg.* 2012;78:323-6.
9. Jungwirth-Weinberger A, Borbas P, Schweizer A, Nagy L. Influence of plate size and design upon healing of ulna-shortening osteotomies. *J Wrist Surg.* 2016;5:284-9.
10. Hulsizer D, Weiss AP, Akelman E. Ulna-shortening osteotomy after failed arthroscopic debridement of the triangular fibrocartilage complex. *J Hand Surg Am.* 1997;22:694-8.
11. Magee W, Hettwer W, Badra M, Bay B, Hart R. Biomechanical comparison of a fully threaded, variable pitch screw and a partially threaded lag screw for internal fixation of type II dens fractures. *Spine (Phila Pa 1976).* 2007;32:E475-9.
12. Chen NC, Wolfe SW. Ulna shortening osteotomy using a compression device. *J Hand Surg Am.* 2003;28:88-93.
13. Darlis NA, Ferraz IC, Kaufmann RW, Sotereanos DG. Step-cut distal ulnar-shortening osteotomy. *J Hand Surg Am.* 2005;30:943-8.
14. Mizuseki T, Tsuge K, Ikuta Y. Precise ulna-shortening osteotomy with a new device. *J Hand Surg Am.* 2001;26:931-9.
15. Van Sanden S, De Smet L. Ulnar shortening after failed arthroscopic treatment of triangular fibrocartilage complex tears. *Chir Main.* 2001;20:332-6.
16. Wehbe MA, Cautilli DA. Ulnar shortening using the AO small distractor. *J Hand Surg Am.* 1995;20:959-64.
17. Chen F, Osterman AL, Mahony K. Smoking and bony union after ulna-shortening osteotomy. *Am J Orthop (Belle Mead NJ).* 2001;30:486-9.
18. Gaspar MP, Kane PM, Zohn RC, Buckley T, Jacoby SM, Shin EK. Variables prognostic for delayed union and nonunion following ulnar shortening fixed with a dedicated osteotomy plate. *J Hand Surg Am.* 2016;41:237-43.
19. Pomerance J. Plate removal after ulnar-shortening osteotomy. *J Hand Surg Am.* 2005;30:949-53.
20. Mih AD, Cooney WP, Idler RS, Lewallen DG. Long-term follow-up of forearm bone diaphyseal plating. *Clin Orthop Relat Res.* 1994;(299):256-8.