

Corrigenda: Table Correction

Nonalcoholic Fatty Liver Disease in Diabetes. Part I: Epidemiology and Diagnosis

Yong-ho Lee^{1,2}, Yongin Cho^{1,2}, Byung-Wan Lee^{1,2}, Cheol-Young Park³, Dae Ho Lee⁴, Bong-Soo Cha^{1,2},
Eun-Jung Rhee³

¹Department of Internal Medicine, ²Institute of Endocrine Research, Yonsei University College of Medicine, Seoul,

³Division of Endocrinology and Metabolism, Department of Internal Medicine, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul,

⁴Department of Internal Medicine, Wonkwang University Hospital, Wonkwang University School of Medicine, Iksan, Korea

Diabetes Metab J 2019;43:31-45. <https://doi.org/10.4093/dmj.2019.0011>

In this article, the description of one equation named “NAFLD liver fat score” in the Table 3 was miswritten.

NAFLD liver fat score = $-2.89 + 1.18 \times \text{MetS} + 0.45 \times \text{diabetes (yes=2/no=0)} + 0.15 \times (\text{fSinsulin, } \mu\text{U/L}) + 0.04 \times \text{AST} - \underline{\mathbf{0.94}} \times \text{AST/ALT}$

The corrected part is written in bold characters with underlines. This Table has been corrected in the online versions.

We apologize for any inconvenience that this may have caused.