

Noncanonical Wnt Signaling in Bone

Fanxin Long

Department of Medicine and Department of Developmental Biology, Washington University School of Medicine, St. Louis, MO, USA

INTRODUCTION

The Wnt family of proteins plays essential roles conserved from flies to human in many developmental processes including cell fate specification, polarity, migration and proliferation. It has becoming increasingly clear that Wnt proteins activate multiple intracellular signaling cascades to achieve the diverse biological functions. One such cascade that has been most intensely studied to date involves the multi-functional protein β -catenin; this cascade is commonly known as the canonical Wnt pathway. In this pathway, extracellular Wnt proteins (19 members in mammals) engage both the low-density lipoprotein receptor-related proteins (LRP-5 and LRP-6) [18, 22,27,31] and the Frizzled family of membrane receptors (11 members in mammals), resulting in stabilization of the cytosolic β -catenin through inhibition of GSK3 β [13,32]. The stabilized β -catenin accumulates and transports into the nucleus where it interacts with transcription factors including lymphoid enhancer-binding factor-1 (Lef-1) and T cell factors (Tcf1-4), leading to transcriptional activation of downstream target genes [7].

For nearly a decade, bone biologists have studied Wnt signaling intensely. This strong interest was kindled by the discovery that loss-of-function mutations in the Wnt co-receptor LRP-5 causes the hereditary human disease osteoporosis-pseudoglioma syndrome [8]. Further supporting the role of LRP5 in bone mass regulation, multiple gain-of-function mutations in LRP5 were found to cause a high-bone-mass syndrome in humans [3,16]. Moreover, loss-of-function mutations in a secreted Wnt antagonist sclerostin (SOST) were discovered to cause rare high-bone-mass syndromes known as sclerosteosis and the van Buchem disease [1,2,4,25]. The role of Wnt signaling in bone formation, as revealed by the human genetic studies, was further demonstrated in mouse genetic models. Spe-

cifically, mice deficient in *Lrp-5* (*Lrp-5*^{-/-}) exhibit low bone mass postnatally due to reduced osteoblast proliferation and function [14]. Moreover, mouse embryos lacking β -catenin in the skeletal progenitors failed to form osteoblasts altogether [6,11,12,23]. Overall, genetic evidence from both human and mouse strongly support the notion that Wnt signaling positively regulates bone formation.

NONCANONICAL WNT SIGNALING AND OSTEOBLAST DIFFERENTIATION

In addition to the β -catenin-mediated pathway, Wnt signaling through β -catenin-independent mechanisms, collectively termed noncanonical pathways, also plays fundamental roles in the vertebrate embryo [29]. Most notably, a β -catenin-independent pathway that involves the Rac1 and Cdc42 small GTPases, as well as the downstream kinase JNK, has been implicated in the directed migration and intercalation of cells towards the midline during gastrulation, a process known as the convergent extension (CE) movement [5,9,10,20]. This pathway is often considered to be similar to the planar cell polarity (PCP) pathway extensively studied in *Drosophila*. In addition, studies in *Xenopus* embryos have shown that Wnt proteins can activate heterotrimeric G-proteins and lead to calcium mobilization within the cell [15]. This process, often known as the “Ca⁺⁺ pathway”, is sensitive to pertussis toxin and therefore most likely mediated through the Gi subfamily of α subunits. Although the vast majority of studies on non-canonical Wnt signaling have been on its role on cell polarity and directed movement, recent evidence has revealed that β -catenin-independent mechanisms also play significant roles in osteoblast differentiation.

In an effort to uncover downstream effectors through which Wnt3a induces osteoblast differentiation, we serendipitously discovered a PKC δ -dependent pathway that contributes to Wnt-induced bone formation [28] (Fig. 1). By comparing the protein profiles of ST2 cells treated with Wnt3a versus untreated cells by proteomics analyses, we found that the protein MARCKS (meris-

Corresponding author: Fanxin Long
Department of Medicine and Department of Developmental Biology, Washington University School of Medicine, 660 S. Euclid Avenue, St. Louis, MO 63110, USA
Tel: +1.314-454-8795, Email: flong@wustl.edu

toylated alanine-rich C kinase substrate), a prototypic PKC substrate, was robustly phosphorylated by Wnt3a signaling. Subsequent studies identified PKC δ as the critical PKC isoform that mediates the Wnt-induced phosphorylation. More importantly, specific knockdown of PKC δ inhibited Wnt-induced osteoblast differentiation, and PKC δ -null mouse embryos exhibited a notable delay in bone collar formation. We further investigated the upstream mechanism through which Wnt3a activates PKC δ and found that heterotrimeric G-proteins composed of the Gq subfamily of α subunits are important. Because this pathway is insensitive to pertussis toxin (which inhibits Gi but not Gq subfamily), it therefore represents a separate G-protein-mediated pathway distinct from the aforementioned Ca⁺⁺ pathway. Adding to the physiological significance of this mechanism, we found that Wnt7b, a ligand selectively expressed in osteogenic cells in vivo, indeed activates the PKC δ pathway but not the β -catenin pathway, and nonetheless induces osteoblast differentiation in cell cultures. Thus, this study has provided evidence that a PKC δ -dependent mechanism, contributes to the osteogenic function of Wnt proteins independent of β -catenin. The critical target for PKC δ however is currently unknown.

More recently, Takada et al. [26] discovered that non-canonical

signaling by Wnt5a promotes osteoblast differentiation while inhibiting adipogenesis from ST2 cells (Fig. 1). In particular, the authors showed that Wnt5a activates a kinase cascade CaMKII-TAK1-NLK which leads to phosphorylation of a histone methyltransferase that suppress PPAR γ -target gene activation through inhibitory histone methylation of the promoters. Due to the suppression of the adipocyte fate, the authors argue that osteoblast differentiation is enhanced from the bi-potential mesenchymal progenitor. Because CaMKII (calcium/calmodulin-dependent protein kinase II) is activated by Ca⁺⁺, the kinase cascade uncovered here could be a part of the Ca⁺⁺ pathway as discussed above. This connection however was not directly demonstrated in this study. Overall, this study has revealed a mechanism through which Wnt proteins can alter gene transcription and hence cell fate decisions through a β -catenin-independent mechanism.

RAC1-JNK2 SIGNALING AS A CRITICAL COMPONENT OF CANONICAL WNT SIGNALING

Although nuclear localization of β -catenin is a prerequisite for canonical Wnt signaling, the underlying mechanism is poorly un-




Fig. 1. A summary of two recent studies about non-canonical Wnt signaling in osteoblast differentiation. Tu et al. [28] demonstrated a PKC δ -mediated pathway that promotes the Runx2-Osterix (Osx) transition through a yet unknown mechanism. Takada et al. [26] showed that CaMKII-mediated activation of a histone methyltransferase (SETDB1-SET domain bifurcated 1) represses PPAR γ transactivation of adipogenic target genes, thereby secondarily enhancing osteoblast differentiation. In both cases, the involvement of Frizzled family of receptors (Fz) has not been demonstrated (hence the question mark). In addition, Takada et al. did not provide evidence for the mechanism upstream of CaMKII. G: heterotrimeric G-proteins composed of α , β and γ subunits. P. T.: pertussis toxin. DAG: diacylglycerol.

derstood. In a prevailing conventional view, β -catenin is stabilized in the cytoplasm to a threshold level that allows passive entry into the nucleus. A recent study from our group however challenges this view, and provides evidence that nuclear localization is a necessary regulatory step. In particular, Rac1 and JNK2 activation appears to be required for β -catenin nuclear localization and signaling [33].

Perhaps the most surprising aspect of this finding is that the Rac1-JNK signaling module, conventionally believed to function exclusively in a non-canonical Wnt pathway, performs a critical function in β -catenin signaling. While the role of Rac1-JNK in PCP and CE is well established, the new finding highlights the realization that the same signaling molecules may perform different roles in the cell, depending on their specific interaction partners and whether they are activated in a particular protein complex. Indeed, we found that β -catenin is present in a protein complex with Rac1 and JNK1/2 in the cytoplasm of ST2 cells, and Wnt3a selectively activates JNK2 in the complex. More importantly, Wnt7b, which does not activate β -catenin signaling even though it stabilizes the protein, does not activate JNK2 in the Rac1-JNK1/2- β -catenin protein complex, despite the fact that JNK2 activation can be detected in the whole cell lysate.

The role of Rac1 and JNK2 in β -catenin signaling has been confirmed by others in a colon adenoma model [21]. There, the authors found that loss of the tumor suppressor APC (adenomatous polyposis coli), which robustly stabilizes β -catenin, is insufficient for causing β -catenin nuclear localization, whereas addition activation of KRAS is effective. The authors further showed that KRAS activated Rac1 and JNK2, and inhibition of the latter two molecules abolished β -catenin nuclear localization and signaling in these cancer cells. Thus, the colon adenoma cancer cells appear to have hijacked the same mechanisms used by normal Wnt signaling.

RETHINKING CANONICAL VERSUS NONCANONICAL WNT PATHWAYS

Although “canonical” versus “noncanonical” Wnt pathways have been customarily used in the Wnt field, the new evidence calls for an understanding of these terms. First, it has become clear that the same Wnt ligand can activate multiple intracellular pathways in the same cell. This was evident in our studies where Wnt3a activates both the β -catenin and the PKC δ pathway [28]. The divergence in intracellular signaling pathways likely occurs at the

level of cell-surface receptors, including both the Frizzled family members and the atypical receptor kinase Ryk [17,24], as well as the orphan receptor tyrosine kinase Ror2 [19]. Second, the same Wnt ligand can activate different pathways in different cells. For instance, we’ve shown that Wnt7b fails to activate β -catenin signaling in ST2 cells [28,33], but others have shown that it can activate the pathway in epithelial and vascular smooth muscle cells [30]. The difference could be caused by the presence of different receptors in different cells. Third, whereas β -catenin stabilization is clearly important, it is not synonymous with β -catenin signaling in the nucleus. Indeed, the additional regulatory steps for β -catenin signaling may well be “non-canonical”, as demonstrated by the involvement of Rac1 and JNK2 [33]. Finally, whereas “canonical” is defined as β -catenin-dependent, the broad category of “non-canonical pathways” may include an unknown number of cascades that remain to be discovered. Thus, when it comes to Wnt signaling, one would be wise to keep a very open mind for the years to come.

REFERENCES

1. Balemans W, Ebeling M, Patel N, Van Hul E, Olson P, Dioszegi M, Lacza C, Wuyts W, Van Den Ende J, Willems P, Paes-Alves AF, Hill S, Bueno M, Ramos FJ, Tacconi P, Dikkers FG, Stratakis C, Lindpaintner K, Vickery B, Foerzler D, Van Hul W: Increased bone density in sclerosteosis is due to the deficiency of a novel secreted protein (SOST). *Hum Mol Genet* 10:537-543, 2001
2. Balemans W, Patel N, Ebeling M, Van Hul E, Wuyts W, Lacza C, Dioszegi M, Dikkers FG, Hilderling P, Willems PJ, Verheij JB, Lindpaintner K, Vickery B, Foerzler D, Van Hul W: Identification of a 52 kb deletion downstream of the SOST gene in patients with van Buchem disease. *J Med Genet* 39:91-97, 2002
3. Boyden LM, Mao J, Belsky J, Mitzner L, Farhi A, Mitnick MA, Wu D, Insogna K, Lifton RP: High bone density due to a mutation in LDL-receptor-related protein 5. *N Engl J Med* 346:1513-1521, 2002
4. Brunkow ME, Gardner JC, Van Ness J, Paepers BW, Kovacevich BR, Proll S, Skonier JE, Zhao L, Sabo PJ, Fu Y, Alisch RS, Gillett L, Colbert T, Tacconi P, Galas D, Hamersma H, Beighton P, Mulligan J: Bone dysplasia sclerosteosis results from loss of the SOST gene product, a novel cystine knot-containing protein. *Am J Hum Genet* 68:577-589, 2001
5. Choi SC, Han JK: *Xenopus* Cdc42 regulates convergent extension movements during gastrulation through Wnt/Ca²⁺ signaling pathway. *Dev Biol* 244:342-357, 2002
6. Day TE, Guo X, Garrett-Beal L, Yang Y: Wnt/beta-catenin signaling in mesenchymal progenitors controls osteoblast and chondrocyte differentiation during vertebrate skeletogenesis. *Dev Cell* 8:739-750, 2005
7. Eastman Q, Grosschedl R: Regulation of LEF-1/TCF transcription factors by Wnt and other signals. *Curr Opin Cell Biol* 11:233-240, 1999
8. Gong Y, Slee RB, Fukui N, Rawadi G, Roman-Roman S, Reginato AM,

- Wang H, Cundy T, Glorieux FH, Lev D, Zacharin M, Oexle K, Marcelino J, Suwairi W, Heeger S, Sabatakos G, Apte S, Adkins WN, Allgrove J, Arslan-Kirchner M, Batch JA, Beighton P, Black GC, Boles RG, Boon LM, Borrone C, Brunner HG, Carle GF, Dallapiccola B, De Paepe A, Floege B, Halfhide ML, Hall B, Hennekam RC, Hirose T, Jans A, Jüppner H, Kim CA, Keppler-Noreuil K, Kohlschütter A, LaCombe D, Lambert M, Lemyre E, Letteboer T, Peltonen L, Ramesar RS, Romanengo M, Somer H, Steichen-Gersdorf E, Steinmann B, Sullivan B, Superti-Furga A, Swoboda W, van den Boogaard MJ, Van Hul W, Vikkula M, Votruba M, Zabel B, Garcia T, Baron R, Olsen BR, Warman ML; Osteoporosis-Pseudoglioma Syndrome Collaborative Group: LDL receptor-related protein 5 (LRP5) affects bone accrual and eye development. *Cell* 107:513-523, 2001
9. Habas R, Dawid IB, He X: Coactivation of Rac and Rho by Wnt/Frizzled signaling is required for vertebrate gastrulation. *Genes Dev* 17:295-309, 2003
10. Habas R, Kato Y, He X: Wnt/Frizzled activation of Rho regulates vertebrate gastrulation and requires a novel Formin homology protein Daam1. *Cell* 107:843-854, 2001
11. Hill TP, Später D, Taketo MM, Birchmeier W, Hartmann C: Canonical Wnt/beta-catenin signaling prevents osteoblasts from differentiating into chondrocytes. *Dev Cell* 8:727-738, 2005
12. Hu H, Hilton MJ, Tu X, Yu K, Ornitz DM, Long F: Sequential roles of Hedgehog and Wnt signaling in osteoblast development. *Development* 132:49-60, 2005
13. Huelsen J, Birchmeier W: New aspects of Wnt signaling pathways in higher vertebrates. *Curr Opin Genet Dev* 11:547-553, 2001
14. Kato M, Patel MS, Levasseur R, Lobov I, Chang BH, Glass DA 2nd, Hartmann C, Li L, Hwang TH, Brayton CF, Lang RA, Karsenty G, Chan L: Cbfa1-independent decrease in osteoblast proliferation, osteopenia, and persistent embryonic eye vascularization in mice deficient in Lrp5, a Wnt coreceptor. *J Cell Biol* 157:303-314, 2002
15. Kühl M, Sheldahl LC, Park M, Miller JR, Moon RT: The Wnt/Ca2+ pathway: a new vertebrate Wnt signaling pathway takes shape. *Trends Genet* 16:279-283, 2000
16. Little RD, Carulli JP, Del Mastro RG, Dupuis J, Osborne M, Folz C, Manning SP, Swain PM, Zhao SC, Eustace B, Lappe MM, Spitzer L, Zweier S, Braunschweiger K, Benchekroun Y, Hu X, Adair R, Chee L, FitzGerald MG, Tulig C, Caruso A, Tzellas N, Bawa A, Franklin B, McGuire S, Nogue X, Gong G, Allen KM, Anisowicz A, Morales AJ, Lomedico PT, Recker SM, Van Eerdewegh P, Recker RR, Johnson ML: A mutation in the LDL receptor-related protein 5 gene results in the autosomal dominant high-bone-mass trait. *Am J Hum Genet* 70:11-19, 2002
17. Lu W, Yamamoto V, Ortega B, Baltimore D: Mammalian Ryk is a Wnt coreceptor required for stimulation of neurite outgrowth. *Cell* 119:97-108, 2004
18. Mao J, Wang J, Liu B, Pan W, Farr GH 3rd, Flynn C, Yuan H, Takada S, Kimelman D, Li L, Wu D: Low-density lipoprotein receptor-related protein-5 binds to Axin and regulates the canonical Wnt signaling pathway. *Mol Cell* 7:801-809, 2001
19. Mikels AJ, Nusse R: Purified Wnt5a protein activates or inhibits beta-catenin-TCF signaling depending on receptor context. *PLoS Biol* 4:e115, 2006
20. Penzo-Mendéz A, Umbhauer M, Djiane A, Boucaut JC, Riou JF: Activation of Gbetagamma signaling downstream of Wnt-11/Xfz7 regulates Cdc42 activity during *Xenopus* gastrulation. *Dev Biol* 257:302-314, 2003
21. Phelps RA, Chidester S, Dehghanizadeh S, Phelps J, Sandoval IT, Rai K, Broadbent T, Sarkar S, Burt RW, Jones DA: A two-step model for colon adenoma initiation and progression caused by APC loss. *Cell* 137:623-634, 2009
22. Pinson KI, Brennan J, Monkley S, Avery BJ, Skarnes WC: An LDL-receptor-related protein mediates Wnt signalling in mice. *Nature* 407:535-538, 2000
23. Rodda SJ, McMahon AP: Distinct roles for Hedgehog and canonical Wnt signaling in specification, differentiation and maintenance of osteoblast progenitors. *Development* 133:3231-3244, 2006
24. Schmitt AM, Shi J, Wolf AM, Lu CC, King LA, Zou Y: Wnt-Ryk signaling mediates medial-lateral retinotectal topographic mapping. *Nature* 439:31-37, 2006
25. Staehling-Hampton K, Proll S, Paepcr BW, Zhao L, Charnley P, Brown A, Gardner JC, Galas D, Schatzman RC, Beighton P, Papapoulos S, Hamersma H, Brunkow ME: A 52-kb deletion in the SOST-MEOX1 intergenic region on 17q12-q21 is associated with van Buchem disease in the Dutch population. *Am J Med Genet* 110:144-152, 2002
26. Takada I, Mihara M, Suzawa M, Ohtake F, Kobayashi S, Igarashi M, Youn MY, Takeyama K, Nakamura T, Mezaki Y, Takezawa S, Yogiashi Y, Kitagawa H, Yamada G, Takada S, Minami Y, Shibuya H, Matsumoto K, Kato S: A histone lysine methyltransferase activated by non-canonical Wnt signalling suppresses PPAR-gamma transactivation. *Nat Cell Biol* 9:1273-1285, 2007
27. Tamai K, Semenov M, Kato Y, Spokony R, Liu C, Katsuyama Y, Hess F, Saint-Jeannet JP, He X: LDL-receptor-related proteins in Wnt signal transduction. *Nature* 407:530-535, 2000
28. Tu X, Joeng KS, Nakayama KI, Nakayama K, Rajagopal J, Carroll TJ, McMahon AP, Long F: Noncanonical Wnt signaling through G protein-linked PKCdelta activation promotes bone formation. *Dev Cell* 12:113-127, 2007
29. Veeman MT, Axelrod JD, Moon RT: A second canon. Functions and mechanisms of beta-catenin-independent Wnt signaling. *Dev Cell* 5:367-377, 2003
30. Wang Z, Shu W, Lu MM, Morrissey EE: Wnt7b activates canonical signaling in epithelial and vascular smooth muscle cells through interactions with Fzd1, Fzd10, and LRP5. *Mol Cell Biol* 25:5022-5030, 2005
31. Wehrli M, Dougan ST, Caldwell K, O'Keefe L, Schwartz S, Vaizel-Ohayon D, Schejter E, Tomlinson A, DiNardo S: Arrow encodes an LDL-receptor-related protein essential for Wingless signalling. *Nature* 407:527-530, 2000
32. Wodarz A, Nusse R: Mechanisms of Wnt signaling in development. *Annu Rev Cell Dev Biol* 14:59-88, 1998
33. Wu X, Tu X, Joeng KS, Hilton MJ, Williams DA, Long F: Rac1 activation controls nuclear localization of beta-catenin during canonical Wnt signaling. *Cell* 133:340-353, 2008