

## Overview of the Renin-Angiotensin System

Wang Seong Ryu, MD, Sang Wook Kim, MD and Chee Jeong Kim, MD

Department of Internal Medicine, College of Medicine, Chung-Ang University, Seoul, Korea

### ABSTRACT

Angiotensin II is an octapeptide hormone of the renin-angiotensin system (RAS), and it regulates a wide variety of physiological responses including salt and water balance, the blood pressure and the vascular tone. Clinical trials with angiotensin-converting enzyme (ACE) inhibitors have demonstrated survival benefits in subjects with congestive heart failure and myocardial infarction, and this support the importance of angiotensin II in the pathogenesis of cardiovascular diseases. Through activation of small G proteins such as Ras, Rho, and Rac, angiotensin II induces remodeling of vascular smooth muscle cells (VSMC), including proliferation, migration, hypertrophy and inflammation. Angiotensin (1-7) appears to be the main effector peptide of ACE2 with vasodilatory, natriuretic and antiinflammatory properties. The cross-talk between the angiotensin II receptors may play an important role in maintaining cardiovascular homeostasis. (*Korean Circulation J* 2007;37:91–96)

**KEY WORDS :** Renin ; Angiotensin II ; Angiotensin-converting enzyme inhibitors ; Angiotensin II receptor ; Angiotensin-converting enzyme 2.

### Introduction

The renin-angiotensin system, acting through the major effector peptide angiotensin II, has potent effects on the blood pressure, the water and sodium homeostasis and the end-organ damage in the heart, vessels, brain and kidneys.<sup>1-3)</sup> Consequently, inhibiting the RAS has been an important therapeutic strategy for the treatment of hypertension and the related end-organ damage.<sup>4-7)</sup> Although the RAS has been studied for more than a century, our understanding of its mechanisms remains a bit incomplete. ACE inhibitors and angiotensin II type 1 receptor blockers(ARBs) have many other beneficial effects in addition to decreasing the blood pressure.<sup>8-10)</sup> The transmembrane protease ACE2 has recently emerged as a negative regulator of the RAS as it counterbalances the multiple functions of ACE.<sup>11)12)</sup> This review will explore the molecules and pathways that are involved in the RAS, and we will identify potential novel therapeutic targets for cardiovascular diseases.

### Renin

Professor Robert Tigerstedt, a noted Finnish physio-

logist, discovered the renin system in 1898 at the Karolinska Institute in Sweden.<sup>13)</sup> Renin catalyzes the first, rate-limiting step in the RAS. Renin is an aspartyl proteolytic enzyme and its substrate is angiotensinogen.<sup>14)</sup> Renin cleaves a leucine-valine bond in the N-terminal region of human angiotensinogen or it cleaves a leucine-leucine bond in the angiotensinogens of other species to produce the decapeptide angiotensin I. Angiotensinogen is a large globular protein that's mainly derived from the pericentral zone of liver lobules. Juxtaglomerular cells(JG) cells of the afferent arteriole of the kidney are the locus for renin synthesis, storage and release. Translation of renin mRNA in these cells produces pre-prorenin, which in turn is then converted to prorenin by the removal of a single peptide and glycosylation. Some of the prorenin to renin conversion takes place in the juxtaglomerular cells, and both renin and prorenin are secreted from these cells. Prorenin is the more abundant circulating form of renin.<sup>15-17)</sup> Active renin is secreted in response to four regulatory mechanisms: the renal baroreceptor, the macula densa and the beta<sub>1</sub>-receptor stimulation via renal nerves and humoral factors. An intrarenal vascular stretch receptor in the afferent arterioles stimulates renin secretion in response to reduced renal perfusion pressure. When the macula densa senses a decrease of the distal tubular salt delivery, then a feedback signal initiates a series of steps that ultimately stimulate renin secretion. The JG cells are directly innervated by sympathetic nerves. Direct stimulation of


**Correspondence :** Wang Seong Ryu, MD, Department of Internal Medicine, College of Medicine, Chung-Ang University, Hangangro 3-ga, Yongsan-gu, Seoul 140-757, Korea  
Tel: 82-2-748-9882, Fax: 82-2-798-4745  
E-mail: w2283@nate.com

the renal nerves increases renin's release from the JG cells through a beta<sub>1</sub>-adrenergic mediated mechanism. Many humoral factors have also been implicated in the secretion of renin. The primary stimulatory intracellular "second messenger" for renin release is the cyclic nucleotide cAMP, which is probably the second messenger for stimulation via the beta-adrenergic nerves and prostaglandins.<sup>18)</sup>

Developing a renin inhibitor has been marred by such issues as potency, bioavailability, duration of action and the cost of synthesizing it. Aliskiren is the first representative member of a new class of non-peptide, orally active renin inhibitors. Clinical studies on hypertension patients have recently demonstrated dose-dependent reductions of the blood pressure with administering Aliskiren, and the drug's safety and tolerability were similar to those of the ARBs.<sup>19)</sup>

## Angiotensins

Angiotensins are peptide hormones that are produced by a series of proteolytic reactions starting with the cleavage of angiotensinogen by renin. Angiotensin I has no known biologic role aside from serving as the precursor of angiotensin II and III.<sup>20)</sup> Angiotensin II is the key effector hormone of the RAS, and it causes vasoconstriction, increased afterload and the retention of sodium and fluid. All of these actions affect the cardiovascular system and increase the blood pressure.<sup>21)</sup> Angiotensin II has been shown to stimulate VSMC growth, increase the expression of enzymes that produce mediators of inflammation such as phospholipase A<sub>2</sub>, and NAD(P)H oxidase, stimulate the JAK/STAT pathway and activate gene transcription of such proto-oncogenes as *c-fos*.<sup>22-25)</sup> Angiotensin II is produced both systematically and locally in the vessel wall by the actions of ACE, which cleaves angiotensin I to form angiotensin II. The importance of small GTP-binding proteins(G proteins) in angiotensin II signaling is becoming much clearer.<sup>2)</sup> Small G proteins are monomeric G proteins with a low molecular weight of between 20 to 40 kDa. A small G protein acts as a molecular switch that cycles between its inactive GDP-bound and active GTP-bound forms. Through the activation of small G proteins such as Ras, Rho, and Rac, angiotensin II induces vascular smooth muscle cell remodeling, including proliferation, migration and hypertrophy of the smooth muscle cells(Fig. 1).<sup>2,26-29)</sup> Although the circulating angiotensin II levels decrease in response to acute ACE inhibitor treatment, the circulating angiotensin II levels tend to increase in patients who are taking ACE inhibitors over long periods.<sup>30)</sup> The formation of angiotensin II, despite achieving effective ACE inhibition, may be explained by the activity of alternative enzymes that are capable of catalyzing the conversion of angiotensin I to angiotensin II. One such enzyme, which



**Fig. 1.** The Ras activation pathway by angiotensin II in the vascular smooth muscle cells. Angiotensin II stimulates the formation of Ras-GTP and the association of Ras-Raf. AT<sub>1</sub> was proposed to activate Ras through Gq/phospholipase C-mediated intracellular Ca<sup>2+</sup> elevation. A tyrosine kinase has been implicated in Ras activation by angiotensin II as well.<sup>2)</sup> Ang II: angiotensin II, AT<sub>1</sub>: angiotensin II type 1 receptor, EGFR: epidermal growth factor receptor, HB-EGF: heparin-binding EGF-like growth factor, ADAM: a disintegrin and metalloprotease, PLC: phospholipase-C, ROS: reactive oxygen species, GDP: guanosine diphosphate, GTP: guanosine triphosphate, MEK: mitogen-activated protein kinase (MAPK) kinase, ERK: extracellular signal-regulated kinase, MnK: MAPK interacting kinase, PI3K: phosphoinositide 3 kinase, PKB: protein kinase B, mTOR: mammalian target of rapamycin, PHAS-1: phosphorylated heat-and acid-stable protein, eIF4E: eukaryotic initiation factor-4E.


may be important in humans, is human heart chymase. Other enzymes known to catalyze this conversion include cathepsin G, and trypsin.<sup>31)</sup> In humans, direct comparison of the renal hemodynamic response to angiotensin II antagonist and ACE inhibitors has suggested that 30% to 40% of generating angiotensin II is non-ACE dependent when the renin system is stimulated by a low-salt diet. This percentage is even higher when the renin system is suppressed by a high-salt diet.<sup>32)</sup>

Data from the current well-documented studies on RAS's role in the development and maintenance of arterial hypertension has implicated angiotensin II in cardiovascular and renal pathologies, and these include cardiac left ventricular hypertrophy(LVH) and structural alterations of the vasculature, heart and kidney(e.g., neointima formation, post-infarct remodeling and nephrosclerosis).<sup>33)</sup> In this regard, angiotensin II has been recognized as a growth-promoting factor that contributes to structural alterations in various organs.<sup>34-37)</sup> Angio-

tensin III [Angiotensin(2-8)] is more lipid-soluble than is angiotensin II, and it is found in relatively high concentrations in the brain and cerebrospinal fluid. Angiotensin III is less potent than angiotensin II for eliciting biological responses. This is due both to the lower affinity of angiotensin III for the angiotensin II binding sites and also to the more rapid degradation of angiotensin III. Angiotensin IV [Angiotensin(3-8)] has effects on the brain that suggest it has a role in memory. It also stimulates the release of plasminogen activator inhibitor from endothelial cells.<sup>38</sup> Angiotensin(1-7) is a weak agonist that binds with low affinity to the AT<sub>1</sub> receptor, so the net effect of large amounts of that peptide is to inhibit the action of angiotensin II at that receptor. In addition, angiotensin(1-7) induces formation of nitric oxide and vasodilator prostaglandins, and it retards vascular smooth muscle growth via a mechanism that does not involve the AT<sub>1</sub> receptor.<sup>39</sup>

## ACE and ACE2

Angiotensin-converting enzyme(ACE) is a zinc metallopeptidase that's widely distributed on the surface of endothelial cells and epithelial cells.<sup>40</sup> Because of its dual role in regulating the levels of angiotensin II and bradykinin, the positive clinical effects of ACE inhibitors were thought to be the consequence of concomitant reductions in the production of angiotensin II and the degradation of bradykinin(Fig. 2).<sup>40,41</sup> Recent evidence has



**Fig. 2.** Action site of angiotensin-converting enzyme (ACE). ACE cleaves the C-terminal dipeptide from angiotensin I and bradykinin. Thus, ACE is strategically poised to regulate the balance between the renin-angiotensin system and the kallikrein-kinin system.<sup>40</sup>

**Table 1.** Pharmacology of the ACE inhibitors<sup>40</sup>

	Dosage (mg)	Route of elimination	Half-life (h)	Zinc ligand	Bioavailability (%)
Captopril	6.25-300	Kidney	1.7	Sulfhydryl	75-91
Enalapril	2.5-40	Kidney	11	Carboxyl	60
Lisinopril	5-40	Kidney	12	Carboxyl	6-60
Benazepril	5-80	Kidney	10-11	Carboxyl	>37
Quinapril	5-80	Kidney	1.9-2.5, 25 terminal	Carboxyl	>60
Ramipril	1.25-20	Kidney	Triphasic 4, 9-18, >50	Carboxyl	50-60
Trandolapril	1-8	Kidney, liver	15-24 terminal	Carboxyl	70
Fosinopril	10-80	Liver, kidney	12	Phosphinyl	36

ACE: angiotensin-converting enzyme

suggested that some of the beneficial effects of ACE inhibitors on cardiovascular function and homeostasis can be attributed to other mechanisms. These include the accumulation of the ACE substrate N-acetyl-seryl-aspartyl-lysyl-proline, which blocks collagen deposition in the injured heart, as well as the activation of the ACE signalling cascade, which involves the activation of kinase CK2 and the c-Jun N-terminal kinase in endothelial cells, and this leads to changes in gene expression.<sup>30</sup> It is clear that ACE inhibitors represent one of the major advances for cardiovascular therapeutics over the past 20 years (Table 1). They have proved effective for treating hypertension, they decrease the mortality of patients with congestive heart failure and left ventricular dysfunction after myocardial infarction, and they delay the progression of diabetic nephropathy.<sup>42</sup> The development of hypertensive left ventricular hypertrophy(LVH) is an early and important finding that may have direct pathophysiological implications on the progression from early hypertension to cardiovascular death.<sup>43</sup> It is well established that a reduction in blood pressure will reduce hypertensive LVH.<sup>44</sup> However, experimental evidence has suggested that various classes of antihypertensive agents have different effects on the left ventricular mass. A meta-analysis of the randomized controlled echocardiographic trials in humans, with the trials having a duration of 6 months or more, found that there were greater reductions in the left ventricular mass with administering ACE inhibitors than with administering beta-blocker, calcium antagonists or diuretics.<sup>45</sup>


The plasma ACE levels are stable when they are measured repeatedly in the same individual, whereas large inter-individual differences of these levels have been observed. This suggests that there is strong long term control of the ACE plasma levels, and this control may possibly have a genetic origin. Studies on the structure of the human ACE gene have revealed a polymorphism involving the presence(insertion, I) or absence(deletion, D) of a 287-bp sequence of DNA in intron 16.<sup>46-49</sup> The mean ACE activity levels in DD carriers were approximately twice that found in the individuals with the II genotype. Some investigations have suggested an association of the D allele with an increased risk for hyper-

tension, myocardial infarction or diabetic nephropathy, and the I allele has been associated with enhanced performance in athletes.<sup>12)41)</sup>

Genome-based strategies have identified a homologue of ACE with 42% sequence homology. Like ACE, this enzyme, which is named ACE2, is also a zinc-containing membrane-associated exopeptidase.<sup>12)</sup> ACE2 cleaves only a single C-terminal amino acid. It converts angiotensin II to angiotensin(1-7) and angiotensin I to angiotensin(1-9). ACE2 was initially found to be expressed in endothelial cells of the heart and in the tubular epithelial cells of the kidney. Subsequent studies have shown that the ACE2 gene expression also occurs in the gastrointestinal tract and to a lesser extent, in other organs such as lungs. Angiotensin(1-7) appears to be the main effector peptide of ACE2, and it has vasodilatory, natriuretic and antiinflammatory properties. A new regulator has entered the established metabolic RAS pathways with the discovery of ACE2(Fig. 3).<sup>19)</sup>

## Angiotensin II Receptors

Angiotensin II binds to at least three known high-affinity receptor subtypes that are termed AT<sub>1</sub>, AT<sub>2</sub> and




**Fig. 3.** Overview of the renin-angiotensin pathway. After its formation from angiotensinogen by renin, angiotensin I can be further cleaved by angiotensin-converting enzyme (ACE) to yield angiotensin II, or it is alternatively converted by ACE2 to angiotensin (1-9). BK: bradykinin, NO: nitric oxide.<sup>19)</sup>

AT<sub>4</sub>, and this converts the interaction with angiotensin II into a cellular response via signal transduction. The AT<sub>1</sub> and AT<sub>2</sub> receptors have similar affinities for angiotensin II, but they have distinctly different effects.<sup>1)</sup> The AT<sub>1</sub> is the major angiotensin II receptor expressed in adults. Angiotensin II binds to the AT<sub>1</sub> receptor and this activates G protein-coupled phospholipase C(PLC). PLC is an essential enzyme for transmembrane signal transduction by generating second messenger molecules like inositol triphosphate(IP<sub>3</sub>) and diacylglycerol(DG). IP<sub>3</sub> induces the release of Ca<sup>2+</sup> from internal stores, and DG is the physiological activator of protein kinase.<sup>17)48)</sup> On the other hand, angiotensin II binding to the AT<sub>2</sub> receptor activates a counterregulatory pathway to induce vasorelaxation via activation of the kinin/NO/cGMP system.<sup>49)50)</sup> Distribution of the AT<sub>1</sub> receptor in the adult is ubiquitous, including the vasculature, kidney, adrenal gland, heart, liver and brain. With some exceptions, most of the pathologic cardiovascular effects of angiotensin II are mediated through the AT<sub>1</sub> receptor: hypertension, coagulation, inflammation and vascular smooth muscle cell growth.<sup>2)</sup> Several AT<sub>1</sub> receptor blockers have become available for the treatment of hypertension or congestive heart failure(Table 2). In contrast, the AT<sub>2</sub> receptor is mostly involved in developmental processes in the fetus; it is abundant in the fetal brain, kidney and other sites. The AT<sub>2</sub> receptor reappears in adults, and it generally acts as a counterweight to the pathologic effects of AT<sub>1</sub>; it inhibits angiotensin II-induced growth and proinflammatory activity, and it decreases the blood pressure. The AT<sub>1</sub> and AT<sub>2</sub> receptors have both been cloned. They belong to the superfamily of G-protein-coupled receptors that contain 7 transmembrane regions. Their amino acid sequence seems to be highly conserved across species and across the tissues within a species.<sup>50)</sup> AT<sub>1</sub> and AT<sub>2</sub> receptors share only 34% homology and they have distinct signal transduction pathways. In rodents, AT<sub>1</sub> receptor have been further subdivided into AT<sub>1A</sub> and AT<sub>1B</sub>.<sup>51)</sup> In amphibians and in neuroblastoma cell lines, an angiotensin II receptor that was not inhibited by either losartan or PD 123177 has been classified as AT<sub>3</sub>. Evidence suggests that the AT<sub>4</sub> receptor, via angiotensin IV lan-

**Table 2.** Pharmacokinetic properties of the angiotensin II receptor blockers<sup>51)</sup>

Drug (active metabolite)	Dosage (mg/day)	Active metabolite	Half-Life (h)	Food effect	Bioavailability (%)
Losartan (EXP3174)	50-100	Yes	2 (6-9)	No	33
Valsartan	80-320	No	9	Yes, -40%	25
Candesartan (TCV116) (CV11974)	4-16 (32)	Yes	3.5-4 3-11	No	... 42
Irbesartan	150-300	No	11-15	No	70
Eprosartan	200-800	No	5-7	No	15
Telmisartan	40-80	No	24	No	43
Olmesartan	10-40	Yes	13	No	26



**Fig. 4.** Pathways of angiotensin II formation and the proposed role of angiotensin II receptors. ACE: angiotensin-converting enzyme, AT<sub>1</sub>R, AT<sub>2</sub>R, AT<sub>3</sub>R, AT<sub>4</sub>R: angiotensin II types 1, 2, 3 and 4 receptors, CAGE: chymostatin-sensitive angiotensin II-generating enzyme, PAI-1: plasminogen activator inhibitor-1.<sup>1)</sup>

giotensin(3-8)], is an important mediator of the expression of plasminogen activator inhibitor-1(Fig. 4).<sup>51)</sup>

## Conclusion

This review summarizes the molecules and pathways of the RAS. Therapeutic agents that block the RAS at several points along its pathway have significantly reduced the cardiovascular and renal morbidity and mortality. Both ACE and ACE2 are involved in the production of the biologically active peptides angiotensin II and angiotensin(1-7) from angiotensin I. There are currently ongoing studies that will elucidate the physiological roles of ACE2 in myocardial function and its contribution to kidney damage. It still remains to be seen whether more complete blockade of the RAS will improve the clinical outcomes of patients suffering with cardiovascular diseases.

## REFERENCES

- Duprez DA. Role of the renin-angiotensin-aldosterone system in vascular remodeling and inflammation: a clinical review. *J Hypertens* 2006;24:983-91.
- Ohtsu H, Suzuki H, Nakashima H, et al. Angiotensin II signal transduction through small GTP-binding proteins: mechanism and significance in vascular smooth muscle cells. *Hypertension* 2006;48:534-40.
- Kim KB. The role of angiotensin receptor antagonist in patients with hypertension and cardiovascular diseases. *Korean Circ J* 1999;29:335-40.
- Dzau V. The cardiovascular continuum and renin-angiotensin-aldosterone system blockade. *J Hypertens Suppl* 2005;23:S9-17.
- Burnier M, Zanchi A. Blockade of the renin-angiotensin-aldosterone system: a key therapeutic strategy to reduce renal and cardiovascular events in patients with diabetes. *J Hypertens* 2006;24:11-25.
- Weber MA. Hypertension treatment and implications of recent cardiovascular outcome trials. *J Hypertens Suppl* 2006;24:S37-44.
- Sokol SI, Portnay EL, Curtis JP, et al. Modulation of the renin-angiotensin-aldosterone system for the secondary prevention of stroke. *Neurology* 2004;63:208-13.
- Sleight P, Yusuf S. New evidence on the importance of the renin-angiotensin system in the treatment of higher-risk patients with hypertension. *J Hypertens* 2003;21:1599-608.
- Jeon ES. Role of angiotensin II receptor blockers in the treatment of congestive heart failure. *Korean Circ J* 2002;32:1039-45.
- Mancia G. Total cardiovascular risk: a new treatment concept. *J Hypertens Suppl* 2006;24:S17-24.
- Danilezyk U, Penninger JM. Angiotensin-converting enzyme II in the heart and the kidney. *Circ Res* 2006;98:463-71.
- Farmer JA. Renin angiotensin system and ASCVD. *Curr Opin Cardiol* 2000;15:141-50.
- Ulfendahl HR, Aurell M. Renin-Angiotensin: a centenary symposium of the discovery of the renin-angiotensin system. London: Portland Press; 1998. p.1-12.
- Suh SY, Park CG, Chwe UR, et al. Plasma renin activity and clinical implication in Korean hypertensive patients. *Korean Circ J* 2005;35:658-64.
- Duff JL, Marrero MB, Paxton WG, Schieffer B, Bernstein KE, Berk BC. Angiotensin II signal transduction and the mitogen-activated protein kinase pathway. *Cardiovasc Res* 1995;30:511-7.
- Pan L, Gross KW. Transcriptional regulation of renin. *Hypertension* 2005;45:3-8.
- Rosendorff C. The renin-angiotensin system and vascular hypertrophy. *J Am Coll Cardiol* 1996;28:803-12.
- Beierwaltes WH. Renin synthesis and secretion. In: Izzo JL, Black HR, editor. *Hypertension Primer*. 3rd ed. Philadelphia: Lippincott Williams & Wilkins; 2003. p.14-7.
- Krum H, Gilbert RE. Novel therapies blocking the renin-angiotensin-aldosterone system in the management of hypertension and related disorders. *J Hypertens* 2007;25:25-35.
- Berk BC, Corson MA. Angiotensin II signal transduction in vascular smooth muscle: role of tyrosine kinases. *Circ Res* 1997;80:607-16.
- de Leeuw PW. How do angiotensin II receptor antagonists affect blood pressure? *Am J Cardiol* 1999;84:5K-6K.
- Min LJ, Mogi M, Li JM, Iwanami J, Iwai M, Horiuchi M. Aldosterone and angiotensin II synergistically induce mitogenic response in vascular smooth muscle cells. *Circ Res* 2005;97:434-42.
- Zhang Y, Griendling KK, Dikalova A, Owens GK, Taylor WR. Vascular hypertrophy in angiotensin II-induced hypertension is mediated by vascular smooth muscle cell-derived H<sub>2</sub>O<sub>2</sub>. *Hypertension* 2005;46:732-7.
- Ushio-Fukai M, Zuo L, Ikeda S, Tojo T, Patrushev NA, Alexander RW. cAb1 tyrosine kinase mediates reactive oxygen species and caveolin-dependent AT<sub>1</sub> receptor signaling in vascular smooth muscle. *Circ Res* 2005;97:829-36.
- Dikalova A, Clempus R, Lassegue B, et al. Nox1 overexpression potentiates angiotensin II-induced hypertension and vascular smooth muscle hypertrophy in transgenic mice. *Circulation* 2005;112:2668-76.
- Shimokawa H. Rho-kinase as a novel therapeutic target in treatment of cardiovascular diseases. *J Cardiovasc Pharmacol* 2002;39:319-27.
- Calo LA, Pessina AC. RhoA/Rho-kinase pathway: much more than just a modulation of vascular tone: evidence from studies in humans. *J Hypertens* 2007;25:259-64.
- Smith NJ, Luttrell LM. Signal switching, crosstalk, and arrestin scaffolds: novel G protein-coupled receptor signaling in cardiovascular disease. *Hypertension* 2006;48:173-9.

- 29) Yang CH, Lee SH, Park TH, et al. *Gene expression profile of volume-overloaded human ventricular myocardium prior to developing heart failure*. Korean Circ J 2005;35:649-57.
- 30) Fleming I. *Signaling by the angiotensin-converting enzyme*. Circ Res 2006;98:887-96.
- 31) Willenheimer R, Dahloef B, Rydberg E, Erhardt L. *AT<sub>1</sub>-receptor blockers in hypertension and heart failure: clinical experience and future directions*. Eur Heart J 1999;20:997-1008.
- 32) Hollenberg NK. *Hypertension, small arteries, and pathways for angiotensin II generation*. Circulation 2000;101:1641-2.
- 33) Unger T, Culman J, Peter G. *Angiotensin II receptor blockade and end-organ protection: pharmacological rationale and evidence*. J Hypertens Suppl 1998;16:S3-9.
- 34) Diez J. *Profibrotic effects of angiotensin II in the heart*. Hypertension 2004;43:1164-5.
- 35) Wolf G, Wenzel UO. *Angiotensin II and cell cycle regulation*. Hypertension 2004;43:693-8.
- 36) Kim YH, Son JH, Piao H, et al. *Role of angiotensin II type 1A receptor-mediated signaling in the development of pressure overload-induced left ventricular hypertrophy and fibrosis*. Korean Circ J 2006;36:626-34.
- 37) Kim KM, Kim IS, Park SJ, et al. *Effects of angiotensin II on the growth of vascular smooth muscle cells*. Korean Circ J 1999;29:209-15.
- 38) Dielis AW, Smid M, Spronk HM, et al. *The prothrombotic paradox of hypertension: role of the renin-angiotensin and kallikrein-kinin systems*. Hypertension 2005;46:1236-42.
- 39) Sampaio WO, Souza dos Santos RA, Faria-Silva R, de Mata Machado LT, Schiffrin EL, Touyz RM. *Angiotensin-(1-7) through receptor Mas mediates endothelial nitric oxide synthase activation via Akt-dependent pathways*. Hypertension 2007;49:185-92.
- 40) Brown NJ, Vaughan DE. *Angiotensin-converting enzyme inhibitors*. Circulation 1998;97:1411-20.
- 41) Sayed-Tabatabaei FA, Oostra BA, Isaacs A, van Duijn CM, Witteman JC. *ACE polymorphisms*. Circ Res 2006;98:1123-33.
- 42) Goldberg LR, Jessup M. *Stage B heart failure: management of asymptomatic left ventricular systolic dysfunction*. Circulation 2006;113:2851-60.
- 43) Kahan T. *The importance of left ventricular hypertrophy in human hypertension*. J Hypertens Suppl 1998;16:S23-9.
- 44) Alfakin K, Reid S, Hall A, Sivananthan MU. *The assessment of left ventricular hypertrophy in hypertension*. J Hypertens 2006;24:1223-30.
- 45) Gosse P. *Left ventricular hypertrophy as a predictor of cardiovascular risk*. J Hypertens Suppl 2005;23:S27-33.
- 46) Rigat B, Hubert C, Alhenc-Gelas F, Cambien F, Corvol P, Soubrier F. *An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels*. J Clin Invest 1990;86:1343-6.
- 47) Rigat B, Hubert C, Corvol P, Soubrier F. *PCR detection of the insertion/deletion polymorphism of the human angiotensin-converting enzyme gene (DCP1) (dipeptidyl carboxypeptidase I)*. Nucleic Acids Res 1992;20:1433.
- 48) Rosendorff C. *Vascular hypertrophy in hypertension: role of the renin-angiotensin system*. Mt Sinai J Med 1998;65:108-17.
- 49) Schiffrin EL. *Effects of antihypertensive drugs on vascular remodeling: do they predict outcome in response to antihypertensive therapy?* Curr Opin Nephrol Hypertens 2001;10:617-24.
- 50) Schulman IH, Zhou MS, Raij L. *Cross-talk between angiotensin II receptor types 1 and 2: potential role in vascular remodeling in humans*. Hypertension 2007;49:270-1.
- 51) Burnier M. *Angiotensin II type 1 receptor blockers*. Circulation 2001;103:904-12.