

An Assay of Measuring Platelet Reactivity Using Monoclonal Antibody against Activated Platelet Glycoprotein IIb/IIIa in Patients Taking Clopidogrel

Seung-Jae Joo, MD¹, Joon-Hyoun Choi, MD¹, Song-Yi Kim, MD¹, Ki-Seok Kim, MD¹, Young Ree Kim, MD², and Sung Ha Kang, MD²

¹Department of Cardiology, and ²Laboratory Medicine, Jeju National University Hospital, Jeju, Korea

Background and Objectives: Residual platelet reactivity in patients who are taking clopidogrel is commonly measured with VerifyNow assay, which is based on the principle of light transmission aggregometry. However, to evaluate the residual platelet reactivity, it would be more accurate if the reactivity of platelet glycoprotein (GP) IIb/IIIa is directly monitored. In this study, PAC1, a monoclonal antibody against activated platelet GP IIb/IIIa, was used to measure the residual platelet reactivity.

Subjects and Methods: Twenty seven patients with coronary artery disease taking clopidogrel were enrolled. Platelets in whole blood were stained with fluorescein isothiocyanate (FITC)-conjugated PAC1. Mean fluorescence intensity (MFI) and % positive platelets (PP) were measured with flow cytometry, and the binding index (BI; MFI x %PP/100) was calculated. P2Y12 reaction unit (PRU) and % inhibition of VerifyNow assay were also measured in the usual manner.

Results: PRU of VerifyNow assay correlated significantly with MFI, %PP, and BI at 10 μ M ($r=0.59$, 0.73 , and 0.60 , respectively, all $p<0.005$) and 20 μ M of adenosine diphosphate (ADP; $r=0.61$, 0.75 , and 0.63 , respectively, all $p<0.005$). The % inhibition also correlated significantly with MFI, %PP, and BI at 10 μ M ($r=-0.60$, -0.69 , and -0.59 , respectively, all $p<0.005$) and 20 μ M of ADP ($r=-0.63$, -0.71 , and -0.62 , respectively, all $p<0.005$).

Conclusion: Direct measurements of the reactivity of platelet GP IIb/IIIa were feasible using PAC1 and flow cytometry in patients taking clopidogrel. Further clinical studies are required to determine the cut-off values which would define high residual platelet reactivity in patients on this treatment protocol. (**Korean Circ J 2015;45(5):378-385**)

KEY WORDS: Blood platelets; Glycoprotein IIb/IIIa; Platelet function test; Flow cytometry; Clopidogrel.

Introduction

Although platelet activation and aggregation is an essential part of hemostasis, it also initiates acute coronary syndrome or thrombotic complications related to percutaneous coronary stent

implantation. Dual antiplatelet therapy, including aspirin and P2Y12 inhibitors, is usually recommended in patients with acute myocardial infarction or unstable angina, especially those who have undergone percutaneous coronary intervention (PCI) with drug-eluting stents.¹⁻³⁾ One of the most commonly used P2Y12 inhibitors is clopidogrel, which needs to be metabolized in vivo to become an active drug. However, individual response to oral clopidogrel to inhibit P2Y12 receptor is variable,^{4,5)} and, despite taking clopidogrel, high residual platelet reactivity in patients with PCI has been associated with death, myocardial infarction, or stent thrombosis.⁶⁾

Platelet function tests such as light transmission aggregometry (LTA), VerifyNow P2Y12 assay, platelet function analyser, or flow cytometric analysis of vasodilator-stimulated phosphoprotein (VASP) phosphorylation or P-selectin are used to measure on-treatment high residual platelet reactivity;⁴⁻⁹⁾ however, no single test can evaluate the complex mechanisms of platelet activation and aggregation.⁴⁾ In patients undergoing coronary stent implantation,

Received: March 2, 2015

Revision Received: April 10, 2015

Accepted: April 28, 2015

Correspondence: Seung-Jae Joo, MD, Department of Cardiology, Jeju National University Hospital, 15 Aran 13-gil, Jeju 63241, Korea
Tel: 82-64-717-1701, Fax: 82-64-725-2593
E-mail: sejjoo@jejunu.ac.kr

• The authors have no financial conflicts of interest.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

the diagnostic accuracy of each test to predict cardiovascular events was not high.¹⁰⁾

Activation and prothrombin binding of platelet glycoprotein (GP) IIb/IIIa is a final common pathway of platelet aggregation.^{11,12)} If the reactivity of platelet GP IIb/IIIa is directly measured, it would be a more accurate assay to evaluate the residual platelet reactivity. PAC1, a monoclonal antibody having high affinity to activated platelet GP IIb/IIIa,¹³⁻¹⁵⁾ was used to monitor the effect of GP IIb/IIIa antagonists on platelet activation.¹⁶⁻¹⁸⁾ However, the direct measurement of GP IIb/IIIa activation with PAC1 to assess the residual platelet reactivity in patients taking clopidogrel has not yet been systemically performed. Feasibility of flow cytometric analysis using PAC1 in whole blood to measure on-treatment residual platelet reactivity was investigated in this study.

Subjects and Methods

Study patients

A total of 27 patients with coronary artery disease, who were

taking clopidogrel 75 mg per day for at least 7 days, or for at least 48 hours after the 300-600 mg loading dose, were included in this study, after acquiring their written informed consent. The study protocol was approved by the institutional review board of Jeju National University Hospital.

Reagents

Sodium chloride (NaCl; Prod. No. S3014), potassium chloride (KCl; Prod. No. P9541), magnesium chloride (MgCl₂; Prod. No. M8266), dextrose (Prod. No. D9434), bovine serum albumin (BSA; Prod. No. A2513), 4-(2-hydroxyethyl)-1-piperazineethane-sulfonic acid (HEPES; Prod. No. H3375), paraformaldehyde (Prod. No. P6148), adenosine 5'-diphosphate (ADP; Prod. No. A2754), and prostaglandin I₂ (PGI₂; Prod. No. P6188) were purchased from Sigma Chemical Co. (St. Louis, MO, USA). Monoclonal antibodies, fluorescein isothiocyanate (FITC)-conjugated PAC1 (Cat. No. 340507) and FITC-conjugated mouse IgM, κ isotype (Cat. No. 555583), were from Becton Dickinson Co. (San Jose, CA, USA). PE-conjugated anti-CD41 (Prod. No. R7058), and phycoerythrin (PE)-conjugated mouse IgG₁ (Prod. No. X0928) were




Fig. 1. Flow cytometric analysis of whole blood. (A) Platelets were separated from erythrocytes and white blood cells, on the basis of their characteristic forward- and side- scatter profiles, and a narrow gate was placed around the platelets. (B) Staining of resting platelets with PE-conjugated anti-CD41, a monoclonal antibody against glycoprotein IIb. (C) Staining of platelets with control FITC-conjugated mouse IgMκ. (D) Staining of platelets with FITC-conjugated PAC1. Antibody-positive platelets were defined as those with a fluorescence intensity >99.0% to 99.5%, of non-stimulated platelets stained with FITC-conjugated mouse IgM κ. SSC: side-scatter, FSC: forward-scatter, FITC: fluorescein isothiocyanate, PP: positive platelet.

from DAKO Co. (Glostrup, Denmark).

Blood sampling

Whole blood was withdrawn in the morning, from an antecubital vein using a 21-gauge needle. After discarding the first 2 mL, the blood was collected into a sodium citrate-coated tube, and was processed within 30 minutes of collection.

VerifyNow P2Y12 assay

Whole blood was processed according to the user's manual of VerifyNow system (Accumetrics, San Diego, CA, USA). P2Y12 reaction unit (PRU), measured in ADP/PGE1 channel, reports ADP-mediated aggregation specific to P2Y12 receptor. BASE is an independent measurement based on platelet aggregation from thrombin receptors, and protease activated receptors (PAR) PAR-1 and PAR-4. Percent inhibition (%) was the percent change from baseline aggregation, and was calculated from PRU and BASE using the following equation: % inhibition = (BASE - PRU) × 100 / BASE.

Flow cytometry

In a mixture consisting of 5 µL of ADP and 25 µL of isotonic HEPES buffer (137 mM NaCl, 2.7 mM KCl, 1 mM MgCl₂, 5.6 mM dextrose, 1 mg/mL BSA, 20 mM HEPES, pH 7.4), 15 µL FITC-conjugated PAC1 (final concentration 60 µg/mL) was incubated for 15 minutes with 5 µL of whole blood. Total volume of the reaction mixture was 50 µL, and the final concentration of ADP was either 1, 10, or 20 µM. To measure the baseline activation of GP IIb/IIIa at ADP 0 µM, 1 µM PGI₂ was added instead of ADP. FITC-conjugated mouse IgM, κ isotype (IgMκ), was used as a control antibody. For staining of the total active and inactive GP IIb/IIIa, 5 µL of whole blood was incubated for 15 minutes with 10 µL of PE-conjugated anti-CD41, 5 µL of PGI₂ (final concentration 1 µM), and 30 µL of isotonic HEPES buffer. The reaction was stopped by adding 500 µL of HEPES buffer. After fixation with 500 µL of 1% paraformaldehyde, the samples were stored at 4°C in the dark, until flow cytometric analysis.

Flow cytometric analysis was performed with BD FACSCalibur (San Jose, CA, USA). Platelets were distinguished from erythrocytes and white blood cells as described by Shattil et al.¹⁴ Briefly, platelets were separated in whole blood by their characteristic forward- and side-scatter profiles. A gate was set around the platelets, and 10000 cells were analyzed to quantify the amount of platelet-bound FITC-PAC1 (Fig. 1A). Over 90% of cells in this gate were platelets identified by PE-anti-CD41 (Fig. 1B). The FITC-PAC1 binding was expressed as mean fluorescence intensity (MFI), percentage of positive platelets for antibody (%PP), or binding index (BI). Antibody-positive platelets were defined as those having a

fluorescence intensity >99.0% to 99.5% of non-stimulated platelets stained with FITC-conjugated mouse IgMκ (Fig. 1C and D).¹⁴ Binding index was calculated as MFI/%PP/100.

Statistics

Values are expressed as mean ± standard deviation or, where

Table 1. Baseline characteristics of patients (N=27)

Variables	
Age (years)	65.4±9.0
Male	17 (63)
Height (cm)	159.3±7.6
Weight (kg)	62.2±9.6
Body mass index (kg/m ²)	24.5±3.2
Smoker	8 (30)
Hypertension	11 (41)
Diabetes mellitus	5 (19)
Chronic kidney disease	3 (11)
Previous myocardial infarction	7 (26)
Clinical diagnosis	
Acute myocardial infarction	6 (22)
Unstable angina	2 (8)
Stable angina	19 (70)
Laboratory findings	
Hemoglobin (g/dL)	13.1±2.1
Platelet count (×10 ³ /µL)	197.2±51.4
Mean platelet volume (fL)	9.9±0.8
Total cholesterol (mg/dL)	158.4±25.6
Triglyceride (mg/dL)	106.0±57.7
HDL-C (mg/dL)	46.0±10.6
LDL-C (mg/dL)	82.0±26.2
Creatinine (mg/dL)	1.5±1.6
Medications	
Aspirin	27 (100)
Clopidogrel	27 (100)
Cilostazol	1 (4)
Beta-blockers	16 (59)
Calcium channel blockers	10 (37)
Angiotensin-converting enzyme inhibitors	5 (19)
Angiotensin receptor blockers	5 (19)
Diuretics	1 (4)
Nitrate	7 (26)
Statins	26 (96)

Data are expressed as number (%) or mean ± standard deviation. HDL-C: high density lipoprotein-cholesterol, LDL-C: low density lipoprotein-cholesterol

appropriate, mean±standard error of mean. Changes of MFI, %PP or binding index after stimulation with increasing concentration of ADP, were compared by paired t-test. Correlations between variables were evaluated by Pearson's correlation. Values were considered to be statistically significant, when $p < 0.05$. All statistical analyses were performed using SPSS for window version 12 (SPSS Inc. Chicago, IL, USA).

Results

Baseline characteristics of patients are shown in Table 1. Mean

age was 65.4 ± 9.0 years. Clinical diagnoses were acute myocardial infarction in 6 patients, unstable angina in 2 patients, and stable angina in 19 patients. Platelet count was $197.2 \pm 51.4 \times 10^3/\mu\text{L}$, and mean platelet volume was 9.9 ± 0.8 fL. All patients were on aspirin and clopidogrel treatment.

Examples of flow cytometric data and analysis are demonstrated in Fig. 2. In a patient with low residual platelet reactivity (PRU 101 and % inhibition 48%), the MFI, %PP, and binding index were 607, 57%, and 347 respectively, after ADP stimulation at $20 \mu\text{M}$ (Fig. 2A). In contrast, a patient with high residual platelet reactivity (PRU 297 and % inhibition 19%) showed much greater MFI, %PP, and binding




Fig. 2. Examples of flow cytometric data and analysis in a patient with (A) low residual platelet reactivity (B) high residual platelet reactivity. A patient with high residual platelet reactivity showed much greater mean fluorescence intensity (MFI), % positive platelets (%PP), and binding index (BI) at $10 \mu\text{M}$ or $20 \mu\text{M}$ of ADP. ADP: adenosine diphosphate, FITC: fluorescein isothiocyanate.




Fig. 3. Changes of mean fluorescence intensity (MFI), % positive platelets, and binding index after ADP stimulation. Values are mean±standard error of mean. They reached maximal values at $10 \mu\text{M}$ concentration. ADP: adenosine diphosphate.




Fig. 4. Correlation between mean fluorescence intensity (MFI), % positive platelets, and binding index with VerifyNow PRU at (A) 10 μ M or (B) 20 μ M of ADP. PRU: P2Y12 reaction unit, ADP: adenosine diphosphate.

index at 20 μ M of ADP (2235, 88%, and 1959, respectively) (Fig. 2B). MFI, %PP, and binding index increased after ADP stimulation in a dose-dependent manner, and reached maximal values at 10 μ M concentration (Fig. 3). MFI, %PP, or binding index correlated significantly with VerifyNow PRU at 10 μ M ($r=0.59$, 0.73 , and 0.60 respectively, all $p<0.005$) and 20 μ M of ADP ($r=0.61$, 0.75 , and 0.63 respectively, all $p<0.005$) (Fig. 4). There were also significant correlations between P2Y12 % inhibition and MFI, %PP, or binding index after ADP stimulation at 10 μ M ($r=-0.60$, -0.69 , and -0.59 respectively, all $p<0.005$) and 20 μ M ($r=-0.63$, -0.71 , and -0.62 respectively, all $p<0.005$) (Fig. 5).

Discussion

This study showed that a direct measurement of the reactivity of platelet GP IIb/IIIa using PAC1 and flow cytometry was possible to assess on-treatment residual platelet reactivity in patients taking clopidogrel. The MFI, %PP, or BI of this assay had a significant correlation with PRU or % inhibition of VerifyNow assay.

GP IIb/IIIa is the most abundant receptor among all the platelet membrane proteins. The most important characteristic of GP IIb/IIIa is its affinity modulation of binding to fibrinogen, the main ligand involved in platelet aggregation. Low affinity state for fibrinogen binding in resting platelets becomes high affinity state after activation by agonists, such as ADP, thrombin or arachidonic acid.¹¹⁾¹²⁾ PAC1, which binds to the activated form of GP IIb/IIIa but not to the resting platelets,¹³⁻¹⁵⁾ is a suitable murine monoclonal antibody for evaluating platelet reactivity. In this study, PAC1-binding to platelets was very low in resting platelets, but increased dramatically after ADP stimulation, especially in on-treatment patients with high residual platelet reactivity.

Several clinical tests are used to measure the residual platelet reactivity in patients taking clopidogrel. LTA is a historical gold standard test which measures the change of turbidity after platelets aggregate by agonist-stimulation. Although LTA is the most widely used method to monitor the effects of antiplatelet agents and predict clinical outcomes, it requires large sample volumes, immediate separation of platelet-rich and platelet-poor plasma




Fig. 5. Correlation between mean fluorescence intensity (MFI), % positive platelets, and binding index with VerifyNow P2Y12 % inhibition at (A) 10 μM or (B) 20 μM of ADP. ADP: adenosine diphosphate.

from whole blood, lengthy processing time, and a trained technician, all of which result in variable reproducibility, making it difficult for a point-of-care method in daily clinical situations.⁴⁷⁾ Presently, VerifyNow P2Y12 assay has been the most commonly used point-of-care system to measure the residual platelet reactivity in clinical studies. Because it does not need the separation of platelet-rich plasma, and it correlated with ADP-stimulated platelet aggregation by LTA,⁴⁾ high PRU has been the most frequently used criteria in many clinical studies of patients with coronary stents to predict clinical outcomes.⁶⁾⁹⁾¹⁹⁾ However, diagnostic accuracy of high platelet reactivity during on-treatment, defined by PRU \geq 236 to predict 1-year clinical outcomes, was only 13% in patients undergoing elective coronary stent implantation and taking clopidogrel.¹⁰⁾ In Gauging Responsiveness with A VerifyNow assay-Impact on Thrombosis And Safety (GRAVITAS) trial, PRU-guided high-dose clopidogrel therapy among patients with high on-treatment platelet reactivity (PRU \geq 230) after PCI with drug-eluting stents, did not reduce fatalities from cardiovascular causes, non-fatal myocardial infarction, or stent thrombosis, when compared with standard-dose

clopidogrel.²⁰⁾

Flow cytometry is also a valuable tool to evaluate platelet reactivity. P-selectin expression on platelet surface, or platelet-leukocyte aggregate, was monitored by flow cytometry to evaluate the inhibitory effect of clopidogrel.⁸⁾²¹⁾ Flow cytometric assessment of VASP phosphorylation, which measures activation-dependent intracellular signaling of P2Y12 receptor, has been frequently used to differentiate high on-treatment platelet reactivity.¹⁹⁾²²⁾ The activated state of platelet GP IIb/IIIa was evaluated to monitor the treatment effect of GP IIb/IIIa antagonists.¹⁶⁻¹⁸⁾

In patients taking clopidogrel, direct measurement of platelet GP IIb/IIIa activation by ADP may be a more valuable tool to monitor the residual platelet reactivity. In this study, activation of GP IIb/IIIa detected by flow cytometric analysis of PAC1-binding to platelets correlated significantly with both PRU and % inhibition of VerifyNow P2Y12 assay, especially at ADP 20 μM, which was the same concentration included in ADP channel of the VerifyNow system. Although flow cytometry was calibrated every day before sample running, MFI might be influenced by variations during

staining process of fluorescent PAC1 with platelets. PAC1-binding platelets were defined as those with fluorescence intensity >99.0% to 99.5% of non-stimulated platelets, processed in the presence of 1 μ M PGI₂. This index (%PP), is relatively independent on daily sample processing and measurement, and may be more accurate than MFI. Notwithstanding, MFI as well as %PP showed good correlations with PRU or % inhibition of VerifyNow system, which suggests both indices can be used to measure the reactivity of GP IIb/IIIa.

Although whole blood flow cytometry has advantages such as low sample volume or the non-requirement of centrifugation or washing, it also has several drawbacks.⁴⁾⁷⁾ It needs expensive flow cytometric analysis and experienced staff. Clear separation of platelets from erythrocytes and white blood cells is an essential requirement in flow cytometric analysis of whole blood. Platelets in whole blood could easily be discriminated by their characteristic small forward- and side-scatter profiles. The two-color method may more accurately separate platelets in whole blood. However, considering that an earlier study showed similar flow cytometric results when either one-color or two-color method was used,¹⁴⁾ and since the one-color method (which is also used in VASP phosphorylation assay)²³⁾ is more easily applicable considering the cost or processing time, in the present study the platelets were distinguished in the whole blood without a second antibody staining. Over 90% of cells in the platelet gate were platelets, as confirmed by anti-CD41, a monoclonal antibody against GP IIb. A time delay during PAC1 staining and fixation may lead to spontaneous activation of the platelets.¹³⁾ To minimize it, whole blood was processed within 30 minutes after sampling from the body; it was not difficult to maintain this sample processing time if all the reagents were ready in a tube before blood sampling.

This study has several limitations. First, cut-off values to define high on-treatment residual platelet reactivity could not be determined due to the small number of patients. A further clinical study to determine an optimal cut-off point is needed. Second, this assay may be expensive because it requires costly monoclonal antibodies and flow cytometry. However, the total cost of this method at our institute was comparable to that of VerifyNow assay, and VASP phosphorylation assay using flow cytometry is frequently used in clinical studies. Third, a need for sample processing without any uncertainty by an experienced staff may make it difficult for it to be used as a point-of-care assay.

In conclusion, direct measurements of the reactivity of platelet GP IIb/IIIa were feasible using PAC1 and flow cytometry. Cut-off values of this assay to define high on-treatment residual platelet reactivity need to be determined in a further clinical study.

Acknowledgements

This study was supported by a research grant from The Korean Society of Cardiology in 2011.

References

1. Amsterdam EA, Wenger NK, Brindis RG, et al. 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: executive summary: a report of the American College of Cardiology/American Heart Association task force on practice guidelines. *Circulation* 2014;130:2354-94.
2. O'Gara PT, Kushner FG, Ascheim DD, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: executive summary: a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines. *Circulation* 2013;127:529-55.
3. Levine GN, Bates ER, Blankenship JC, et al. 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention: executive summary: a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines and the Society for Cardiovascular Angiography and Interventions. *Circulation* 2011;124:2574-609.
4. Gurbel PA, Becker RC, Mann KG, Steinhubl SR, Michelson AD. Platelet function monitoring in patients with coronary artery disease. *J Am Coll Cardiol* 2007;50:1822-34.
5. Kulickowski W, Witkowski A, Polonski L, et al. Interindividual variability in the response to oral antiplatelet drugs: a position paper of the Working Group on antiplatelet drugs resistance appointed by the Section of Cardiovascular Interventions of the Polish Cardiac Society, endorsed by the Working Group on Thrombosis of the European Society of Cardiology. *Eur Heart J* 2009;30:426-35.
6. Brar SS, ten Berg J, Marcucci R, et al. Impact of platelet reactivity on clinical outcomes after percutaneous coronary intervention. A collaborative meta-analysis of individual participant data. *J Am Coll Cardiol* 2011;58:1945-54.
7. Michelson AD. Platelet function testing in cardiovascular diseases. *Circulation* 2004;110:e489-93.
8. Geiger J, Teichmann L, Grossmann R, et al. Monitoring of clopidogrel action: comparison of methods. *Clin Chem* 2005;51:957-65.
9. Price MJ. Bedside evaluation of thienopyridine antiplatelet therapy. *Circulation* 2009;119:2625-32.
10. Breet NJ, van Werkum JW, Bouman HJ, et al. Comparison of platelet function tests in predicting clinical outcome in patients undergoing coronary stent implantation. *JAMA* 2010;303:754-62.
11. Quinn MJ, Byzova TV, Qin J, Topol EJ, Plow EF. Integrin α IIb β 3 and its antagonism. *Arterioscler Thromb Vasc Biol* 2003;23:945-52.
12. Joo SJ. Mechanisms of platelet activation and integrin α II β 3. *Korean Circ J* 2012;42:295-301.
13. Shattil SJ, Hoxie JA, Cunningham M, Brass LF. Changes in the platelet membrane glycoprotein IIb. IIIa complex during platelet activation. *J Biol Chem* 1985;260:11107-14.
14. Shattil SJ, Cunningham M, Hoxie JA. Detection of activated platelets

- in whole blood using activation-dependent monoclonal antibodies and flow cytometry. *Blood* 1987;70:307-15.
15. Abrams CS, Ellison N, Budzynski AZ, Shattil SJ. Direct detection of activated platelets and platelet-derived microparticles in humans. *Blood* 1990;75:128-38.
 16. Harder S, Klinkhardt U, Graff J, et al. In vitro dose response to different GPIIb/IIIa-antagonists: inter-laboratory comparison of various platelet function tests. *Thromb Res* 2001;102:39-48.
 17. Furman MI, Kereiakes DJ, Krueger LA, et al. Quantification of abciximab-induced platelet inhibition is assay dependent: a comparative study in patients undergoing percutaneous coronary intervention. *Am Heart J* 2003;145:e6.
 18. Furman MI, Krueger LA, Linden MD, Barnard MR, Frelinger AL 3rd, Michelson AD. Release of soluble CD40L from platelets is regulated by glycoprotein IIb/IIIa and actin polymerization. *J Am Coll Cardiol* 2004;43:2319-25.
 19. Sibbing D, Byrne RA, Bernlochner I, Kastrati A. High platelet reactivity and clinical outcome - fact and fiction. *Thromb Haemost* 2011;106:191-202.
 20. Price MJ, Berger PB, Teirstein PS, et al. Standard- vs high-dose clopidogrel based on platelet function testing after percutaneous coronary intervention: the GRAVITAS randomized trial. *JAMA* 2011;305:1097-105.
 21. Xiao Z, Thérroux P. Clopidogrel inhibits platelet-leukocyte interactions and thrombin receptor agonist peptide-induced platelet activation in patients with an acute coronary syndrome. *J Am Coll Cardiol* 2004;43:1982-8.
 22. Cayla G, Macia JC, Rabesandratana H, et al. Flow cytometric assessment of vasodilator-stimulated phosphoprotein: prognostic value of recurrent cardiovascular events after acute coronary syndromes. *Arch Cardiovasc Dis* 2008;101:743-51.
 23. Siller-Matula JM, Christ G, Lang IM, Delle-Karth G, Huber K, Jilma B. Multiple electrode aggregometry predicts stent thrombosis better than the vasodilator-stimulated phosphoprotein phosphorylation assay. *J Thromb Haemost* 2010;8:351-9.