

Original Article



Comparison of Supervised Hospital-based versus Educated Home-based Exercise Training in Korean Heart Failure Patients

Minsu Kim , MD, Min-Seok Kim , MD, PhD, Seo-Jin Lim , RN, Jung-Min Ahn , MD, Jae-Joong Kim , MD, PhD, and Seung-Jung Park , MD, PhD

Department of Internal Medicine, Heart Institute, University of Ulsan College of Medicine, Asan Medical Center, Seoul, Korea



Received: Mar 20, 2017
Revised: May 23, 2017
Accepted: Jun 6, 2017

Correspondence to

Min-Seok Kim, MD, PhD

Department of Internal Medicine, Heart Institute, University of Ulsan College of Medicine, Asan Medical Center, 88, Olympic-ro 43-gil, Songpa-gu, Seoul 05505, Korea.
Tel: +82-2-3010-3948
Fax: +82-2-486-5918
E-mail: msk@amc.seoul.kr

Copyright © 2017. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Minsu Kim
<https://orcid.org/0000-0001-9230-3137>
Min-Seok Kim
<https://orcid.org/0000-0002-8512-797X>
Seo-Jin Lim
<https://orcid.org/0000-0001-6369-6517>
Jung-Min Ahn
<https://orcid.org/0000-0003-4031-391X>
Jae-Joong Kim
<https://orcid.org/0000-0002-2714-2282>
Seung-Jung Park
<https://orcid.org/0000-0002-9187-5405>

ABSTRACT

Background and Objectives: Although many clinical trials have shown that exercise training (ET) improves functional capacity and clinical outcomes in heart failure (HF) patients, data comparing supervised hospital-based and educated home-based ET in HF patients is lacking.

Subjects and Methods: This was a single-center, non-randomized, prospective study of 82 HF patients with reduced ejection fraction ($\leq 40\%$) who completed ET. The hospital-based group (n=30) underwent supervised ET at 60% of peak oxygen consumption (VO_2), while a physiotherapist-educated group (n=52) exercised at home without monitoring. The 2 groups were compared before and after the 3-month ET program with respect to functional capacity, quality of life (QOL), and cardiac events (all-cause mortality or hospitalization with worsening HF).

Results: After ET, peak VO_2 increased in the hospital-based group (19.4 ± 4.4 to 21.4 ± 4.3 mL/min/kg, $p=0.006$) and remained unchanged in the home-based group (18.9 ± 4.6 to 18.4 ± 4.6 mL/min/kg, $p=0.660$). The change in peak VO_2 after ET was greater in the hospital-based group compared to the home-based group by 2.5 mL/min/kg ($p=0.014$). QOL improved in the hospital-based group (43.1 ± 18.0 to 28.1 ± 21.6 , $p=0.003$). During one year of follow-up, a comparison of the 2 groups did not reveal a statistical difference in cardiac events (hazard ratio, 0.66; 95% confidence interval, 0.2–2.8; $p=0.570$).

Conclusion: Hospital-based ET was beneficial for HF patients, improving functional capacity and QOL. However, no significant advantages were observed in terms of a composite endpoint compared to home-based ET. Further investigations are required to address the effects and roles of the 2 ET programs for HF patients.

Keywords: Heart failure; Exercise; Rehabilitation

INTRODUCTION

Heart failure (HF) is an increasingly common cardiovascular syndrome and is the final pathway in several cardiac disorders.¹⁾ Many HF patients experience dyspnea, fatigue, diminished exercise capacity, and poor quality of life (QOL).

Conflict of Interest

The authors have no financial conflicts of interest.

Author Contributions

Conceptualization: Kim M, Kim MS, Lim SJ, Ahn JM, Kim JJ, Park SJ; Data curation: Kim M, Lim SJ; Formal analysis: Kim M; Funding acquisition: Kim M, Kim MS; Investigation: Kim M, Kim MS; Methodology: Kim JJ; Project administration: Kim M, Kim MS; Resources: Kim M, Kim MS; Software: Ahn JM; Supervision: Park SJ; Validation: Kim JJ; Visualization: Kim M, Kim MS; Writing - original draft: Kim M; Writing - review & editing: Kim MS.

Progressive functional deterioration is common in HF patients because of exercise intolerance, but can be avoided by discouraging a sedentary lifestyle and promoting physical activity.²⁾ Many recent randomized controlled trials have shown that exercise training (ET) programs enhance aerobic capacity, delay the onset of anaerobic metabolism, improve autonomic balance, decrease HF hospitalization, and increase survival.³⁻⁵⁾ These trials comparing supervised hospital-based ET with educated home-based ET in HF patients lack data, especially among the Korean population. This study aims to compare the effects of both ET programs on the exercise capacity, QOL, and clinical outcomes of HF patients.

SUBJECTS AND METHODS**Patients**

Patients were enrolled between September 2010 and February 2014 from referrals to the Cardiovascular Disease Prevention and Rehabilitation Center at Asan Medical Center, Seoul, Korea, which offers a multidisciplinary, outpatient cardiac rehabilitation program, providing supervised hospital-based ET, dietary and psychological counseling, and nursing education and support.

Symptomatic outpatients classified as New York Heart Association (NYHA) functional class I–III were selected. Included patients had at least 6 months of optimal and standard pharmacological treatment, left ventricular ejection fraction (LVEF) of 40% or less (according to the modified Simpson's method), and stable status for at least 3 months. Exclusion criteria were history of ventricular arrhythmia, implantation of a cardioverter-defibrillator or cardiac resynchronization therapy, pulmonary disease (vital capacity and/or forced expiratory volume in one second <80% of age-dependent predicted value), unstable angina, untreated coronary artery stenosis >50%, decompensated HF, hemodynamically significant valvular heart disease, uncontrolled hypertension (resting systolic blood pressure >150 mmHg or diastolic blood pressure >100 mmHg), anemia (hemoglobin <12 g/dL), clinically relevant musculoskeletal disease, clinically relevant arrhythmia, or change in cardiovascular medication within the previous four weeks. Of the 82 initial patients, 8 did not complete the ET program, 30 did not perform the follow-up cardiopulmonary exercise testing, and 44 completed the protocol (**Figure 1**).

The institution's Ethical Review Board approved this prospective study, and informed consent was obtained from each patient. The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki, as reflected in a prior approval by the institution's human research committee.

Design

A non-randomized prospective trial compared a supervised hospital-based ET group with an educated home-based ET group. The study group was assigned according to the patient's decision. Patients underwent a physical examination, cardiopulmonary exercise testing, a 6-minute walk test, and blood sampling at baseline and after 3 months of ET. The Minnesota Living with Heart Failure Questionnaire (MLWHFQ) was used to assess QOL. Sonographers performed 2-dimensional echocardiographic examinations on all patients where LVEF was measured in the apical 4-chamber view and 2-chamber view using the modified Simpson's method.

Patients performed symptom-limited cardiopulmonary graded exercise testing on a COSMED Quark CPET metabolic cart (COSMED, Rome, Italy) with a Lode cycle ergometer (Lode




Figure 1. Flow of participants through the trial.
ET = exercise training.

Excalibur Sport; Lode B.V., Groningen, the Netherlands) beginning at a 20-watt workload and increasing by 15 to 25-watt increments every 2 minutes. Peak oxygen consumption (VO_2) was measured directly breath-by-breath and defined as the average of 3 measurements recorded in the last minute of exercise. A 12-lead electrocardiogram was continuously recorded, blood pressure was measured every 2 minutes, and cardiopulmonary variables were calculated using printouts of 10-second averaged values. The hospital-based group participated in a structured, group-based exercise program with a goal length of 30–40 minutes, with one or 3 sessions each week for a total of 12 or 36 sessions in 3 months. Training intensity was tailored to 60% to 70% of the target heart rate reserve (maximum heart–resting heart rate).⁶ Exercises for the hospital group included treadmill walking, stationary cycling, arm ergometry, and stair climbing.

Patients in the home-based group participated in one session each month for 3 months with a formal exercise prescription. All patients received a detailed booklet of self-management educational materials at the time of enrollment, including information on medications, fluid management, symptom exacerbation, sodium intake, and moderate-intensity activity of 30 minutes (as tolerated) on most days of the week, consistent with guidelines from the American College of Cardiology and the American Heart Association.⁷

Clinical outcomes

Follow-up started on the day the ET protocol began and ended with an adverse event. All patients were followed up every 2 months for at least one year after enrollment. The primary endpoint was a composite of all-cause mortality or hospitalization with worsening HF, defined as admission with new or increasing symptoms and signs of fluid retention, or objective evidence of HF and initiation of intravenous diuretic agents or vasoactive agents, mechanical ventilation, or mechanical support. All events were carefully verified and adjudicated by independent clinicians.

Statistical analysis

Data are expressed as mean±standard deviation (SD) for continuous variables and as frequency for categorical variables. Differences between groups were assessed using χ^2 analysis for categorical variables and Student's t-test for continuous variables. Changes within the groups during follow-up were assessed by the t-test for paired variables. Survival curves for all-cause mortality or hospitalization with worsening HF were compared using Kaplan-Meier estimates and tested for statistical significance using the log-rank test at $p < 0.050$. Event (or censoring) times for all patients were measured from the time of allocation (time zero). Statistical analysis was performed using commercially available software (SPSS 21 for Windows; SPSS Inc., Chicago, IL, USA).

RESULTS

Baseline characteristics

Clinical characteristics of the 82 study patients according to ET type are presented in **Table 1**. There were no significant differences between the groups in terms of age, body mass index, NYHA functional classification, underlying diseases, medications, LVEF, or Dutch HF knowledge scales. The proportions of male sex and current smokers in the educated home-based group were higher than those in the supervised hospital-based group.

Table 1. Baseline characteristics of patients in the 2 treatment groups

Variables	Hospital-based group (n=30)	Home-based group (n=52)	p value
Clinical characteristics			
Age (years)	56.5±13.8	52.9±12.7	0.230
Male sex	18 (60.0)	43 (82.7)	0.020
Height (cm)	164.6±9.7	167.5±9.0	0.180
Weight (kg)	66.7±15.7	71.1±10.7	0.180
BMI (kg/m ²)	24.5±3.9	24.6±2.9	0.960
NYHA functional class			0.250
I	5 (16.7)	13 (25.0)	
II	21 (70.0)	35 (67.3)	
III	4 (13.3)	4 (7.7)	
Diabetes	8 (26.7)	13 (25.0)	0.890
Hypertension	11 (36.7)	20 (38.5)	0.870
Atrial fibrillation	9 (30.0)	12 (23.1)	0.490
Current smoker	3 (10.0)	16 (33.3)	0.020
Dilated CMP	9 (30.0)	19 (36.5)	0.550
Ischemic CMP	6 (20.0)	19 (36.5)	0.120
Hypertensive HF	4 (13.3)	2 (3.8)	0.190
LVEF	30.4±10.4	28.9±7.3	0.450
Beta blocker	26 (86.7)	46 (88.5)	0.810
Calcium channel blocker	4 (13.3)	10 (19.2)	0.490
ARB/ACEi	20 (66.7)	31 (59.6)	0.530
Spironolactone	12 (40.0)	18 (34.6)	0.630
Diuretics	24 (80.0)	39 (75.0)	0.610
Antiplatelet agent	10 (33.3)	16 (30.8)	0.810
Anticoagulation	10 (33.3)	18 (34.6)	0.910
Dutch HF knowledge scale	12.5±1.5	12.0±1.4	0.190

Values are presented as mean±SD or number (%).

ACEi = angiotensin-converting enzyme inhibitor; ARB = angiotensin receptor blocker; BMI = body mass index; CMP = cardiomyopathy; HF = heart failure; LVEF = left ventricular ejection fraction; NYHA = New York Heart Association; SD = standard deviation.

Exercise training effects

The changes from baseline in exercise time, peak VO₂, 6-minute walk, and MLWHFQ scale at 3 months are presented in **Tables 2** and **3**. Significant improvements of exercise time and peak VO₂ were observed in the supervised hospital-based ET group between the pre- and post-exercise periods. There were no such differences for the educated home-based ET group. The post-exercise differences between the 2 groups in exercise time and peak VO₂ were statistically significant ($p=0.010$ for exercise time, $p=0.014$ for peak VO₂). There was no significant difference between the pre- and post-exercise periods in the 6-minute walk for either group. Self-reported total MLWHFQ score improved significantly in the hospital-based group, but the difference between the groups was not statistically significant. There were no serious adverse events leading to early termination of ET such as ventricular arrhythmia, syncope, unstable angina, or worsening HF in any of the groups during the program.

Clinical outcomes

There were eight cardiac events during the one-year follow-up period, including 1 all-cause mortality in the supervised hospital-based ET group. Two patients in the hospital-based group and 5 patients in the home-based group were hospitalized with worsening HF (**Table 4**). The composite endpoint of all-cause mortality or hospitalization with worsening HF was not statistically different in the 2 groups (**Figure 2**; hazard ratio [HR], 0.66; 95% confidence

Table 2. Comparison of the cardiopulmonary exercise test results and QOL between the pre- and post-exercise periods

Variables	Hospital-based group (n=24)	Home-based group (n=20)
Exercise time (seconds)		
Base	599.0±130.1	682.5±241.5
F/u	650.3±153.6	668.5±239.7
Change	51.3 (15.6 to 87.0)	-14.0 (-52.8 to 24.9)
p value	0.007	0.460
Peak VO₂ (mL/min/kg)		
Base	19.4±4.4	18.9±4.6
F/u	21.4±4.3	18.4±4.6
Change	2.0 (0.6 to 3.4)	-0.4 (-1.8 to 1.0)
p value	0.006	0.660
6-minute walk (m)		
Base	561.9±49.4	560.4±90.0
F/u	583.9±72.3	562.4±68.9
Change	21.9 (-12.7 to 56.6)	2.0 (-24.5 to 28.5)
p value	0.200	0.870
MLWHFQ scale		
Base	43.1±18.0	49.9±26.5
F/u	28.1±21.6	37.8±29.3
Change	-14.6 (-23.6 to -5.5)	-11.8 (-25.5 to 1.9)
p value	0.003	0.087

Data are mean±SD or mean (95% CI). Complete case analysis. Expected 81 patients at 3 months. CI = confidence interval; F/u = follow-up; MLWHFQ = Minnesota Living with Heart Failure Questionnaire; QOL = quality of life; SD = standard deviation; VO₂ = oxygen consumption.

Table 3. Comparison of the cardiopulmonary exercise test results and QOL between the 2 groups


Variables	Hospital-based group (n=24)	Home-based group (n=20)	Difference between groups	p value
Change in				
Exercise time (seconds)	51.3 (15.6 to 87.0)	-14.0 (-52.8 to 24.9)	65.2 (14.0 to 116.5)	0.010
Peak VO ₂ (mL/min/kg)	2.0 (0.6 to 3.4)	-0.4 (-1.8 to 1.0)	2.5 (0.5 to 4.4)	0.014
6-minute walk (m)	21.9 (-12.7 to 56.6)	2.0 (-24.5 to 28.5)	19.8 (-21.2 to 61.1)	0.330
MLWHFQ scale	-14.6 (-23.6 to -5.5)	-11.8 (-25.5 to 1.9)	-2.8 (-17.9 to 12.4)	0.720

Data are mean (95% CI). Complete case analysis. Expected 81 patients at 3 months. CI = confidence interval; MLWHFQ = Minnesota Living with Heart Failure Questionnaire; QOL = quality of life; VO₂ = oxygen consumption.

Table 4. Clinical outcomes in the 2 groups

Clinical outcome	Hospital-based group (n=30)	Home-based group (n=52)	HR* (95% CI)	p value
All-cause mortality or hospitalization with worsening HF	3	5	0.66 (0.20–2.80)	0.570
All-cause mortality	1	0	59.80 (infinite)	0.620
Hospitalization with worsening HF	2	5	0.46 (0.09–2.40)	0.360


CI = confidence interval; HF = heart failure; HR = hazard ratio.
*HR of the hospital-based group compared to the home-based group.



No. at risk					
Hospital-based	30	29	29	29	20
Home-based	52	50	19	19	17

Figure 2. Kaplan-Meier curves for time to composite endpoint (all-cause mortality or hospitalization with worsening HF). The composite endpoint was not statistically different in either group at 1 year after the start of the ET program.

ET = exercise training; HF = heart failure.



No. at risk					
36 sessions	15	15	15	15	11
12 sessions	15	14	14	14	9
3 sessions	19	19	12	12	10
1 session	33	31	8	8	7

Figure 3. Kaplan-Meier curves for time to composite endpoint (all-cause mortality or hospitalization with worsening HF) by the number of exercise sessions. There was no significant difference between the groups by the number of exercise sessions.

HF = heart failure.

interval [CI], 0.2–2.8; p=0.570). We analyzed the post-exercise clinical outcome according to the number of exercise sessions. However, event-free survival did not correlate with number of exercise sessions (**Figure 3**).

DISCUSSION

Significant improvement in functional capacity, exercise capacity, and QOL was observed in the supervised hospital-based group following the 12-week cardiac rehabilitation program. The composite endpoint of all-cause mortality or hospitalization with worsening HF was not different between the groups for one year after cessation of the ET program.

HF is a common clinical syndrome caused by a variety of cardiac diseases. After an initial reduction in cardiac output, intrinsic cardiac decline leads to systemic deterioration with a reduced ability to exercise, early fatigue, weakness, and shortness of breath.⁸⁾ Such symptoms impact the performance of activities of daily living and significantly contribute to reduced participation and poor QOL.⁹⁾

ET is a valuable adjunct in the therapeutic approach to stabilize HF patients and is a class I recommendation from the American College of Cardiology and the American Heart Association.¹⁰⁾ Most studies of exercise in HF patients have focused on hospital-based programs where patients exercise under controlled conditions. Many Korean HF patients are unable to access a formal ET program because they live in remote geographic locations, have difficulty with transportation, or do not have the financial resources to pay for a structured ET program. Furthermore, HF patients with functional limitations demonstrated poor cognitive function and self-care confidence, which can interrupt active participation in a hospital-based ET program.¹¹⁾ Since a home-based program is optimal for these patients, the comparison between hospital- and home-based ET programs has great clinical implications.

Peak VO_2 improved in the hospital-based group by 2.5 mL/min/kg compared to the home-based group (95% CI, 0.5–4.4; $p=0.014$) at cessation of the ET program, along with an improved exercise time of 65.2 seconds (95% CI, 14.0–116.5; $p=0.010$). This is an encouraging result, as the improvement and sustainability of functional performance are the primary goals of an ET program. Exercise capacity increase was modest in the hospital-based ET group, with an improvement rate in measured peak VO_2 of about 10%, relatively lower than in other studies.¹²⁾¹³⁾ There was no improvement in the home-based ET group. Considering the numerical improvement of QOL in both groups, modest or no improvement in exercise capacity might result from the poor clinical conditions or comorbidities of patients rather than inadequate exercise.

Among the exercise parameters, the 6-minute walk was not significantly improved in either group (21.9 vs. 2.0 m, $p=0.330$). The 6-minute walk offers a convenient and economical clinical test of functional performance compared with normal activity,¹⁴⁾ but has several limitations. The test provides no insight into the mechanisms of exercise limitation, and its results can be affected by unrelated factors such as age, sex, height, and weight. The benefits of ET should therefore not be underestimated, irrespective of the program type.

A number of studies have found an increase in QOL with ET.¹⁵⁾¹⁶⁾ Some studies have demonstrated improved QOL with little or no change in exercise capacity.¹⁷⁾¹⁸⁾ This study found a significant improvement of QOL in the supervised hospital-based ET group and numeric improvement of QOL in the educated home-based ET group. These findings suggest that QOL improvement can occur even with modest or no change in exercise capacity. It is further suggested that QOL is determined not only by physiologic change, but also by factors affecting the patient's perception of health.

ET improves cardiac output and regional blood flow, which are associated with increased peak VO_2 . HF patients with a higher peak VO_2 have a greater survival rate. Therefore, improving peak VO_2 can improve survival and reduce hospitalization. Alterations in the autonomic nervous system that decrease sympathetic tone and increase vagal tone might have similar effects.¹⁹⁾ Hospital-based ET did not provide a significant reduction in the composite endpoint of all-cause mortality or hospitalization with worsening HF, which was not in concordance with other studies.³⁾⁵⁾ The home-based ET group in this study differed from groups in other studies, as patients were educated with a formal exercise prescription one or 3 times for effective ET at home. These patients might have increased exercise intensity or duration because of the unblinded assignment, possibly affecting the results between the program types in clinical outcomes over 1 year.

This study has several limitations. It was a non-randomized trial where patients who agreed to and were available for participation could be systematically different from those unable to participate, which could limit the generalizability of the findings. The 82-patient population was relatively small and was derived from a single center, and patients were young compared to the general population with HF. Thirty-eight (20% of the hospital-based group; 62% of the home-based group) of the total study patients terminated the ET program prematurely or did not perform the follow-up functional test because of health insurance issues. However, clinical follow-up was performed every 2 months in the HF outpatient clinic, assessing the occurrence of clinical events in all patients. Clinical outcomes of the non-follow-up patients were not different compared to those of the follow-up patients (**Supplementary Figure 1**). Finally, adherence to the ET was not assessed because it was difficult to objectively evaluate whether or not practicing ET at home made a difference. This is a major limitation of our study and could be a bias that affected the results.

In conclusion, the present study demonstrated that supervised hospital-based ET was beneficial for HF patients with improving functional capacity and QOL. However, hospital-based ET provided a non-significant reduction in the risk for the composite endpoint of all-cause mortality or hospitalization of worsening HF compared to the home-based ET program. In cases that were unsuitable for supervised hospital-based ET, educated home-based ET with a detailed exercise prescription could be considered as an alternative option for the non-pharmacological treatment for HF. Further well-designed large studies are needed to define the role and effect of hospital-based or home-based ET programs.

ACKNOWLEDGMENTS

Thanks to Jung Bok Lee, a biostatistician in our institute, for the statistical consultation.

SUPPLEMENTARY MATERIAL

Supplementary Figure 1

Kaplan-Meier curves for time to composite endpoint (all-cause mortality or hospitalization with worsening HF). The composite endpoint in patients with or without the follow-up functional test was not statistically different at one year after the start of the ET program.

[Click here to view](#)

REFERENCES

1. Lloyd-Jones D, Adams RJ, Brown TM, et al. Executive summary: heart disease and stroke statistics--2010 update: a report from the American Heart Association. *Circulation* 2010;121:948-54.
[PUBMED](#) | [CROSSREF](#)
2. Mancini DM, Walter G, Reichek N, et al. Contribution of skeletal muscle atrophy to exercise intolerance and altered muscle metabolism in heart failure. *Circulation* 1992;85:1364-73.
[PUBMED](#) | [CROSSREF](#)
3. Belardinelli R, Georgiou D, Cianci G, Purcaro A. Randomized, controlled trial of long-term moderate exercise training in chronic heart failure: effects on functional capacity, quality of life, and clinical outcome. *Circulation* 1999;99:1173-82.
[PUBMED](#) | [CROSSREF](#)
4. Zwisler AD, Schou L, Soja AM, et al. A randomized clinical trial of hospital-based, comprehensive cardiac rehabilitation versus usual care for patients with congestive heart failure, ischemic heart disease, or high risk of ischemic heart disease (the DANREHAB trial)--design, intervention, and population. *Am Heart J* 2005;150:899.
[PUBMED](#) | [CROSSREF](#)
5. O'Connor CM, Whellan DJ, Lee KL, et al. Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA* 2009;301:1439-50.
[PUBMED](#) | [CROSSREF](#)
6. Mezzani A, Hamm LF, Jones AM, et al. Aerobic exercise intensity assessment and prescription in cardiac rehabilitation: a joint position statement of the European Association for Cardiovascular Prevention and Rehabilitation, the American Association of Cardiovascular and Pulmonary Rehabilitation and the Canadian Association of Cardiac Rehabilitation. *Eur J Prev Cardiol* 2013;20:442-67.
[PUBMED](#) | [CROSSREF](#)
7. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013;62:e147-239.
[PUBMED](#) | [CROSSREF](#)
8. Jessup M, Brozena S. Heart failure. *N Engl J Med* 2003;348:2007-18.
[PUBMED](#) | [CROSSREF](#)
9. Working Group on Cardiac Rehabilitation & Exercise Physiology and Working Group on Heart Failure of the European Society of Cardiology. Recommendations for exercise training in chronic heart failure patients. *Eur Heart J* 2001;22:125-35.
[PUBMED](#) | [CROSSREF](#)
10. Hunt SA American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *J Am Coll Cardiol* 2005;46:e1-82.
[PUBMED](#) | [CROSSREF](#)
11. Kim JS, Hwang SY, Shim JL, Jeong MH. Cognitive function and self-care in patients with chronic heart failure. *Korean Circ J* 2015;45:310-6.
[PUBMED](#) | [CROSSREF](#)
12. Giannuzzi P, Temporelli PL, Corrà U, Tavazzi LELVD-CHF Study Group. Antiremodeling effect of long-term exercise training in patients with stable chronic heart failure: results of the Exercise in Left Ventricular Dysfunction and Chronic Heart Failure (ELVD-CHF) Trial. *Circulation* 2003;108:554-9.
[PUBMED](#) | [CROSSREF](#)
13. Demopoulos L, Bijou R, Fergus I, Jones M, Strom J, LeJemtel TH. Exercise training in patients with severe congestive heart failure: enhancing peak aerobic capacity while minimizing the increase in ventricular wall stress. *J Am Coll Cardiol* 1997;29:597-603.
[PUBMED](#) | [CROSSREF](#)
14. Lucas C, Stevenson LW, Johnson W, et al. The 6-min walk and peak oxygen consumption in advanced heart failure: aerobic capacity and survival. *Am Heart J* 1999;138:618-24.
[PUBMED](#) | [CROSSREF](#)
15. Parnell MM, Holst DP, Kaye DM. Exercise training increases arterial compliance in patients with congestive heart failure. *Clin Sci (Lond)* 2002;102:1-7.
[PUBMED](#) | [CROSSREF](#)

16. Flynn KE, Piña IL, Whellan DJ, et al. Effects of exercise training on health status in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA* 2009;301:1451-9.
[PUBMED](#) | [CROSSREF](#)
17. Oka RK, De Marco T, Haskell WL, et al. Impact of a home-based walking and resistance training program on quality of life in patients with heart failure. *Am J Cardiol* 2000;85:365-9.
[PUBMED](#) | [CROSSREF](#)
18. Wielenga RP, Erdman RA, Huisveld IA, et al. Effect of exercise training on quality of life in patients with chronic heart failure. *J Psychosom Res* 1998;45:459-64.
[PUBMED](#) | [CROSSREF](#)
19. Coats AJ, Adamopoulos S, Radaelli A, et al. Controlled trial of physical training in chronic heart failure. Exercise performance, hemodynamics, ventilation, and autonomic function. *Circulation* 1992;85:2119-31.
[PUBMED](#) | [CROSSREF](#)