

Original Research



OPEN ACCESS

Received: Sep 24, 2021

Revised: Nov 23, 2021

Accepted: Nov 30, 2021

Published online: Dec 28, 2021

Correspondence to

Jung-Sun Kim, MD, PhD

Division of Cardiology, Severance Cardiovascular Hospital, Yonsei University College of Medicine, 50-1, Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea.
Email: kjs1218@yuhs.ac

Yun-Hyeong Cho, MD, PhD

Division of Cardiology, Department of Internal Medicine, Myongji Hospital, Hanyang University College of Medicine, 55, Hwasu-ro 14 beon-gil, Deogyang-gu, Goyang 10475, Korea.
Email: princette@gmail.com

*Yong-Joon Lee and Yongsung Suh contributed equally to this work.

Copyright © 2022. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Yong-Joon Lee

<https://orcid.org/0000-0003-4526-6120>

Yongsung Suh

<https://orcid.org/0000-0002-1975-123X>

Jung-Sun Kim

<https://orcid.org/0000-0003-2263-3274>

Ticagrelor Monotherapy After 3-Month Dual Antiplatelet Therapy in Acute Coronary Syndrome by High Bleeding Risk: The Subanalysis From the TICO Trial

Yong-Joon Lee , MD^{1,*}, Yongsung Suh , MD^{2,*}, Jung-Sun Kim , MD, PhD¹, Yun-Hyeong Cho , MD, PhD², Kyeong Ho Yun , MD³, Yong Hoon Kim , MD⁴, Jae Young Cho , MD³, Ae-Young Her , MD⁴, Sungsoo Cho , MD⁵, Dong Woon Jeon , MD⁶, Sang-Yong Yoo , MD⁷, Deok-Kyu Cho , MD⁸, Bum-Kee Hong , MD⁹, Hyuckmoon Kwon , MD⁹, Sung-Jin Hong , MD¹, Chul-Min Ahn , MD¹, Dong-Ho Shin , MD¹, Chung-Mo Nam , PhD¹⁰, Byeong-Keuk Kim , MD¹, Young-Guk Ko , MD¹, Donghoon Choi , MD¹, Myeong-Ki Hong , MD¹, Yangsoo Jang , MD¹¹, and for the TICO investigators

¹Division of Cardiology, Severance Cardiovascular Hospital, Yonsei University College of Medicine, Seoul, Korea

²Division of Cardiology, Department of Internal Medicine, Myongji Hospital, Hanyang University College of Medicine, Goyang, Korea

³Division of Cardiology, Wonkwang University Hospital, Iksan, Korea

⁴Division of Cardiology, Kangwon National University School of Medicine, Chuncheon, Korea

⁵Division of Cardiology, Dankook University Hospital, Dankook University College of Medicine, Cheonan, Korea

⁶Division of Cardiology, National Health Insurance Service Ilsan Hospital, Goyang, Korea

⁷Division of Cardiology, Gangneung Asan Hospital, University of Ulsan College of Medicine, Gangneung, Korea

⁸Division of Cardiology, Yongin Severance Hospital, Yonsei University College of Medicine, Yongin, Korea

⁹Division of Cardiology, Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul, Korea

¹⁰Department of Preventive Medicine and Biostatistics, Yonsei University College of Medicine, Seoul, Korea

¹¹Division of Cardiology, CHA Bundang Medical Center, CHA University College of Medicine, Seongnam, Korea

AUTHOR'S SUMMARY

This study demonstrated that for acute coronary syndrome patients treated with drug eluting stents, high bleeding risk (HBR) was associated with higher rates of net adverse clinical events (NACEs), not only including bleeding but also ischemic events. Ticagrelor monotherapy after 3-month dual antiplatelet therapy (DAPT) was associated with lower rates of NACEs and major bleeding than ticagrelor-based 12-month DAPT regardless of HBR, with no significant interaction between therapy and HBR. These findings were consistent regardless of HBR definition by Academic Research Consortium for HBR criteria or by Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent DAPT score.

ABSTRACT

Background and Objectives: Identifying patients with high bleeding risk (HBR) is important when making decisions for antiplatelet therapy strategy. This study evaluated the impact of

Yun-Hyeong Cho 
<https://orcid.org/0000-0001-7581-9545>
 Kyeong Ho Yun 
<https://orcid.org/0000-0003-4911-8854>
 Yong Hoon Kim 
<https://orcid.org/0000-0002-9669-3598>
 Jae Young Cho 
<https://orcid.org/0000-0001-7972-6223>
 Ae-Young Her 
<https://orcid.org/0000-0002-9990-6843>
 Sungsoo Cho 
<https://orcid.org/0000-0003-2059-1584>
 Dong Woon Jeon 
<https://orcid.org/0000-0001-7409-4378>
 Sang-Yong Yoo 
<https://orcid.org/0000-0002-9054-4698>
 Deok-Kyu Cho 
<https://orcid.org/0000-0001-6130-3994>
 Bum-Kee Hong 
<https://orcid.org/0000-0002-6456-0184>
 Hyuckmoon Kwon 
<https://orcid.org/0000-0001-9901-5015>
 Sung-Jin Hong 
<https://orcid.org/0000-0003-4893-039X>
 Chul-Min Ahn 
<https://orcid.org/0000-0002-7071-4370>
 Dong-Ho Shin 
<https://orcid.org/0000-0002-7874-5542>
 Chung-Mo Nam 
<https://orcid.org/0000-0003-0985-0928>
 Byeong-Keuk Kim 
<https://orcid.org/0000-0003-2493-066X>
 Young-Guk Ko 
<https://orcid.org/0000-0001-7748-5788>
 Donghoon Choi 
<https://orcid.org/0000-0002-2009-9760>
 Myeong-Ki Hong 
<https://orcid.org/0000-0002-2090-2031>
 Yangsoo Jang 
<https://orcid.org/0000-0002-2169-3112>

Trial Registration

ClinicalTrials.gov Identifier: NCT02494895

Funding

This work was supported by the Cardiovascular Research Center (Seoul, South Korea) and funded by Biotronik (Bülach, Switzerland). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Conflict of Interest

The authors have no financial conflicts of interest.

ticagrelor monotherapy after 3-month dual antiplatelet therapy (DAPT) according to HBR in acute coronary syndrome (ACS) patients treated with drug eluting stents (DESs).

Methods: In this post-hoc analysis of the TICO trial, HBR was defined by 2 approaches: meeting Academic Research Consortium for HBR (ARC-HBR) criteria or Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent DAPT (PRECISE-DAPT) score ≥ 25 . The primary outcome was a 3–12 months net adverse clinical event (composite of major bleeding and adverse cardiac and cerebrovascular events).

Results: Of the 2,980 patients without adverse events during the first 3 months after DES implantation, 453 (15.2%) were HBR by ARC-HBR criteria and 504 (16.9%) were HBR by PRECISE-DAPT score. The primary outcome rate was higher in HBR versus non-HBR patients (by ARC-HBR criteria: hazard ratio [HR], 2.87; 95% confidence interval [CI], 1.76–4.69; $p < 0.001$; by PRECISE-DAPT score: HR, 3.09; 95% CI, 1.92–4.98; $p < 0.001$). Ticagrelor monotherapy after 3-month DAPT was associated with lower primary outcome rate than ticagrelor-based 12-month DAPT regardless of HBR by ARC-HBR criteria, with similar magnitudes of therapy effect for HBR and non-HBR patients (p -interaction=0.400). Results were consistent by PRECISE-DAPT score (p -interaction=0.178).

Conclusions: In ACS patients treated with DESs, ticagrelor monotherapy after 3-month DAPT was associated with lower rate of adverse clinical outcomes regardless of HBR, with similar magnitudes of therapy effect between HBR and non-HBR.

Trial Registration: ClinicalTrials.gov Identifier: [NCT02494895](https://clinicaltrials.gov/ct2/show/study/NCT02494895)

Keywords: Acute coronary syndrome; Ticagrelor; Hemorrhage

INTRODUCTION

In patients undergoing percutaneous coronary intervention (PCI) for acute coronary syndrome (ACS), identifying and managing patients with high bleeding risk (HBR) are of major concern since prolonged 12-month dual antiplatelet therapy (DAPT) with aspirin and P2Y₁₂ receptor inhibitor is recommended.¹⁻³ Recently, 2 definitions for HBR, the Academic Research Consortium for HBR (ARC-HBR) criteria and the Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent DAPT (PRECISE-DAPT) score, have been proposed to provide consistency in defining HBR and aid clinical decision-making for patients undergoing PCI.^{3,4} Studies have shown that HBR by these 2 definitions is associated with increased risk of both bleeding and ischemic events.⁵⁻⁷ Furthermore, newer antiplatelet agents, such as ticagrelor or prasugrel, with better speed, potency, and consistent effect have become available, and ticagrelor monotherapy after short-term DAPT may minimize bleeding risk without increasing ischemic risk.⁸⁻¹⁰ The Ticagrelor Monotherapy After 3 Months in the Patients Treated With New Generation Sirolimus-eluting Stent for Acute Coronary Syndrome (TICO) trial demonstrated that ticagrelor monotherapy after 3-month DAPT was associated with a lower incidence of a composite outcome of major bleeding and adverse cardiac and cerebrovascular events at 1 year than currently recommended ticagrelor-based 12-month DAPT in ACS patients treated with drug eluting stents (DESs).¹¹

Thus, the aims of the present post-hoc analysis of the TICO trial were: 1) to apply HBR by ARC-HBR criteria and PRECISE-DAPT score for clinical outcomes; and 2) to investigate the impact of ticagrelor monotherapy after 3-month DAPT in HBR compared to non-HBR patients.

Data Sharing Statement

The data generated in this study is available from the corresponding author(s) upon reasonable request.

Author Contributions

Conceptualization: Lee YJ, Suh Y, Kim JS, Cho YH, Hong SJ, Ahn CM, Shin DH, Nam CM, Kim BK, Ko YG, Choi D, Hong MK, Jang Y; Data curation: Kim JS, Cho YH, Yun KH, Kim YH, Cho JY, Her AY, Cho S, Jeon DW, Yoo SY, Cho DK, Hong BK, Kwon H, Hong SJ, Shin DH, Nam CM, Kim BK, Ko YG, Choi D, Hong MK, Jang Y; Formal analysis: Lee YJ, Suh Y, Kim JS, Nam CM; Funding acquisition: Jang Y; Investigation: Kim JS, Kim YH, Cho JY, Her AY, Cho S, Jeon DW, Yoo SY, Cho DK, Hong BK, Kwon H, Hong SJ, Ahn CM, Shin DH, Nam CM, Kim BK, Ko YG, Choi D, Hong MK, Jang Y; Methodology: Lee YJ, Suh Y, Kim JS, Cho YH, Yun KH, Hong SJ, Ahn CM, Shin DH, Nam CM, Kim BK, Ko YG, Choi D, Hong MK, Jang Y; Resources: Jang Y; Supervision: Kim JS, Jang Y; Writing - original draft: Lee YJ, Suh Y; Writing - review & editing: Lee YJ, Suh Y, Kim JS, Cho YH, Yun KH, Kim YH, Cho JY, Her AY, Cho S, Jeon DW, Yoo SY, Cho DK, Hong BK, Kwon H, Hong SJ, Ahn CM, Shin DH, Nam CM, Kim BK, Ko YG, Choi D, Hong MK, Jang Y.

METHODS

Ethical statement

The trial was approved by the Institutional Review Board of Yonsei University College of Medicine (approval No. 1-2014-0066) and at each participating center and followed the ethical principles of the Declaration of Helsinki 2013. All patients gave written informed consent prior to participation in the trial.

Study subjects and design

The study design and rationale for the TICO trial have been described in detail previously.¹¹⁾¹²⁾ Briefly, the multicenter, prospective, open-label, randomized superiority trial evaluated ticagrelor monotherapy followed by 3-month DAPT versus ticagrelor-based 12-month DAPT in 3,056 patients treated with DESs for ACS (unstable angina, non-ST-elevation myocardial infarction [MI], or ST-elevation MI) in South Korea. Consenting patients were randomized in a 1:1 ratio to either therapy and observed for up to 12 months.

In this post-hoc analysis, patients were classified into ticagrelor monotherapy after 3-month DAPT and ticagrelor-based 12-month DAPT on an intention-to-treat basis for primary analysis, and those who encountered adverse events during the first 3 months after PCI were excluded (n=76). Consequently, a total of 2980 TICO patients were included in this study. Of those, 1,489 received ticagrelor monotherapy after 3-month DAPT, and 1,491 received ticagrelor-based 12-month DAPT. A per-protocol analysis was also performed to evaluate the robustness of the results obtained by intention-to-treat analysis.

Study proceedings

Study patients were classified by HBR using 2 different approaches: 1) by meeting ARC-HBR criteria or 2) by PRECISE-DAPT score for HBR.³⁾⁴⁾ Patients were considered to be at HBR if at least 1 major or 2 minor ARC-HBR criteria were met.³⁾ Major criteria available for analysis were severe or end-stage chronic kidney disease (estimated glomerular filtration rate <30 mL/min/1.73 m² or dialysis), hemoglobin <11 g/dL, previous spontaneous bleeding requiring hospitalization or transfusion within the past 6 months, and active malignancy (excluding nonmelanoma skin cancer) within the past 12 months. Minor criteria available were age ≥75 years, moderate chronic kidney disease (estimated glomerular filtration rate ≥30 and <60 mL/min/1.73 m²), hemoglobin ≥11 and <13 g/dL for male and ≥11 and <12 g/dL for female, use of non-steroidal anti-inflammatory drugs or steroids, and any ischemic stroke beyond the past 12 months. PRECISE-DAPT scores were assessed using an online calculator (<http://www.precisedaptscore.com>) with 5 variables (age, creatine clearance, hemoglobin, white blood cell count, and previous spontaneous bleeding).⁴⁾ Scores were categorized into 4 groups of bleeding risks (very low risk: score ≤10; low risk: score 11–17; moderate risk: score 18–24; and high risk: score ≥25).

Study endpoints

The primary outcome was a net adverse clinical event (NACE), defined as a composite of major bleeding and major adverse cardiac and cerebrovascular events (MACCEs) from 3–12 months after index PCI. MACCE included all-cause death, MI, stent thrombosis, stroke, and target-vessel revascularization. Secondary outcomes included major bleeding and MACCEs.¹²⁾

Major bleeding was defined in accordance with the thrombolysis in MI criteria: intracranial bleeding, hemorrhage associated with a ≥5 g/dL decrease in hemoglobin, or fatal bleeding

that resulted in death within 7 days.¹³ MI after discharge from the hospital was defined as clinical symptoms, electrocardiography changes, or abnormal imaging findings, combined with a creatine kinase MB fraction above the upper normal limits or a troponin T or troponin I level >99th percentile of the upper normal limit.¹⁴ Stent thrombosis was defined as definite or probable stent thrombosis according to the Academic Research Consortium.¹⁵ Stroke was defined as an acute cerebrovascular event that resulted in death or a neurological deficit lasting >24 hours or as an acute infarction demonstrated by imaging studies.¹⁶ Target-vessel revascularization was defined as a repeat PCI or bypass surgery of the target-vessel with either: 1) ischemia symptoms or a positive stress test and angiographic diameter stenosis >50%; or 2) angiographic diameter stenosis >70% without ischemia symptoms or a positive stress test.^{12,17} Routine follow-up of angiography was not recommended in the trial. Adverse events, including bleeding and ischemic events, were categorized by an independent clinical event committee blinded to the treatment assignments and primary results of the trial.¹²

Statistical analysis

Continuous variables were reported as mean \pm SD and compared using Student's t-tests or Mann-Whitney tests. Categorical variables were reported as number (percentage) and compared using χ^2 tests or Fisher's exact tests.

Event rates were estimated using Kaplan-Meier survival analysis and compared using log-rank tests. Hazard ratios (HRs) with 95% confidence intervals (CIs) were computed using Cox regression analysis. Cox regression analysis with tests for interaction were used to assess for differential therapy effects by HBR. All tests were 2-sided, and $p < 0.05$ was considered statistically significant. Statistical analyses were performed using IBM SPSS, version 25.0 (IBM Corporation, Chicago, IL, USA) and R 3.5.3 software (R foundations for Statistical Computing, Vienna, Austria).

RESULTS

Baseline characteristics

Of 3,056 patients enrolled in the TICO trial, there were 2,980 patients who did not encounter adverse events during the first 3 months after index PCI; 453 (15.2%) were considered to be HBR by ARC-HBR criteria, and 504 (16.9%) were considered to be HBR by PRECISE-DAPT score (**Figure 1**). Baseline characteristics by HBR are presented in **Table 1**. By ARC-HBR criteria, HBR patients were older and showed a higher frequency of hypertension, diabetes mellitus, chronic kidney disease, prior MI, prior PCI, prior coronary bypass graft, prior stroke, transfemoral approach, and multivessel coronary artery disease. On the other hand, HBR patients had lower body mass index and a lower frequency of males, current smokers, and primary PCI as compared to non-HBR patients. By PRECISE-DAPT score, HBR patients showed similar characteristics as those with HBR by ARC-HBR criteria, although HBR patients by PRECISE-DAPT score showed a higher frequency of admission via emergency room, clinical presentation of MI, a longer stent length per patient, as well as no difference in prior MI, prior coronary bypass graft, and primary PCI compared to non-HBR patients.

Baseline characteristics within HBR versus non-HBR subgroups for patients treated with ticagrelor monotherapy after 3-month DAPT versus ticagrelor-based 12-month DAPT are presented in **Supplementary Tables 1 and 2**. Based on ARC-HBR criteria, baseline characteristics did not differ between the 2 antiplatelet therapy groups in HBR and non-HBR




Figure 1. Distribution of TICO patients by HBR. Study patients were classified by HBR in 2 different approaches: 1) by meeting ARC-HBR criteria (at least 1 major or 2 minor criteria were met) or 2) by PRECISE-DAPT score for HBR (score ≥ 25).

ARC-HBR = Academic Research Consortium for High Bleeding Risk; HBR = high bleeding risk; PRECISE-DAPT = Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent Dual Antiplatelet Therapy.

patients. Based on PRECISE-DAPT score, baseline characteristics did not differ between the 2 therapy groups in HBR and non-HBR patients, except for prior MI, which was more frequent in non-HBR patients who received ticagrelor monotherapy after 3-month DAPT. Baseline characteristics of patients by per-protocol analysis are presented in **Supplementary Tables 3-5**.

Clinical outcomes by HBR

Clinical outcomes by HBR are presented in **Figure 2** and **Table 2**. The primary outcome rate was significantly higher in HBR versus non-HBR patients regardless of HBR definition. Based on ARC-HBR criteria, the rate of NACE was significantly higher in HBR versus non-HBR patients (5.4% vs. 1.9%; HR, 2.87; 95% CI, 1.76–4.69; $p < 0.001$). Furthermore, in addition to a significantly higher rate of major bleeding in HBR versus non-HBR patients (2.7% vs. 0.6%; HR, 4.91; 95% CI, 2.27–10.61; $p < 0.001$), the rate of MACCE was also significantly higher (3.2% vs. 1.4%; HR, 2.34; 95% CI, 1.26–4.36, $p = 0.006$). Based on PRECISE-DAPT score, the rate of NACE was significantly higher in HBR versus non-HBR patients (5.5% vs. 1.9%; HR, 3.09; 95% CI, 1.92–4.98; $p < 0.001$). Furthermore, in addition to a significantly higher rate of major bleeding in HBR versus non-HBR patients (2.9% vs. 0.5%; HR, 5.96; 95% CI, 2.76–12.88; $p < 0.001$), the rate of MACCE was also significantly higher (3.1% vs. 1.4%; HR, 2.31; 95% CI, 1.25–4.25; $p = 0.006$). These findings were also confirmed in per-protocol analysis (**Supplementary Figure 1**).

Clinical outcomes by HBR and antiplatelet therapy strategy

The therapy effect of ticagrelor monotherapy after 3-month DAPT versus ticagrelor-based 12-month DAPT in HBR and non-HBR patients is presented in **Figure 3** and **Table 3**. With

Table 1. Baseline characteristics by HBR

	ARC-HBR			PRECISE-DAPT		
	HBR (n=453)	Non-HBR (n=2,527)	p value	HBR (n=504)	Non-HBR (n=2,476)	p value
Age (years)	70.6±9.0	59.1±10.1	<0.001	70.8±8.0	58.8±10.1	<0.001
Male	290 (64.0)	2,090 (82.7)	<0.001	321 (63.7)	2,059 (83.2)	<0.001
Body mass index (kg/m ²)	24.2±3.4	25.1±3.2	<0.001	24.3±3.2	25.1±3.2	<0.001
Hypertension	315 (69.5)	1,181 (46.7)	<0.001	335 (66.5)	1,161 (46.9)	<0.001
Diabetes mellitus	223 (49.2)	582 (23.0)	<0.001	228 (45.2)	577 (23.3)	<0.001
Chronic kidney disease*	314 (69.3)	276 (10.9)	<0.001	351 (69.6)	239 (9.7)	<0.001
Dyslipidemia	259 (57.2)	1,545 (61.1)	0.124	296 (58.7)	1,508 (60.9)	0.390
Current smoker	96 (21.2)	1,019 (40.3)	<0.001	124 (24.6)	991 (40.0)	<0.001
Prior myocardial infarction	25 (5.5)	83 (3.3)	0.027	26 (5.2)	82 (3.3)	0.059
Prior percutaneous coronary intervention	69 (15.2)	189 (7.5)	<0.001	57 (11.3)	201 (8.1)	0.025
Prior coronary bypass graft	8 (1.8)	10 (0.4)	0.002	6 (1.2)	12 (0.5)	0.121
Prior stroke	72 (15.9)	50 (2.0)	<0.001	37 (7.3)	85 (3.4)	<0.001
Admission via emergency department	299 (66.0)	1,744 (69.0)	0.220	376 (74.6)	1,667 (67.4)	0.002
Clinical presentation			0.275			0.016
Unstable angina	149 (32.9)	763 (30.2)		131 (26.0)	781 (31.5)	
Myocardial infarction	304 (67.1)	1,764 (69.8)		373 (74.0)	1,695 (68.5)	
Primary percutaneous coronary intervention†	123 (27.2)	905 (35.8)	<0.001	190 (37.7)	838 (33.8)	0.108
Transfemoral approach	222 (49.0)	1,093 (43.3)	0.026	260 (51.6)	1,055 (42.6)	<0.001
Multivessel coronary artery disease	314 (69.3)	1,339 (53.0)	<0.001	339 (67.1)	1,315 (53.1)	<0.001
Multi-lesion intervention	100 (22.1)	503 (19.9)	0.320	110 (21.8)	493 (19.9)	0.361
Multivessel intervention	84 (18.5)	421 (16.7)	0.360	94 (18.7)	411 (16.6)	0.292
Treated lesions per patient	1.26±0.52	1.23±0.50	0.280	1.27±0.55	1.23±0.50	0.186
Total number of stents per patient	1.40±0.67	1.36±0.67	0.362	1.42±0.74	1.36±0.65	0.090
Total stent length per patient (mm)	36.0±20.8	34.5±20.5	0.145	36.6±21.7	34.3±20.3	0.026
Antithrombotic drug before intervention‡						
Unfractionated heparin	280 (61.8)	1,563 (61.9)	1.000	326 (64.7)	1,517 (61.3)	0.165
Low-molecular-weight heparin	31 (6.8)	230 (9.1)	0.140	42 (8.3)	219 (8.8)	0.776
Glycoprotein IIb/IIIa inhibitors	24 (5.3)	170 (6.7)	0.302	39 (7.7)	155 (6.3)	0.260

Data are presented as mean±SD or number (%).

ARC-HBR = Academic Research Consortium for High Bleeding Risk; HBR = high bleeding risk; PRECISE-DAPT = Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent Dual Antiplatelet Therapy.

*Chronic kidney disease was defined as an estimated glomerular filtration rate of less than 60 mL per minute per 1.73 m² of body-surface area; †Primary percutaneous coronary intervention was defined as an emergent percutaneous coronary intervention without previous treatment of fibrinolytic agents; ‡Drugs before intervention were those which were given in the hospital immediately before the procedure.

regard to NACE, there was no significant interaction between therapy and HBR by ARC-HBR criteria (p-interaction=0.400). Ticagrelor monotherapy after 3-month DAPT was associated with lower rate of NACE than ticagrelor-based 12-month DAPT in both HBR (HR, 0.30; 95% CI, 0.11–0.80) and non-HBR patients (HR, 0.49; 95% CI, 0.27–0.89). Significant interactions

Table 2. Clinical outcomes by HBR

	ARC-HBR				PRECISE-DAPT			
	HBR (n=453)	Non-HBR (n=2,527)	HR (95% CI)	p-value	HBR (n=504)	Non-HBR (n=2,476)	HR (95% CI)	p-value
Primary outcome								
NACE	24 (5.4)	48 (1.9)	2.87 (1.76–4.69)	<0.001	27 (5.5)	45 (1.9)	3.09 (1.92–4.98)	<0.001
Secondary outcome								
Major bleeding	12 (2.7)	14 (0.6)	4.91 (2.27–10.61)	<0.001	14 (2.9)	12 (0.5)	5.96 (2.76–12.88)	<0.001
MACCE	14 (3.2)	34 (1.4)	2.34 (1.26–4.36)	0.006	15 (3.1)	33 (1.4)	2.31 (1.25–4.25)	0.006
Death	10 (2.3)	5 (0.2)	11.37 (3.89–33.25)	<0.001	9 (1.9)	6 (0.2)	7.59 (2.70–21.31)	<0.001
Myocardial infarction	2 (0.5)	11 (0.4)	1.03 (0.23–4.65)	0.968	3 (0.6)	10 (0.4)	1.52 (0.42–5.51)	0.524
Stent thrombosis	1 (0.2)	2 (0.1)	2.84 (0.26–31.29)	0.373	0 (0)	3 (0.1)	NA	NA
Stroke	2 (0.5)	6 (0.2)	1.89 (0.38–0.38)	0.426	3 (0.6)	5 (0.2)	3.04 (0.73–12.71)	0.109
Target-vessel revascularization	1 (0.2)	16 (0.6)	0.36 (0.05–2.68)	0.295	1 (0.2)	16 (0.7)	0.32 (0.04–2.39)	0.239

Data are presented as number (% of the cumulative rates at 12 months according to Kaplan-Meier event rates). The p-values are from log-rank tests.

ARC-HBR = Academic Research Consortium for High Bleeding Risk; CI = confidence interval; HBR = high bleeding risk; HR = hazard ratio; MACCE = major adverse cardiac and cerebrovascular event; NA = not applicable; NACE = net adverse clinical event; PRECISE-DAPT = Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent Dual Antiplatelet Therapy.




Figure 2. Time-to-event curves for clinical events by HBR. NACE (A, D), major bleeding (B, E), and MACCEs (C, F). ARC-HBR = Academic Research Consortium for High Bleeding Risk; CI = confidence interval; HBR = high bleeding risk; HR = hazard ratio; MACCE = major adverse cardiac and cerebrovascular event; NACE = net adverse clinical event; PCI = percutaneous coronary intervention; PRECISE-DAPT = Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent Dual Antiplatelet Therapy.




Figure 3. Time-to-event curves for clinical events by HBR and antiplatelet therapy strategy. NACE (A, D), major bleeding (B, E), and MACCEs (C, F). ARC-HBR = Academic Research Consortium for High Bleeding Risk; CI = confidence interval; HBR = high bleeding risk; HR = hazard ratio; MACCE = major adverse cardiac and cerebrovascular event; NACE = net adverse clinical event; PCI = percutaneous coronary intervention; PRECISE-DAPT = Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent Dual Antiplatelet Therapy.

Table 3. Clinical outcomes by HBR and antiplatelet therapy strategy

	ARC-HBR				PRECISE-DAPT								
	HBR		Non-HBR		HBR		Non-HBR		Therapy × HBR Interaction p-value	Therapy × HBR Interaction p-value			
	Ticagrelor-based 12-month DAPT (n=241)	Ticagrelor monotherapy after 3-month DAPT (n=212)	HR (95% CI)	Ticagrelor-based 12-month DAPT (n=1,250)	Ticagrelor monotherapy after 3-month DAPT (n=1,277)	HR (95% CI)	Ticagrelor-based 12-month DAPT (n=263)	Ticagrelor monotherapy after 3-month DAPT (n=241)					
Primary outcome													
NACE	19 (8.0)	5 (2.4)	0.30 (0.11-0.80)	32 (2.6)	16 (1.3)	0.49 (0.27-0.89)	22 (8.5)	5 (2.2)	0.25 (0.09-0.66)	29 (2.4)	16 (1.3)	0.54 (0.29-0.99)	0.178
Secondary outcome													
Major bleeding	11 (4.7)	1 (0.5)	0.10 (0.01-0.81)	12 (1.0)	2 (0.2)	0.16 (0.04-0.73)	13 (5.1)	1 (0.4)	0.09 (0.01-0.65)	10 (0.8)	2 (0.2)	0.20 (0.04-0.89)	0.518
MACCE	10 (4.2)	4 (2.0)	0.46 (0.15-1.47)	20 (1.6)	14 (1.1)	0.69 (0.35-1.36)	11 (4.3)	4 (1.7)	0.41 (0.13-1.27)	19 (1.6)	14 (1.1)	0.73 (0.36-1.45)	0.389
Death	6 (2.5)	4 (2.0)	0.77 (0.22-2.74)	2 (0.2)	3 (0.2)	1.48 (0.25-8.84)	5 (1.9)	4 (1.7)	0.90 (0.24-3.36)	3 (0.2)	3 (0.2)	0.99 (0.20-4.89)	0.932
Myocardial infarction	2 (0.8)	0 (0)	NA	8 (0.7)	3 (0.2)	0.37 (0.10-1.39)	3 (1.2)	0 (0)	NA	7 (0.6)	3 (0.2)	0.42 (0.11-1.63)	NA
Stent thrombosis	1 (0.4)	0 (0)	NA	0 (0)	2 (0.2)	NA	0 (0)	0 (0)	NA	1 (0.1)	2 (0.2)	1.97 (0.18-21.75)	NA
Stroke	2 (0.8)	0 (0)	NA	3 (0.2)	3 (0.2)	0.98 (0.20-4.87)	3 (1.2)	0 (0)	NA	2 (0.2)	3 (0.2)	1.48 (0.25-8.86)	NA
Target-vessel revascularization	1 (0.4)	0 (0)	NA	8 (0.7)	8 (0.6)	0.98 (0.37-2.62)	1 (0.4)	0 (0)	NA	8 (0.7)	8 (0.7)	0.99 (0.37-2.63)	NA

Data are presented as number (% of the cumulative rates at 12 months according to Kaplan-Meier event rates). The p-values are from Cox regression test of therapy × HBR interaction. ARC-HBR = Academic Research Consortium for High Bleeding Risk; CI = confidence interval; DAPT = dual-antiplatelet therapy; HBR = high bleeding risk; HR = hazard ratio; MACCE = major adverse cardiac and cerebrovascular event; NA = not applicable; NACE = net adverse clinical event; PRECISE-DAPT = Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent Dual Antiplatelet Therapy.

were not observed with regard to major bleeding (HBR: HR, 0.10; 95% CI, 0.01–0.81; non-HBR: HR, 0.16; 95% CI, 0.04–0.73; p-interaction=0.728) and MACCE (HBR: HR, 0.46; 95% CI, 0.15–1.47; non-HBR: HR, 0.69; 95% CI, 0.35–1.36; p-interaction=0.560). Results were consistent by PRECISE-DAPT score. With respect to NACE, there was no significant interaction between therapy and HBR by PRECISE-DAPT score (p-interaction=0.178). Ticagrelor monotherapy after 3-month DAPT was associated with lower rate of NACE than ticagrelor-based 12-month DAPT in both HBR (HR, 0.25; 95% CI, 0.09–0.66) and non-HBR patients (HR, 0.54; 95% CI, 0.29–0.99). Significant interactions were not observed with respect to major bleeding (HBR: HR, 0.09; 95% CI, 0.01–0.65; non-HBR: HR, 0.20; 95% CI, 0.04–0.89; p-interaction=0.518) and MACCE (HBR: HR, 0.41; 95% CI, 0.13–1.27; non-HBR: HR, 0.73; 95% CI, 0.36–1.45; p-interaction=0.389). These findings were also confirmed in per-protocol analysis and without excluding the patients with adverse events within 3 months (**Supplementary Figures 2 and 3**). There were no significant interactions between therapy and HBR for NACE, major bleeding, and MACCE.

DISCUSSION

The main findings of this post-hoc analysis were as follows: 1) for patients presenting with ACS and treated with DESs, HBR was associated with higher rate of primary outcome, not only including bleeding but also ischemic events; 2) ticagrelor monotherapy after 3-month DAPT was associated with lower rate of primary outcome and major bleeding than ticagrelor-based 12-month DAPT regardless of HBR, with no significant interaction between therapy and HBR; and 3) These findings were consistent regardless of HBR definition by ARC-HBR criteria or by PRECISE-DAPT score.

Current guidelines recommend 12-month DAPT including novel antiplatelet agents with higher potency in patients treated with DESs for ACS.^{1,2,8,9} However, major concerns toward increased risk of bleeding by prolonged DAPT have been raised and shortened duration of DAPT may be considered in patients with HBR.¹⁸ Therefore, it is of great importance to identify patients with HBR. The TWILIGHT trial included patients with not only high ischemic risk but also HBR, and it showed that ticagrelor monotherapy followed by 3-month DAPT after PCI was associated with lower bleeding events than ongoing ticagrelor plus aspirin, without an increased risk for ischemic events.¹⁰ The TICO trial exclusively focused on patients with ACS and demonstrated that ticagrelor monotherapy after 3-month DAPT was associated with a lower incidence of NACE than currently recommended ticagrelor-based 12-month DAPT.¹¹ In this post-hoc analysis, we classified the TICO subjects into HBR and non-HBR patients by 2 different definitions of HBR that were proposed recently: the ARC-HBR criteria and the PRECISE-DAPT score.^{3,4} The ARC-HBR criteria, which consists of 20 clinical criteria as major and minor by consensus and supported by published evidence, and the PRECISE-DAPT score, which consists of 5 variables developed from a large pooled dataset of recent randomized clinical trials, were excellent in identifying those with HBR after PCI and associated with higher risk of bleeding as well as ischemic events when validated to Bern PCI registry.^{3,7} In our study, the rate of NACE, including major bleeding and MACCE, was significantly higher in HBR versus non-HBR patients, regardless of the HBR definition which was consistent to those of previous studies.

Several studies have investigated the association between HBR and the duration of DAPT. Costa et al. reported that compared to a short DAPT duration (3–6 months), a long DAPT duration

(12–24 months) increased bleeding in patients with HBR but not in those with non-HBR (p for interaction=0.007).⁴⁾ According to the secondary analysis, 20.8% of patients underwent complex PCI, which is associated with higher ischemic risk and can be alleviated by a long DAPT duration.¹⁹⁾ However, the benefit of a long DAPT duration for reducing ischemic events was present only in patients with non-HBR for both complex (absolute risk difference: -3.86%, p=0.05) and noncomplex PCI (absolute risk difference: -1.14%, p=0.04). The benefit was not present in HBR patients for both complex (absolute risk difference: +1.30%, p=0.76) and noncomplex PCI (absolute risk difference: +1.45%, p=0.39), which indicated that bleeding, more than ischemic risk, should inform on making decision for DAPT duration after PCI.¹⁹⁾ Different from previous studies, only ultrathin bioresorbable polymer sirolimus-eluting stents (Orsiro; Biotronik AG, Bülach, Switzerland) were used in the TICO trial, which have shown superior clinical outcomes over other DESs, probably due to reduced thrombus formation, inflammation, and neointimal proliferation.¹¹⁾¹²⁾²⁰⁾²¹⁾ Likewise, as for DAPT strategies utilizing P2Y₁₂ receptor inhibitor, ticagrelor was used, which showed superior clinical outcomes over clopidogrel in reducing ischemic events.⁸⁾ Therefore, long DAPT duration for the purpose of preventing ischemic events is less likely needed compared to other DESs and P2Y₁₂ receptor inhibitors, which raises the issue of bleeding risk to be considered as a priority over ischemic risk. Furthermore, although the definition of short and long duration of DAPT was heterogeneous (3–6 months vs. 12–24 months) with aspirin continuation after DAPT in previous studies, the definition was strictly confined to 3-month DAPT with P2Y₁₂ receptor inhibitor continuation versus 12-month DAPT in the TICO trial.⁴⁾¹¹⁾¹⁹⁾ These factors may have contributed to the findings of this study, which showed the beneficial effect of ticagrelor monotherapy after short DAPT duration in both HBR and non-HBR patients. However, the benefit of ticagrelor monotherapy after 3-month DAPT tended to be more prominent in HBR patients which is in line with the secondary analysis of the TWILIGHT trial regarding HBR.²²⁾

This study has several limitations. First, the present study was not prespecified in the TICO trial protocol; however, the definition of HBR by ARC-HBR criteria or PRECISE-DAPT score was not developed at the time of trial design. Therefore, our findings need to be interpreted only as hypothesis-generating and require further prospective randomized trials. Second, the number of HBR patients is relatively small compared with non-HBR patients. Not all criteria for ARC-HBR were available for analysis, and while some of those were not routinely collected in the TICO trial, others were in fact exclusion criteria for the trial since the patients with an extremely high risk of bleeding who may be contraindicated for antiplatelet agents were excluded as in other trials regarding antiplatelet therapy strategies. Nevertheless, statistical significance for the benefit of ticagrelor monotherapy after 3-month DAPT toward primary outcome and major bleeding was found even in relatively small number of HBR patients. Third, the patients who encountered adverse events during the first 3 months after PCI were excluded in primary analysis, since the therapy during the first 3 months was the same in both therapy strategies. However, key findings of this post-hoc analysis (interactions between therapy and HBR for clinical outcomes) were also confirmed without excluding the patients with adverse events within 3 months. Fourth, although the superiority assumption was met, the lower than expected event rate of the primary outcome was noted in the TICO trial, which was consistent in this post-hoc analysis.

In conclusions, in patients with ACS treated with DESs, ticagrelor monotherapy after 3-month DAPT was associated with a lower incidence of NACE and major bleeding than currently recommended ticagrelor-based 12-month DAPT regardless of HBR, and the magnitude of therapy effect did not differ between HBR and non-HBR.

SUPPLEMENTARY MATERIALS

Supplementary Table 1

Baseline characteristics by ARC-HBR criteria and antiplatelet therapy strategy

[Click here to view](#)

Supplementary Table 2

Baseline characteristics by PRECISE-DAPT score based HBR and antiplatelet therapy strategy

[Click here to view](#)

Supplementary Table 3

Baseline characteristics by HBR for per-protocol population

[Click here to view](#)

Supplementary Table 4

Baseline characteristics by ARC-HBR criteria and antiplatelet therapy strategy for per-protocol population

[Click here to view](#)

Supplementary Table 5

Baseline characteristics by PRECISE-DAPT score based HBR and antiplatelet therapy strategy for per-protocol population

[Click here to view](#)

Supplementary Figure 1

Time-to-event curves for clinical events by HBR for per-protocol population. NACE (A, D), major bleeding (B, E), and MACCEs (C, F).

[Click here to view](#)

Supplementary Figure 2

Time-to-event curves for clinical events by HBR and antiplatelet therapy strategy for per-protocol population. NACE (A, D), major bleeding (B, E), and MACCEs (C, F).

[Click here to view](#)

Supplementary Figure 3

Time-to-event curves for clinical events by HBR and antiplatelet therapy strategy for overall TICO patients. NACE (A, D), major bleeding (B, E), and MACCEs (C, F).

[Click here to view](#)

REFERENCES

1. Levine GN, Bates ER, Bittl JA, et al. 2016 ACC/AHA guideline focused update on duration of dual antiplatelet therapy in patients with coronary artery disease: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *J Am Coll Cardiol* 2016;68:1082-115.
[PUBMED](#) | [CROSSREF](#)
2. Valgimigli M, Bueno H, Byrne RA, et al. 2017 ESC focused update on dual antiplatelet therapy in coronary artery disease developed in collaboration with EACTS: the task force for dual antiplatelet therapy in coronary artery disease of the European Society of Cardiology (ESC) and of the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J* 2018;39:213-60.
[PUBMED](#) | [CROSSREF](#)
3. Urban P, Mehran R, Collieran R, et al. Defining high bleeding risk in patients undergoing percutaneous coronary intervention. *Circulation* 2019;140:240-61.
[PUBMED](#) | [CROSSREF](#)
4. Costa F, van Klaveren D, James S, et al. Derivation and validation of the predicting bleeding complications in patients undergoing stent implantation and subsequent dual antiplatelet therapy (PRECISE-DAPT) score: a pooled analysis of individual-patient datasets from clinical trials. *Lancet* 2017;389:1025-34.
[PUBMED](#) | [CROSSREF](#)
5. Ueki Y, Karagiannis A, Zanchin C, et al. Validation of high-risk features for stent-related ischemic events as endorsed by the 2017 DAPT guidelines. *JACC Cardiovasc Interv* 2019;12:820-30.
[PUBMED](#) | [CROSSREF](#)
6. Ueki Y, Bar S, Losdat S, et al. Validation of the Academic Research Consortium for High Bleeding Risk (ARC-HBR) criteria in patients undergoing percutaneous coronary intervention and comparison with contemporary bleeding risk scores. *EuroIntervention* 2020;16:371-9.
[PUBMED](#) | [CROSSREF](#)
7. Natsuaki M, Morimoto T, Shiomi H, et al. Application of the academic research consortium high bleeding risk criteria in an all-comers registry of percutaneous coronary intervention. *Circ Cardiovasc Interv* 2019;12:e008307.
[PUBMED](#) | [CROSSREF](#)
8. Wallentin L, Becker RC, Budaj A, et al. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2009;361:1045-57.
[PUBMED](#) | [CROSSREF](#)
9. Wiviott SD, Braunwald E, McCabe CH, et al. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2007;357:2001-15.
[PUBMED](#) | [CROSSREF](#)
10. Mehran R, Baber U, Sharma SK, et al. Ticagrelor with or without aspirin in high-risk patients after PCI. *N Engl J Med* 2019;381:2032-42.
[PUBMED](#) | [CROSSREF](#)
11. Kim BK, Hong SJ, Cho YH, et al. Effect of ticagrelor monotherapy vs ticagrelor with aspirin on major bleeding and cardiovascular events in patients with acute coronary syndrome: the TICO randomized clinical trial. *JAMA* 2020;323:2407-16.
[PUBMED](#) | [CROSSREF](#)
12. Kim C, Hong SJ, Shin DH, et al. Randomized evaluation of ticagrelor monotherapy after 3-month dual-antiplatelet therapy in patients with acute coronary syndrome treated with new-generation sirolimus-eluting stents: TICO trial rationale and design. *Am Heart J* 2019;212:45-52.
[PUBMED](#) | [CROSSREF](#)
13. Mehran R, Rao SV, Bhatt DL, et al. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium. *Circulation* 2011;123:2736-47.
[PUBMED](#) | [CROSSREF](#)
14. Thygesen K, Alpert JS, Jaffe AS, et al. Third universal definition of myocardial infarction. *Circulation* 2012;126:2020-35.
[PUBMED](#) | [CROSSREF](#)
15. Cutlip DE, Windecker S, Mehran R, et al. Clinical end points in coronary stent trials: a case for standardized definitions. *Circulation* 2007;115:2344-51.
[PUBMED](#) | [CROSSREF](#)
16. Sacco RL, Kasner SE, Broderick JP, et al. An updated definition of stroke for the 21st century: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2013;44:2064-89.
[PUBMED](#) | [CROSSREF](#)

17. Kim BK, Hong MK, Shin DH, et al. A new strategy for discontinuation of dual antiplatelet therapy: the RESET Trial (REal Safety and Efficacy of 3-month dual antiplatelet Therapy following Endeavor zotarolimus-eluting stent implantation). *J Am Coll Cardiol* 2012;60:1340-8.
[PUBMED](#) | [CROSSREF](#)
18. Capodanno D, Alfonso F, Levine GN, Valgimigli M, Angiolillo DJ. ACC/AHA versus ESC guidelines on dual antiplatelet therapy: JACC guideline comparison. *J Am Coll Cardiol* 2018;72:2915-31.
[PUBMED](#) | [CROSSREF](#)
19. Costa F, Van Klaveren D, Feres F, et al. Dual antiplatelet therapy duration based on ischemic and bleeding risks after coronary stenting. *J Am Coll Cardiol* 2019;73:741-54.
[PUBMED](#) | [CROSSREF](#)
20. Kandzari DE, Mauri L, Koolen JJ, et al. Ultrathin, bioresorbable polymer sirolimus-eluting stents versus thin, durable polymer everolimus-eluting stents in patients undergoing coronary revascularisation (BIOFLOW V): a randomised trial. *Lancet* 2017;390:1843-52.
[PUBMED](#) | [CROSSREF](#)
21. Roguin A, Kandzari DE, Marcusohn E, et al. Subgroup analysis comparing ultrathin, bioresorbable polymer sirolimus-eluting stents versus thin, durable polymer everolimus-eluting stents in acute coronary syndrome patients. *Circ Cardiovasc Interv* 2018;11:e007331.
[PUBMED](#) | [CROSSREF](#)
22. Escaned J, Cao D, Baber U, et al. Ticagrelor monotherapy in patients at high bleeding risk undergoing percutaneous coronary intervention: TWILIGHT-HBR. *Eur Heart J* 2021;42:4624-34.
[PUBMED](#) | [CROSSREF](#)