



Clinical Research Article

Korean J Anesthesiol 2021;74(2):134–141
<https://doi.org/10.4097/kja.20114>
pISSN 2005–6419 • eISSN 2005–7563

Received: March 16, 2020

Revised: April 26, 2020 (1st); May 25, 2020 (2nd)

Accepted: June 8, 2020

Corresponding author:

Gaurav Jain, M.D.

Department of Anesthesiology, All India
Institute of Medical Sciences, Virbhadr
Marg, Rishikesh, Uttarakhand 249203, India
Tel: +91-8808631209

Fax: +91-135-2460994

Email: gauravhld@gmail.com

ORCID: <https://orcid.org/0000-0002-1205-7237>

Effectiveness of four ultrasonographic parameters as predictors of difficult intubation in patients without anticipated difficult airway

Rishabh Agarwal, Gaurav Jain, Ankit Agarwal, Nishith Govil

Department of Anesthesiology, All India Institute of Medical Sciences, Rishikesh, India

Background: Predicting difficult intubation (DI) is a key challenge, as no single clinical predictor is sufficiently valid to predict the outcome. We evaluated the effectiveness of four upper airway ultrasonographic parameters in predicting DI. The validity of the models using combinations of ultrasonography-based parameters was also investigated.

Methods: This prospective, observational, double-blinded cohort trial enrolled 1,043 surgical patients classified as American Society of Anesthesiologists physical status I–III without anticipated difficult airway. Preoperatively, their tongue thickness (TT), invisibility of hyoid bone (VH), and anterior neck soft tissue thickness from the skin to thyrohyoid membrane (ST) and hyoid bone (SH) were measured by sublingual and submandibular ultrasonography. The logistic regression, Youden index, and receiver operator characteristic analysis results were reported.

Results: Overall, 58 (5.6%) patients were classified as DI. The TT, SH, ST, and VH had accuracies of 78.4%, 85.0%, 84.7%, and 84.9%, respectively. The optimal values of TT, SH, and ST for predicting DI were > 5.8 cm (sensitivity: 84.5%, specificity: 78.1%, AUC: 0.880), > 1.4 cm (sensitivity: 81%, specificity: 85.2%, AUC: 0.898), and > 2.4 cm (sensitivity: 75.9%, specificity: 85.2%, AUC: 0.885), respectively. VH had a sensitivity and specificity of 72.4% and 85.6% (AUC: 0.790). The AUC values of the five models (with combinations of three or four parameters) ranged from 0.975–0.992. ST and VH had a significant impact on the individual models.

Conclusions: SH had the best accuracy. Individual parameters showed limited validity. The model including all four parameters offered the best diagnostic value.

Keywords: Airway management; Diagnostic ultrasound; General anesthesia; Hyoid bone; Intubation; Laryngoscopy; Tongue.

Introduction

Securing the airway is a vital component in the clinical practice of anesthesia. Difficult intubation (DI) is prone to potential complications, ranging from minimal airway edema to life-threatening events. Predicting DI during the preoperative assessment is a key challenge, as no single clinical predictor is sufficiently valid for predicting the outcomes. Various imaging techniques have been under consideration for evaluation of the airways, but each has specific limitations, such as radiation exposure, high cost, and procedure time, etc.

Ultrasonography is a non-invasive and quick bedside tool that allows easy visualization of the neck anatomy and assessment of the airway [1,2]. Various ultrasonography-related

© The Korean Society of Anesthesiologists, 2021

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

parameters such as tongue thickness (TT), the invisibility of hyoid bone (VH), mandible condylar mobility, and anterior neck soft tissue thickness from the skin to the thyrohyoid membrane (ST) and the hyoid bone (SH), respectively, have the potential of predicting difficult airway. The current literature is, however, limited to small studies, restricted further by the low incidence of DI [3–5]. Thus, the validity of ultrasonography-based parameters in predicting DI requires further exploration. We hypothesized that upper airway ultrasonographic parameters including TT, SH, ST, and VH would reliably predict DI during preoperative assessment in patients without anticipated difficult airway. We preferred these parameters considering the ease and rapidity in locating their anatomical landmarks to allow precise measurements, their potential ability to predict DI, and the limited role of clinical screening in their evaluation. Our primary aim was to evaluate the effectiveness of aforesaid ultrasonographic parameters in predicting DI by comparing them between the DI and easy intubation (EI) groups. We also analyzed the validity of various models with combined ultrasonography-based parameters in predicting DI.

Materials and Methods

After obtaining institutional ethical approval (AIIMS/IEC/18/85) and written informed consent, patients of both sexes classified as American Society of Anesthesiologists physical status I–III, aged 18 to 60 years, undergoing tracheal intubation for surgery under general anesthesia, were included in this prospective, obser-

vational, cohort study conducted between August 2018 to July 2019 (Indian Clinical Trial Registry No: CTRI/2018/07/014786). This clinical research was done following the ethical principles for medical research involving human subjects in accordance with the Helsinki Declaration 2013. An experienced investigator conducted a difficult airway assessment during the preoperative visit. Those with upper airway anatomical anomaly, trauma, or tumor; history of a difficult airway; and difficult airway on pre-anesthetic check-up and those requiring deviation in the study protocol were excluded. A modified Mallampati test (MMP) grade of 3 or 4, small thyromental distance (< 6.5 cm), and small inter-incisor distance (< 3 cm) indicate the presence of difficult airway.

All enrolled patients underwent a duly explained ultrasonographic examination (Logic eR7, GE Medical Systems Co. Ltd., China) of the upper airway in the pre-operative room. A skilled investigator (≥ 5 years of experience in airway ultrasonography) performed the procedure and recorded the measurements. For sublingual ultrasonography, patients were positioned in the sitting, neutral head position [4]. A curved ultrasound probe (4.2–10.0 MHz in a sterile cover) was placed intra-orally under the patient's tongue in a longitudinal orientation (perpendicular to the face) and advanced backwards as far as the patient felt comfortable (Fig. 1A). The hyoid bone was noted in the obtained image (Fig. 1D). For submandibular ultrasonography, all patients were asked to remain in a supine and extended neck posture, to keep their mouth closed, and to remain quiet, with the tip of tongue relaxed and just touching the incisors [6]. A curvilinear ultrasound




Fig. 1. Schematic diagram showing the position of ultrasound probe and the corresponding images of upper airway. (A) Probe placed intraorally under the patient's tongue, (B) positioned beneath the mentum along the mid-sagittal plane, and (C) rotated 90° in the same position. Ultrasonography image showing (D) visibility of hyoid bone, (E) measurement of TT, (F) measurement of SH, and (G) measurement of ST. A: air-membrane interface, E: epiglottis, G: geniohyoid, H: hyoid, M: mentum, S: skin, SH: soft tissue thickness from skin to hyoid bone, ST: soft tissue thickness from skin to thyrohyoid membrane, T: tongue, TT: tongue thickness.

probe (2–5 MHz) was positioned beneath the mentum along the mid-sagittal plane, adjusted to obtain a clear image of the tongue contour (Fig. 1B). The maximum vertical length from the tongue surface to the submental skin was noted and defined as TT (Fig. 1E). In the same position, the transducer was rotated to 90° and SH and ST, were measured (Figs. 1C, 1F and 1G).

Thereafter, the patients were moved to the operation theatre, and standard monitors were applied. After adequate pre-oxygenation, general anesthesia was induced with propofol (1–2 mg/kg IV), midazolam (0.05 mg/kg IV), fentanyl (0.004 mg/kg IV), and vecuronium (0.1 mg/kg IV). After 3 minutes of mask ventilation, an experienced investigator (≥ 5 years of experience in intubation procedure) performed the laryngoscopy (Macintosh blade size 3 or 4) in the sniffing position. To facilitate laryngeal view, external laryngeal manipulation was allowed, and intubation was attempted. In case of failed attempts, standard protocols were followed as per unanticipated DI guidelines [7]. ‘DI’ was defined as the placement of the endotracheal tube by using conventional laryngoscopy that required > 2 attempts, lasted > 10 min, or required alternate methods [7]. The ‘time taken for intubation’ was defined as the time point from initiation of the first direct laryngoscopy attempt to confirmation of successful endotracheal intubation by continuous waveform capnography. The difficult airway cart included intubating stylet (IS), McCoy blade (MB), intubating laryngeal mask airway (ILMA), video laryngoscope (VL), light wand (LW), fiberoptic bronchoscope, and percutaneous cricothyroidotomy. The endpoint of the study was tracheal intubation, based on which all the included patients were categorized as EI or DI.

Clinical airway assessment data and ultrasonography-based data were categorized according to the group for statistical analysis and interpretation. The investigator who performed the ultrasonographic examination was blinded to the preoperative airway assessment data, intubation procedure, and group allocation. Another investigator carried out the preoperative airway assessment, intubation procedure and group allocation, but was blinded to ultrasonographic parameters. Another investigator blinded to study protocols performed the data analysis.

Statistical analysis

The sample size was calculated using the sample size calculator of the University of California, San Francisco, US [8]. Taking an alpha error of 5%, power of 80%, the incidence of unanticipated DI as 5% (a weighted average of data from the literature [1–9.5%]), the sample size was calculated as 1,030 (EI: 978, DI: 52), considering an effect size of 0.8 (estimated from initial pilot ob-

servations) for the primary outcome [3,9]. Statistical analysis was performed using MedCalc software version 19.0.7 (Acaciaaan, Belgium). The results were presented as descriptive statistics, summarized as mean (SD) or number (percentage). Data were analyzed by logistic regression, receiver operator characteristic (ROC) curve, and Youden index to calculate the diagnostic validity profile of the outcome variables. The continuous variables were compared by unpaired student t-test. The categorical variables were compared by Chi-square test/Fisher’s exact test. A $P < 0.05$ was considered significant.

Results

We assessed 1,252 patients for eligibility, of which 1,043 were included in the study (no dropouts). The demographic profile was comparable among the groups (Table 1). In all, 985 patients were classified as EI, as 58 patients as DI (Fig. 2). The EI group had significantly lower MMP grade than the DI group. On direct laryngoscopy, the EI group had a Cormack-Lehane (CL) grade of 1 or 2, while DI group had a CL grade of 2 or 3 with some grade 4 cases as well, which varied significantly on intergroup comparison (Table 1). Further, 802 patients could be intubated in the first attempt, while the remaining required repeat attempts or alternate methods, such as IS+MB in 27 patients, ILMA in 24 patients, VL in 10 patients, and LW in one patient, with success rates of 96.3%, 91.6%, 90%, and 100%, respectively.

The means of the upper airway ultrasonography-based parameters TT, SH, and ST were significantly greater in the DI group than in the EI group ($P < 0.001$, respectively) (Table 2). The VH was 14.4% in the EI group versus 72.4% in the DI group ($P < 0.001$). For validity analysis, an ROC curve was plotted for each ultrasonographic parameter. The optimal criterion for TT to predict DI was found to be > 5.8 cm (sensitivity: 84.5%, specificity: 78.1%) with an area under the curve (AUC) of 0.880. For SH, the optimal value was > 1.4 cm (sensitivity: 81%, specificity: 85.2%), with an AUC of 0.898; for ST, it was > 2.4 cm (sensitivity: 75.9%, specificity: 85.2%), with an AUC of 0.885. The sensitivity and specificity for VH were 72.4% and 85.6%, respectively, with an AUC of 0.790 (Table 2). We also plotted a graph to determine the changes in sensitivity and specificity for each threshold value of TT, SH, and ST. An increase in TT, SH, and ST thresholds led to an increase in the specificity but a decrease in the sensitivity for identifying DI (Fig. 3). The TT, SH, ST, and VH had the accuracies of 78.4%, 85.0%, 84.7%, and 84.9%, respectively. On univariate analysis, the odds ratio (OR) for TT was 1.06, indicating a 6% increase in the log-odds of DI per millimeter increase in TT. Similarly, the ORs for SH, ST, and VH were 1.07, 1.10, and 15.58, re-

Table 1. Characteristics of Included Patients

Variable	Easy intubation (n = 985)	Difficult intubation (n = 58)	P value
Age (yr)	41.2 ± 13.0	39.1 ± 12.9	0.247
Weight (kg)	60.5 ± 12.7	60.0 ± 11.2	0.763
Male	349 (35.4)	15 (25.9)	0.157
MMP			
Grade 1	720 (73.1)	27 (46.5)	< 0.001
Grade 2	265 (26.9)	31 (53.4)	< 0.001
CL			
Grade 1	715 (72.6)	-	< 0.001
Grade 2	267 (27.1)	26 (44.8)	< 0.001
Grade 3	3 (0.3)	27 (46.5)	< 0.001
Grade 4	-	5 (8.6)	< 0.001
Number of intubation attempts			
1	802 (81.4)	0	-
2	183 (18.6)	0	-
3	0	54 (93.1)	-
4	0	4 (6.9)	-
Average time taken for intubation (s)	54.5 ± 6.3	287.1 ± 20.4	< 0.001

Values are presented as mean ± SD or number (%). MMP: Modified Mallampati, CL: Cormack-Lehane. A P value < 0.05 was considered to be statistically significant.




Fig. 2. CONSORT flowchart of patient selection. TT: tongue thickness, SH: anterior neck soft tissue thickness from skin to hyoid bone, ST: anterior neck soft tissue thickness from skin to thyrohyoid membrane, VH: invisibility of hyoid bone.

spectively. A wide CI (8.52, 28.47) was observed for VH, indicating a low level of precision (Table 2). A subgroup analysis was also performed to compare the ultrasonographic parameters in those with CL grade 2 and a similar difference was observed between the ultrasonographic parameters.

The validity of five models based on combined ultrasonography-based parameters was also assessed through multiple logistic regression-derived ROC analysis (Table 3, Fig. 4). ‘Model 1’, which included all the four ultrasonographic parameters, had the highest accuracy with an AUC of 0.992. ‘Model 2’, which included TT, SH,

Table 2. Diagnostic Validity Profile of Ultrasonographic Parameters in Predicting Difficult Intubation

Variable	EI (n = 985)	DI (n = 58)	OC	SE	SP	Accuracy % (95% CI)	OR (95% CI)	AUC	P value
TT (cm)	5.6 ± 0.2	6.1 ± 0.3	> 5.8	0.845	0.781	78.4 (75.8, 80.9)	1.06 (1.04, 1.07)	0.88	< 0.001
SH (cm)	1.3 ± 0.2	1.6 ± 0.2	> 1.4	0.81	0.852	85 (82.6, 87.1)	1.07 (1.06, 1.09)	0.898	< 0.001
ST (cm)	2.2 ± 0.2	2.5 ± 0.2	> 2.4	0.759	0.852	84.7 (82.3, 86.8)	1.1 (1.08, 1.12)	0.885	< 0.001
VH	142 (14.4)	42 (72.4)	-	0.724	0.856	84.9 (82.5, 87.0)	15.58 (8.52, 28.47)	0.79	< 0.001

Values are presented as mean ± SD or number (%). EI: easy intubation group, DI: difficult intubation group, OC: optimal criterion, SE: sensitivity, SP: specificity, OR: odds ratio, AUC: area under curve, TT: tongue thickness, SH: anterior neck soft tissue thickness from skin to hyoid bone, ST: anterior neck soft tissue thickness from skin to thyrohyoid membrane, VH: invisibility of hyoid bone. A P value < 0.05 was considered to be statistically significant.




Fig. 3. Sensitivity and specificity profile for predicting difficult intubation at different values of ultrasonographic parameters. (A) tongue thickness (TT), (B) anterior neck soft tissue thickness from skin to hyoid bone (SH), (C) anterior neck soft tissue thickness from skin to thyrohyoid membrane (ST).

and ST, and 'Model 4', with TT, ST, and VH, had an AUC of 0.981. 'Model 3', which included SH, ST, and VH, had an AUC of 0.975. 'Model 5', which included TT, SH, and VH, had an AUC of 0.978. On evaluating the relative contribution of each parameter, VH had the largest OR, though a wide CI was observed for the OR values of VH. ST was the second variable to have a strong impact on the diagnostic validity of the models (Table 3).

Discussion

This study demonstrated the ability of individual upper airway ultrasonography-based parameters and of models using a combination of these parameters in predicting DI. All the upper-airway ultrasonographic parameters varied significantly between the EI and DI groups. The SH parameter had the highest accuracy, while the TT was least accurate in predicting DI. Among the five models, 'Model 1' with all the four ultrasonographic parameters had the highest validity in terms of the AUC.

For optimal viewing of the glottis during direct laryngoscopy, the soft tissues in the neck need to be mobilized adequately. Adhikari et al. [10] evaluated the anterior neck soft tissue thickness at different ultrasonographic planes and observed that ST and SH correlated strongly with difficult laryngoscopy. We used similar levels for predicting DI and obtained a threshold limit of 1.4 cm for SH and 2.4 cm for ST. Wu et al. [11] found an SH cut-off of 1.28 cm while Adhikari et al. [10] obtained an ST limit of 2.8 cm, for predicting difficult laryngoscopy. Yadav et al. [5] reported cut off values of 0.66 and 2.03 cm, respectively, for SH and ST. It appears that different cut-off targets are required for predicting difficult laryngoscopy and intubation. The small sample sizes in the above studies could have attributed to such variation.

Previous studies have correlated increased TT to difficult laryngoscopy [5,6]. However, the measurements varied significantly according to the anatomic level of the ultrasonographic scans. We measured the TT in the mid-sagittal plane to obtain values of the thickest portion of entire tongue contour. The ROC analysis showed that TT > 5.8 cm predicted risk of DI with a sensitivity of

Table 3. Diagnostic Validity Profile of Different Models in Predicting Difficult Intubation by Combining Ultrasonographic Parameters

Variable	Ultrasonographic parameters	OR (95% CI)	AUC (95% CI)	P value
Model 1	TT	1.06 (1.04, 1.08)	0.992 (0.984, 0.996)	< 0.001
	SH	1.07 (1.04, 1.10)		
	ST	1.11 (1.07, 1.14)		
	VH	21.25 (6.72, 67.20)		
Model 2	TT	1.05 (1.04, 1.07)	0.981 (0.971, 0.988)	< 0.001
	SH	1.07 (1.04, 1.09)		
	ST	1.11 (1.08, 1.14)		
Model 3	SH	1.08 (1.06, 1.10)	0.975 (0.964, 0.984)	< 0.001
	ST	1.10 (1.07, 1.12)		
	VH	23.64 (9.17, 60.10)		
Model 4	TT	1.06 (1.04, 1.08)	0.981 (0.970, 0.988)	< 0.001
	ST	1.11 (1.08, 1.15)		
	VH	18.02 (6.91, 47.03)		
Model 5	TT	1.05 (1.03, 1.06)	0.978 (0.967, 0.986)	< 0.001
	SH	1.07 (1.05, 1.09)		
	VH	21.13 (8.57, 52.09)		

OR: odds ratio, AUC: area under curve, TT: tongue thickness, SH: anterior neck soft tissue thickness from skin to hyoid bone, ST: anterior neck soft tissue thickness from skin to thyrohyoid membrane, VH: invisibility of hyoid bone. A P value < 0.05 was considered to be statistically significant.




Fig. 4. Receiver operating characteristic curves (A-E) showing the usefulness of combined sonographic models (1-5) in predicting difficult intubation. AUC, area under curve; TT, tongue thickness; SH, anterior neck soft tissue thickness from skin to hyoid bone; ST, anterior neck soft tissue thickness from skin to thyrohyoid membrane; VH, invisibility of hyoid bone.

84.5% and specificity of 78.1%, with a 6% increase in the log-odds of DI per millimeter increase in TT values. Yao and Wang [6] observed that a cut-off value of TT as 6.1 cm predicted DI. The observed variation could be due to a difference in baseline demographics and ethnicities of the studied population. The increased TT also relates to the high MMP grade. Our study had a higher proportion of patients with MMP grade 2 in the DI group (mean TT of 6.1 cm) than in the EI group (mean TT of 5.8 cm). A larger TT is expected to mask the visibility of the faucal pillars, which justifies our results. Though MMP grade 1 or 2 patients are classified as EI, previous studies have shown that it as an inadequate stand-alone test for predicting difficult airway [12]. We excluded the patients with MMP grades 3 and 4 to identify the threshold for those in which difficult airway could not be anticipated by clinical examination. As no clinical predictor is sufficient to predict DI, our results serve to complement the pre-anesthetic work-up in anticipating DI. The CL grades of 1 and 2 are considered easy laryngoscopy, but 26 patients with CL grade 2 were classified as DI in our study. The factors that contributed to DI in these patients included visibility of only the posterior part of the glottis or arytenoid during direct laryngoscopy, deep-seated larynx, or reduced space within the oropharynx [13]. These patients were intubated at the third attempt by IS with MB, ILMA, or VL. Hui and Tsui [4] observed a correlation between VH and difficult laryngoscopy. We observed a similar pattern in the DI group, although VH was also observed in 14% of the EI group. The caudal displacement of the hyoid bone reduced its visibility on ultrasonography probe, possibly because of the hypopharyngeal position of the tongue or short rami of the mandible, which compromises the view of the glottis during direct laryngoscopy [4].

We also analyzed the validity of five different models with different combinations of the studied ultrasonographic parameters. The predictive ability of the combinations was significantly better than that of the individual parameters, as evidenced by an increased AUC (0.975, 0.992). 'Model 1' had the highest AUC (0.992) 'Models 2 and 4' were the next best (AUC 0.981), while the 'Model 3' had the lowest AUC (0.975). Considering the inclusion of three parameters in a single submandibular window, with no need of intraoral probe placement, and an acceptable AUC, 'Model 2' seems to be a viable option. To analyze the individual contribution of each variable in the model, we calculated the OR. Taking into account the wide CI in OR values of VH, indicating a low precision level, ST appears to have a significant impact on the validity of models. A large sample size may, however, efficiently delineate the impact of VH. Future studies can attempt to design a scoring system/formula based on combined ultrasonography-based variables, considering the weightage of each ultrasono-

graphic parameter in predicting DI. This study can serve as the base for such trials.

Our study had several limitations. We only analyzed the anatomic parameters of the airway without taking into account the impact of functional components like head positioning, degree of neck extension, skills of performer, etc. which may alter the difficulties encountered during the intubation procedure. To avoid the associated bias, we standardized all the functional parameters in our study. Patients in the EI and DI groups were distributed unevenly, which may have affected the validity of results, but this was as expected, considering the low incidence of unanticipated DI. We could not randomize the patients, but the investigators were blinded. Thus, we expect that the above limitations did not severely distort the results.

In conclusion, the SH had better accuracy than the remaining three ultrasonographic parameters included in the study. Although the individual parameters showed limited validity, a model combining all the four parameters offered better diagnostic profile than each one of them.

Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

Author Contributions

Rishabh Agarwal (Conceptualization; Investigation; Methodology; Resources; Visualization; Writing – original draft; Writing – review & editing)

Gaurav Jain (Conceptualization; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing)

Ankit Agarwal (Investigation; Writing – review & editing)

Nishith Govil (Data curation; Formal analysis; Software; Writing – review & editing)

ORCID

Rishabh Agarwal, <https://orcid.org/0000-0003-2817-0231>

Gaurav Jain, <https://orcid.org/0000-0002-1205-7237>

Ankit Agarwal, <https://orcid.org/0000-0003-4963-7101>

Nishith Govil, <https://orcid.org/0000-0003-3749-6217>

References

1. Singh M, Chin KJ, Chan VW, Wong DT, Prasad GA, Yu E. Use of sonography for airway assessment: an observational study. *J Ultrasound Med* 2010; 29: 79-85.
2. Kristensen MS, Teoh WH, Graumann O, Laursen CB. Ultrasonography for clinical decision-making and intervention in airway management: from the mouth to the lungs and pleurae. *Insights Imaging* 2014; 5: 253-79.
3. Andruszkiewicz P, Wojtczak J, Sobczyk D, Stach O, Kowalik I. Effectiveness and validity of sonographic upper airway evaluation to predict difficult laryngoscopy. *J Ultrasound Med* 2016; 35: 2243-52.
4. Hui CM, Tsui BC. Sublingual ultrasound as an assessment method for predicting difficult intubation: a pilot study. *Anaesthesia* 2014; 69: 314-9.
5. Yadav NK, Rudingwa P, Mishra SK, Pannerselvam S. Ultrasound measurement of anterior neck soft tissue and tongue thickness to predict difficult laryngoscopy - An observational analytical study. *Indian J Anaesth* 2019; 63: 629-34.
6. Yao W, Wang B. Can tongue thickness measured by ultrasonography predict difficult tracheal intubation? *Br J Anaesth* 2017; 118: 601-9.
7. Apfelbaum JL, Hagberg CA, Caplan RA, Blitt CD, Connis RT, Nickinovich DG, et al. Practice guidelines for management of the difficult airway: an updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology* 2013; 118: 251-70.
8. Browner WS, Newman TB, Hulley SB. Estimating sample size and power: applications and Examples. In: *Designing Clinical Research An Epidemiologic Approach*. 4th ed. Edited by Hulley SB, Cummings SR, Browner WS, Grady DG, Newman TB: Philadelphia, Lippincott, Williams & Wilkins Publishers. 2013, p 73.
9. Crosby ET, Cooper RM, Douglas MJ, Doyle DJ, Hung OR, Labrecque P, et al. The unanticipated difficult airway with recommendations for management. *Can J Anaesth* 1998; 45: 757-76.
10. Adhikari S, Zeger W, Schmier C, Crum T, Craven A, Frrokaj I, et al. Pilot study to determine the utility of point-of-care ultrasound in the assessment of difficult laryngoscopy. *Acad Emerg Med* 2011; 18: 754-8.
11. Wu J, Dong J, Ding Y, Zheng J. Role of anterior neck soft tissue quantifications by ultrasound in predicting difficult laryngoscopy. *Med Sci Monit* 2014; 20: 2343-50.
12. Lundstrøm LH, Vester-Andersen M, Møller AM, Charuluxananan S, L'hermite J, Wetterslev J. Poor prognostic value of the modified Mallampati score: a meta-analysis involving 177 088 patients. *Br J Anaesth* 2011; 107: 659-67.
13. Crawley SM, Dalton AJ. Predicting the difficult airway. *BJA Educ* 2015; 15: 253-7.