

# Correction to: Hard- and soft-tissue profiles of the midface region in patients with skeletal Class III malocclusion using cone-beam computed tomography multiplanar-reconstructed image analysis

Bomi Kim<sup>a</sup>, Hyung-Chul Lee<sup>a</sup>, Seong-Hun Kim<sup>a</sup>, Yongil Kim<sup>a</sup>, Woosung Son<sup>a</sup>, Seong Sik Kim<sup>a,b,c</sup>

<sup>a</sup>Department of Orthodontics, Pusan National University Dental Hospital, Yangsan, Korea

<sup>b</sup>Dental Research Institute, Pusan National University Dental Hospital, Yangsan, Korea

<sup>c</sup>Institute of Translational Dental Sciences, Pusan National University, Busan, Korea

<https://doi.org/10.4041/kjod.2018.48.3.143>

In the above article, the 'Class' of skeletal malocclusion was misprinted as 'class' in the main text. The publisher would like to apologize for any inconvenience caused.

© 2018 The Korean Association of Orthodontists.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.