

Antioxidant Effect of CoQ₁₀ on N-nitrosodiethylamine-induced Oxidative Stress in Mice

Ho Sun Song¹, Hee Rae Kim¹, Tae Wook Park¹, Bong Jae Cho¹, Mi Young Choi¹, Chang Jong Kim¹, Uy Dong Sohn², and Sang Soo Sim¹

Departments of ¹Pathophysiology and ²Pharmacology, College of Pharmacy, Chung-Ang University, Seoul 156-756, Korea

The antioxidant effect of CoQ₁₀ on N-nitrosodiethylamine (NDEA)-induced oxidative stress was investigated in mice. Food intake and body weight were similar in both CoQ₁₀ and control groups during the 3-week experimental period. NDEA significantly increased the activities of typical marker enzymes of liver function (AST, ALT and ALP) both in control and CoQ₁₀ groups. However, the increase of plasma aminotransferase activity was significantly reduced in the CoQ₁₀ group. Lipid peroxidation in various tissues, such as heart, lung, liver, kidney, spleen and plasma, was significantly increased by NDEA, but this increase was significantly reduced by 100 mg/kg of CoQ₁₀. Superoxide dismutase activity increased significantly upon NDEA-induced oxidative stress in both the control and CoQ₁₀ groups with the effect being less in the CoQ₁₀ group. Catalase activity decreased significantly in both the control and CoQ₁₀ groups treated with NDEA, again with the effect being less in the CoQ₁₀ group. The lesser effect on superoxide dismutase and catalase in the NDEA-treated CoQ₁₀ group is indicative of the protective effect CoQ₁₀. Thus, CoQ₁₀ can offer useful protection against NDEA-induced oxidative stress.

Key Words: Catalase, CoQ₁₀, Lipid peroxidation, Reactive oxygen species, Superoxide dismutase

INTRODUCTION

Coenzyme Q is a quinone structure that is widely biosynthesized in living organisms such including yeasts, plants, and animals (Nohl et al., 1998). In humans, this compound has 10 isoprene units (5 carbons each) in the side chain and is thus named coenzyme Q₁₀ (CoQ₁₀, Ubiquinone-10, 2,3-dimethoxy, 5-methyl, 6-decaprenyl benzoquinone) (Crane, 2001). Two major physiological activities of CoQ₁₀ have been reported: (a) mitochondrial electron-transport activity and (b) antioxidant activity. CoQ₁₀H₂, a reduced form of CoQ₁₀, is an effective fat-soluble antioxidant and an important cellular antioxidant (Crane, 2001). CoQ₁₀H₂ inhibits lipid peroxidation when cell membranes and low-density lipoproteins are exposed to oxidizing conditions outside the body. In isolated mitochondria, CoQ₁₀ can protect membrane proteins and DNA from the oxidative damage that accompanies lipid peroxidation (Ernster and Dallner, 1995).

N-nitroso compounds, like N-nitrosodimethylamine, N-nitrosodiethylamine (NDEA) and N-nitrosopyrrolidine, constitute an important group of carcinogens and have been found in various foodstuffs, such as milk products, meat products, soft drinks and alcoholic beverages (Bansal et al., 2005). Nitroso compounds are readily formed in the human

body by the reaction of nitrite with amines and amides (Ohkawa et al., 1979). NDEA has been suggested to cause oxidative stress and cellular injury due to involvement of free radicals (Ames et al., 1993; Beckman and Ames, 1998; Noguchi et al., 2000). There are reports that oxygen free radicals and related lipid peroxides play a key role in the pathogenesis of age-related chronic degenerative diseases (Marklund and Marklund, 1974). It is very important to investigate the protective role of CoQ₁₀ on oxidative stress induced by NDEA, which may be formed in human body. Therefore, we measured the effect of CoQ₁₀ on lipid peroxidation, superoxide dismutase and catalase activity in various tissues in NDEA-treated mice.

METHODS

Materials

CoQ₁₀ powder was obtained from Daewoong Pharm. Co. Ltd. (Seoul, Korea). N-nitrosodiethylamine (NDEA), p-nitrophenylphosphate, thiobarbituric acid, hypoxanthine, nitroblue tetrazolium (NBT) and xanthine oxidase were purchased from Sigma Chemical Co. (St. Louis, MO, USA). 2',7'-Dichlorofluorescein diacetate (DCF-DA) was purchased from Molecular Probes (Leiden, Netherlands).

Received June 9, 2009, Revised July 15, 2009,
Accepted July 31, 2009

Corresponding to: Sang Soo Sim, Department of Pathophysiology, College of Pharmacy, Chung-Ang University, 221, Huksuk-dong, Dongjak-gu, Seoul 156-756, Korea. (Tel) 82-2-820-5615, (Fax) 82-2-816-7338, (E-mail) simss@cau.ac.kr

ABBREVIATIONS: ALP, alkaline phosphatase; ALT, alanine aminotransferase; AST, aspartate aminotransferase; CoQ₁₀, coenzyme Q₁₀; NDEA, N-nitrosodiethylamine; ROS, reactive oxygen species; SOD, superoxide dismutase.

Animals

Six-week-old male BALB/c mice, weighing 24 to 31 g, were purchased from Han-Lim Animal (Hwasung-gun, Gyeonggi-do, Korea) and were housed in an environmentally controlled animal room (temperature 24±2°C, humidity 50±5%, illumination 300~500 Lux).

Animal protocols were approved by the ethics committee for the care and use of laboratory animals at Chung-Ang University. The animals were randomized into two groups. CoQ₁₀ in olive oil (100 mg/kg) was orally administered to the CoQ₁₀ group (n=16) for 3 weeks, while an equal volume of olive oil was orally administered to the control group (n=16). After 3 weeks, oxidative stress was induced in half of the animals in each group by i.p. administration of NDEA (200 mg/kg body weight) in normal saline (0.5 ml) and the other half were administered an equal volume of normal saline.

Collection of samples

The mice were sacrificed by light ether anesthesia at 48 h after induction of oxidative stress, and blood was collected by cardiac puncture into test tubes rinsed with a saturated solution of EDTA. The organs including the liver, kidney, spleen, heart, and lung of each animal were removed, washed with normal saline, and weighed. The tissue was stored at -20°C until use.

Biochemical analysis

Plasma aspartate aminotransferase (AST) and alanine aminotransferase (ALT) activities were determined by the colorimetric method described by Reitman and Frenkel (1957). Alkaline phosphatase (ALP) activity was determined using p-nitrophenylphosphate as the substrate (King and Delory, 1959). The color that developed upon hydrolysis of p-nitrophenylphosphate to p-nitrophenol by ALP was read at 410 nm and was proportional to the enzyme activity.

Measurement of lipid peroxidation

The fluorescence of thiobarbituric acid reactive substances was used as a measure of lipid peroxidation (Boland et al., 2000). Tissues were homogenized in phosphate-buffered saline with a Polytron homogenizer. Trichloroacetic acid was added at a final concentration of 5%. Then, an equal volume of 0.325% 2-thiobarbituric acid in 50% glacial acetic acid was added. The solution was incubated at 95°C for 30 min. After cooling, the samples were vigorously mixed with 500 µl of isobutanol and centrifuged at 2,000 rpm for 10 min. The fluorescence of a 200 µl aliquot of the upper organic phase was quantified using excitation and emission wavelengths of 485 and 535 nm, respectively. Results were expressed as µg/mg protein. Protein concentration was measured by the bicinchoninic acid (BCA) method to correct for differences between preparations (Smith et al., 1987).

Superoxide dismutase (SOD) assay

The SOD level in each sample was measured spectrophotometrically via a hypoxanthine/xanthine oxidase generating system coupled with NBT reduction (Kirby and Schmidt,

1997). Reaction mixtures containing 50 mM potassium phosphate buffer (pH 7.4), 1 mM K-EDTA, 0.6 mM hypoxanthine, 0.2 mM NBT, 20 mU/ml xanthine oxidase and sample were incubated at 37°C for 20 min and the absorbance was measured at 590 nm (McCune and Johns, 2002).

Catalase assay

Catalase activity was directly determined by decomposition of H₂O₂ by recording the decrease in absorbance at 240 nm using UV spectrophotometer and was expressed as units per milligram of protein.

Measurement of reactive oxygen species (ROS) generation

RAW 264.7 cells obtained from the Korean Cell Line Bank (Seoul, Korea) were cultured in DMEM supplemented with 10% heat-inactivated FBS and antibiotic-antimycotic at 37°C with 5% CO₂. The level of intracellular ROS was quantified by fluorescence with DCF-DA. RAW 264.7 cells were suspended in 20 ml of Krebs buffer [137 mM NaCl, 2.7 mM KCl, 0.4 mM Na₂HPO₄, 0.5 mM MgCl₂, 10 mM HEPES (pH 7.4), 1.8 mM CaCl₂, 5 mM glucose] and incubated with 20 µM DCF-DA for 30 min at 37°C. The cells were washed twice with Krebs buffer and were suspended in Krebs buffer at a density of 10⁶ cells/ml. The cells were incubated with CoQ₁₀ for 10 min, and ROS generation was then induced with 1 mg/ml silica for 30 min at 37°C. The fluorescence intensity was measured using excitation and emission wavelengths of 485 and 535 nm, respectively (Woo et al., 2000).

Data analysis

The results were represented as means±SD and analyzed statistically by analysis of variance (ANOVA), and the differences between groups were determined with the Newman-Keuls test. The level of significance was set at less than 5%.

RESULTS

Effect of CoQ₁₀ on NDEA-induced liver damage

Food intake and body weight were similar in both the control and CoQ₁₀ groups during the 3-week experimental period. The body weights in the control and CoQ₁₀ groups

Table 1. Protective effect of CoQ₁₀ on NDEA (200 mg/kg)-induced liver damage in mice

Parameters	Control		CoQ ₁₀ (100 mg/kg)	
	Saline	NDEA	Saline	NDEA
AST (U/l)	15.0±3.2	58.3±6.8*	14.2±2.5	42.3±4.5* [#]
ALT (U/l)	14.9±2.5	69.4±5.3*	12.8±3.3	57.3±6.3* [#]
ALP (U/l)	48.5±6.0	83.3±7.2*	47.2±7.0	71.3±5.3* [#]

Values are mean±SD; n=8. AST, aspartate aminotransferase; ALT, alanine aminotransferase; ALP, alkaline phosphatase. *significantly different from saline group (p<0.05), [#]significantly different from NDEA control group (p<0.05).

increased from 25.8 ± 1.1 to 27.8 ± 1.0 and from 26.3 ± 1.3 g to 27.9 ± 1.3 g, respectively. The activities of typical marker enzymes of liver function (AST, ALT and ALP) were also similar in both groups without NDEA (Table 1). NDEA significantly increased these activities both in the control and CoQ₁₀ groups. However, the increase of plasma amino-transferase activity was significantly less in the CoQ₁₀ group as compared with the control group.

Effect of CoQ₁₀ on NDEA-induced lipid peroxidation

Lipid peroxidation induced by NDEA in various tissues, including heart, lung, liver, kidney, spleen and plasma, was significantly increased both in the control and CoQ₁₀ groups. Lipid peroxidation was the most prominent in the heart. CoQ₁₀ alone did not affect the lipid peroxidation level without NDEA treatment, whereas it significantly reduced the level of NDEA-induced lipid peroxidation in all tissues, including plasma, heart, lung, liver, kidney and spleen (Fig. 1).

Effect of CoQ₁₀ on SOD activity in murine tissues treated with NDEA

In all tissues, SOD activity increased significantly upon induction of oxidative stress by NDEA both in the control and CoQ₁₀ groups. In the control, the increase of SOD activity induced by NDEA in plasma, heart and lung was more prominent than that in liver, kidney and spleen. The increase of SOD levels by NDEA in all tissues was significantly reduced by treatment with 100 mg/kg CoQ₁₀ (Fig. 2). In the normal group without NDEA, CoQ₁₀ alone significantly increased the SOD level in plasma, whereas it did not affect SOD levels in other tissues.

Effect of CoQ₁₀ on catalase activity in murine tissues treated with NDEA

Upon induction of oxidative stress by NDEA, catalase activity decreased significantly in the control and CoQ₁₀




Fig. 1. The effect of CoQ₁₀ on NDEA-induced lipid peroxidation in murine tissues. Malondialdehyde (MDA) is an index of lipid peroxidation in oxidative stress induced by 200 mg/kg NDEA. CoQ₁₀ in olive oil (100 mg/kg) was orally administered to the CoQ₁₀ group for 3 weeks, while an equal volume of olive oil was orally administered to the control group. Results are means \pm SD from 8 mice. *significantly different from saline group ($p < 0.05$), #significantly different from NDEA control group ($p < 0.05$).




Fig. 2. The effect of CoQ₁₀ on superoxide dismutase (SOD) activity in murine tissues treated with 200 mg/kg NDEA. CoQ₁₀ in olive oil (100 mg/kg) was orally administered to the CoQ₁₀ group for 3 weeks, while an equal volume of olive oil was orally administered to the control group. Results are means±SD from 8 mice. *significantly different from saline group ($p < 0.05$), #significantly different from NDEA control group ($p < 0.05$).

groups. CoQ₁₀ significantly restored the level of catalase in all tissues treated with NDEA (Fig. 3). CoQ₁₀ alone significantly increased the levels of catalase in heart and lung, but it did not affect the levels in liver, kidney and spleen. In a cell culture system, CoQ₁₀ dose-dependently reduced ROS generation in RAW 264.7 cells stimulated with 1 mg/ml silica (Fig. 4).

DISCUSSION

The food intake and changes in body weight in the control and CoQ₁₀ groups during the 3-week experimental period were similar, which indicate that CoQ₁₀ does not produce any untoward effects. Liver is the typical detoxifying organ of the body and is profoundly affected by external chemicals. Aminotransferases (AST and ALT) and ALP have been used as markers of liver function. The increase in these activities by NDEA seems to be due to the hepatotoxicity of the chemical. It has been previously reported that hepatotoxic

responses to NDEA result in increased levels of AST, ALT, and ketone bodies with nitroso compounds (Taniguchi et al., 1999). A relatively lower degree of hepatic damage in the CoQ₁₀ group is indicative of the hepatoprotective nature of CoQ₁₀. While there is no direct evidence of the hepatoprotective effect of CoQ₁₀, it has been reported that maintenance of the normal level of CoQ₉H₂ in the S-allyl-mercaptocysteine pretreatment group may suppress acetaminophen-induced liver injury (Sumioka et al., 1998).

Our data showed that NDEA profoundly increased the level of lipid peroxidation. Free radicals, mostly ROS, cause cellular injury, the consequences of which are often exhibited and measured as lipid peroxidation (Spiteller, 1996). Lipid peroxidation is the oxidative deterioration of polyunsaturated lipids to form radical intermediates that cause oxidative damage. Malondialdehyde, the major end-product of this reaction, is an index of lipid peroxidation and can be estimated as thiobarbituric acid reactive substances (Kohen and Nyska, 2002). Tremendous increases in malondialdehyde upon NDEA-induced oxidative stress




Fig. 3. The effect of CoQ₁₀ on catalase activity in murine tissues treated with 200 mg/kg NDEA. CoQ₁₀ in olive oil (100 mg/kg) was orally administered to the CoQ₁₀ group for 3 weeks, while an equal volume of olive oil was orally administered to the control group. Results are means \pm SD from 8 mice. *significantly different from each saline group ($p < 0.05$), #significantly different from NDEA group of control ($p < 0.05$).




Fig. 4. The effect of CoQ₁₀ on intracellular reactive oxygen species (ROS) production induced by 1 mg/ml silica in RAW 264.7 cells. Results are means \pm SD from 4 separate experiments.

could be caused by increased production of or decreased destruction of ROS. The lesser effect of CoQ₁₀ on NDEA-induced lipid peroxidation in various tissues may be due to

the presence of an antioxidant system associated with CoQ₁₀ that increases anti-oxidogenic potential. Supplementation of CoQ₁₀ in diet is reported to help maintain a favorable antioxidant/pro-oxidant balance (Sohal and Forster, 2007). Recently, it has been also reported that CoQ₁₀ significantly decreases the level of lipid peroxidation *in vivo* and *in vitro* (Sawicka and Dlugosz, 2008; Sena et al., 2008). This effect of CoQ₁₀ may be related to its antioxidant activity since it significantly decreased ROS generation in silica-stimulated RAW 264.7 cells.

SOD scavenges the superoxide radical by converting it to H₂O₂, which in turn is converted to molecular oxygen by reactions catalyzed by peroxidase and catalase (Thomas, 2003). Catalase is ubiquitously present in all aerobic cells and is a major component of the primary antioxidant defense system. It is most abundant in liver and is responsible for catalytic decomposition of H₂O₂ to oxygen and water. In our studies, catalase activity decreased significantly upon NDEA-induced oxidative stress in control and CoQ₁₀ groups, the effect being less in the CoQ₁₀ group. Decreased

catalase activity indicates that NDEA-induced oxidative stress interferes with the anti-oxygenic potential in the tissues, thus resulting in increased generation of ROS. The H_2O_2 formed by SOD is decomposed by catalase. The decrease in catalase activity in animals administered NDEA may be caused by inactivation of catalase as superoxide anions have been shown to reduce catalase activity (Bartsch et al., 1989; Bansal et al., 1996). Similarly, exposure of erythrocytes to NDEA inhibited catalase activity through excess superoxide anion formation (Bansal et al., 2005). The lesser effect on catalase in the CoQ₁₀ group during NDEA-induced oxidative stress is indicative of the protective effect of CoQ₁₀. Thus, CoQ₁₀ can offer useful protection against NDEA-induced oxidative stress.

REFERENCES

- Ames BN, Shigenga MK, Hagen TM. Oxidants, antioxidants and the degenerative diseases of aging. *Proc Natl Acad Sci* 90: 7915–7922, 1993.
- Bansal AK, Bansal M, Soni G, Bhatnagar D. Modulation of NDEA induced oxidative stress by vitamin E in rat erythrocytes. *Hum Exp Toxicol* 24: 297–302, 2005.
- Bansal AK, Bhatnagar D, Soni GL. *In vitro* effect of N-nitrosodimethylamine on lipid peroxidation and antioxidant system in human erythrocytes. *Toxicol In Vitro* 10: 649–653, 1996.
- Bartsch H, Hietanen E, Malaveille C. Carcinogenic nitrosamines: free radical aspects of their action. *Free Radic Biol Med* 7: 637–644, 1989.
- Beckman KB, Ames BN. The free radical theory of aging matures. *Physiol Rev* 78: 547–581, 1998.
- Boland A, Delapierre D, Mossay D, Hans P, Dresse A. Propofol protects cultured brain cells from iron ion-induced death: comparison with trolox. *Eur J Pharmacol* 404: 21–27, 2000.
- Crane FL. Biochemical functions of coenzyme Q10. *J Am Coll Nutr* 20: 591–598, 2001.
- Ernster L, Dallner G. Biochemical, physiological and medical aspects of ubiquinone function. *Biochim Biophys Acta* 1271: 195–204, 1995.
- King EJ, Delory GE. The rates of enzyme hydrolysis of phosphoric esters. *Biochem J* 33: 11–85, 1959.
- Kirby AJ, Schmidt RJ. The antioxidant activity of Chinese herbs for eczema and of placebo herbs-I. *J Ethnopharmacol* 56: 103–108, 1997.
- Kohen R, Nyska A. Oxidation of biological systems: oxidative stress phenomena, antioxidants, redox reactions and methods for their quantification. *Toxicol Pathol* 30: 620–650, 2002.
- Marklund S, Marklund G. Involvement of the superoxide anion radical in the auto-oxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur J Biochem* 47: 469–474, 1974.
- McCune LM, Johns T. Antioxidant activity in medicinal plants associated with the symptoms of diabetes mellitus used by the indigenous peoples of the north American boreal forest. *J Ethnopharmacol* 82: 197–205, 2002.
- Noguchi N, Watanabe A, Shi H. Diverse functions of antioxidants. *Free Radic Res* 33: 809–817, 2000.
- Nohl H, Gille L, Staniek K. The biochemical, pathophysiological, and medical aspects of ubiquinone function. *Ann NY Acad Sci* 854: 394–409, 1998.
- Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric reaction. *Anal Biochem* 95: 351–358, 1979.
- Reitman S, Frenkel SA. Colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvate transaminases. *Am J Pathol* 28: 56–63, 1957.
- Sawicka E, Dlugosz A. Toluene and P-xylene mixture exerts antagonistic effect on lipid peroxidation *in vitro*. *Int J Occup Med Environ Health* 21: 201–209, 2008.
- Sena CM, Nunes E, Gomes A, Santos MS, Proença T, Martins MI, Seça RM. Supplementation of coenzyme Q10 and alpha-tocopherol lowers glycated hemoglobin level and lipid peroxidation in pancreas of diabetic rats. *Nutr Res* 28: 113–121, 2008.
- Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH, Provenzano MD, Fujimoto EK, Goeke NM, Olson BJ, Klenk DC. Measurement of protein using bicinchoninic acid. *Anal Biochem* 163: 279–282, 1987.
- Sohal RS, Forster MJ. Coenzyme Q, oxidative stress and aging. *Mitochondrion* 7S: S103–S111, 2007.
- Spiteller G. Enzymic lipid peroxidation - a consequence of cell injury? *Free Radic Biol Med* 21: 1003–1009, 1996.
- Sumioka I, Matsura T, Kasuga S, Itakura Y, Yamada K. Mechanisms of protection by S-allylmercaptocysteine against acetaminophen-induced liver injury in mice. *Jpn J Pharmacol* 78: 199–207, 1998.
- Taniguchi M, Yasutake A, Takedomi K, Inoue K. Effects of N-nitrosodimethylamine (NDMA) on the oxidative status of rat liver. *Arch Toxicol* 73: 141–146, 1999.
- Thomas WS. The physiology and pharmacology of singlet oxygen. *Med Hypothesis* 60: 567–572, 2003.
- Woo CH, Eom YW, Yoo MH, You HJ, Han HJ, Song WK, Yoo YJ, Chun JS, Kim JH. Tumor necrosis factor-alpha generates reactive oxygen species via a cytosolic phospholipase A2-linked cascade. *J Biol Chem* 275: 32357–32362, 2000.