

Effects of Erythropoietin on Memory Deficits and Brain Oxidative Stress in the Mouse Models of Dementia

Rohit Kumar, Amteshwar Singh Jaggi, and Nirmal Singh

Department of Pharmaceutical Sciences and Drug Research, Punjabi University, Patiala (Punjab), PIN-147002, India

The present study was undertaken to explore the potential of erythropoietin in memory deficits of mice. Memory impairment was produced by scopolamine (0.5 mg/kg, *i.p.*) and intracerebroventricular streptozotocin (*i.c.v.* STZ, 3 mg/kg, 10 μ l, 1st and 3rd day) in separate groups of animals. Morris water-maze test was employed to assess learning and memory. The levels of brain thio-barbituric acid reactive species (TBARS) and reduced glutathione (GSH) were estimated to assess degree of oxidative stress. Brain acetylcholinesterase enzyme (AChE) activity was also measured. Scopolamine/streptozotocin administration induced significant impairment of learning and memory in mice as indicated by marked decrease in Morris water-maze performance. Scopolamine/streptozotocin administration also produced a significant enhancement of brain AChE activity and brain oxidative stress (an increase in TBARS and a decrease in GSH) levels. Treatment of erythropoietin (500 and 1,000 IU/Kg *i.p.*) significantly reversed scopolamine- as well as streptozotocin-induced learning and memory deficits along with attenuation of those-induced rise in brain AChE activity and brain oxidative stress levels. It may be concluded that erythropoietin exerts a beneficial effect in memory deficits of mice possibly through its multiple actions including potential anti-oxidative effect.

Key Words: Erythropoietin, Memory, Scopolamine, Streptozotocin, Morris water-maze

INTRODUCTION

The human brain is a very complex and intricate machine and many factors can interfere with its functioning. With age, time and medical conditions such as atherosclerosis, stroke, high levels of cholesterol, plasma homocysteine, diabetes and genetic factors a person may lose his ability to solve problems and maintain emotional control, therefore experience personality changes and behavioral problems such as agitation, delusions, hallucinations and cognitive deficits. Such a mental condition characterized by impairment of memory and loss of intellectual ability, sufficiently severe to interfere with one's occupational or social activities is termed as dementia (Sharma et al, 2008a; 2008b). Alzheimer's disease (AD) is the most common serious type of dementia effecting elderly population. AD is a progressive neurodegenerative disorder associated with loss of neurons in distinct brain areas. Currently, over 50% cases of memory impairment account for dementia of AD (Torre et al, 2004). Clinical management of memory dysfunction associated with dementia is still a nightmare for the neurobiologist because of only limited therapeutic modalities. Currently only two categories of the drugs are available in the market *i.e.* cholinesterase inhibitors (such as donepezil,

rivastigmine and galantamine) and N-methyl-D-aspartate (NMDA) receptor antagonist (memantine). Scientists are looking for an agent who will not only reverse memory dysfunction but also progression of dementia.

Erythropoietin (EPO) is basically a glycoprotein that increases red cell mass to improve tissue oxygenation. It is produced by the kidney in response to hypoxia. Recombinant human EPO (r-Hu-EPO) is effective and widely used in treatment of anemia associated with renal failure, HIV infection, cancer and surgery. Like other members of the cytokine family to which it and its receptor belong, both are expressed within the CNS in response to hypoxia (Masuda et al, 1994; Marti et al, 1996; Juul et al, 1999). Hypoxia may not be the only relevant stimulus for brain EPO production, however, as metabolic disturbances, including hypoglycemia and strong neuronal depolarization, generate mitochondrial reactive oxygen species that may increase brain EPO expression through hypoxia inducible factor-1 (Chandel et al, 1998). EPO may thus protect nervous tissue under any condition characterized by a relative deficiency of ATP in the face of increased metabolic demands. EPO has been shown to exhibit classic neurotrophic effects *in vivo* and *in vitro* (Konishi et al, 1993;

Received May 11, 2010, Revised August 20, 2010,
Accepted August 24, 2010

Corresponding to: Nirmal Singh, Department of Pharmaceutical Sciences and Drug Research, Punjabi University, Patiala-147002, Punjab, India. (Tel) 91-(0175)-3046255, (E-mail) nirmal_puru@rediffmail.com

ABBREVIATIONS: STZ, streptozotocin; *i.c.v.*, intracerebroventricularly or intracerebroventricular; EPO, erythropoietin; ACSF, artificial cerebrospinal fluid; AChE, acetylcholinesterase; TBARS, thiobarbituric acid reactive species; GSH, glutathione; AD, Alzheimer's disease; HIV, human immunodeficiency virus; NMDA, N-methyl-D-aspartate; ATP, adenosine triphosphate; MWM, Morris water-maze; ELT, escape latency time; TSTQ, time spent in target quadrant; AChE, acetyl cholinesterase; LTP, long term-potential.

Tabira et al, 1995; Campana et al, 1998; Juul et al, 1999; Brines et al, 2000). In the light of above, the present study has been undertaken to investigate the effects of EPO on memory deficits and brain oxidative stress in scopolamine- and streptozotocin (STZ)-induced dementia mouse models (Lannert and Hoyer, 1998; Sharma and Gupta 2001; Parle and Singh, 2004).

METHODS

Animals

Swiss albino mice, weighing 20~30 g were employed in the present study (procured from Central Research Institute, Kasauli). They were maintained on standard laboratory pellet chow diet (Kisan Feeds Ltd, Chandigarh, India) and water ad libitum. The animals were exposed to natural cycles of light and dark. The mice were acclimatized to the laboratory conditions five days prior to behavioral study. The experimental protocol was duly approved by Institutional animal ethics committee and care of the animals was carried out as per the guidelines of committee for the purpose of control and supervision of experiments on animals (CPCSEA), Ministry of Environment and Forest Government of India (Reg. No. CPCSEA/107/1999).

Drugs and reagents

Erythropoietin was purchased from Claris (India) Pvt. Ltd. Folin-Ciocalteu's Phenol reagent was purchased from Merck Limited, Mumbai, India. 5,5, dithiobis (2-nitro benzoic acid) (DTNB), reduced glutathione (GSH), Bovine serum albumin (BSA) and Thiobarbituric acid were obtained from Loba Chem, Mumbai, India. Streptozotocin (STZ) and scopolamine were purchased from Sigma-Aldrich, St. Louis, U.S.A. All the reagents used in this study were of analytical grade. STZ was dissolved in artificial cerebro spinal fluid (ACSF) prepared according to the method as described by Sakurada et al, (1999). EPO and scopolamine were dissolved in distilled water and given intraperitoneally (i.p.). STZ and ACSF were delivered intracerebroventricularly (i.c.v.).

Intracerebroventricular administration of STZ

Mice were anesthetised with anaesthetic ether for intracerebroventricular (i.c.v.) administrations. Ether has been preferred here due to its ultrashort action and fast reversibility. Moreover brief extent of ether exposure for i.c.v. injection has been reported to exert no significant effect on learning and memory behavior of animals (Haley and McCormick, 1957). Intracerebroventricular injections were made with hypodermic needle of 0.4 mm external diameter attached to a 10 μ l Hamilton microlitre syringe (Top Syringe, Mumbai, India). The needle was covered with a polypropylene tube except 3 mm of the tip region so as to insert this much portion of the needle perpendicularly through the skull into the brain of mouse. STZ was dissolved in artificial CSF (25 mg/ml) solution, made freshly. The injection site was 1 mm to right or left midpoint on the line drawn through to the anterior base of the ears. Injections were performed into right or left ventricle randomly. Two doses of STZ (3 mg/kg) were administered by i.c.v. injection bilaterally. The second dose was administered after 48 h of first dose. The concentration was ad-

justed so as to deliver 10 μ l in an injection. Control group mice were given i.c.v. injection of artificial cerebrospinal fluid (CSF) in a similar fashion.

Intra peritoneal (i.p.) administration of scopolamine

Mice were administered scopolamine (0.5 mg/kg, i.p.) dissolved in distilled water (10 ml/kg, i.p.) daily for 4 days, 30 min prior to acquisition trials on Morris water-maze.

Morris water maze (MWM)

Morris water-maze test was employed to assess learning and memory of the animals (Parle and Singh, 2004; Morris, 1984). MWM is a swimming based model where the animal learns to escape on to a hidden platform. It consisted of large circular pool (150 cm in diameter, 45 cm in height, filled to a depth of 30 cm with water maintained at 28 \pm 1°C). The water was made opaque with white colored non-toxic dye. The tank was divided into four equal quadrants with the help of two threads, fixed at right angle to each other on the rim of the pool. A submerged platform (10 \times 10 cm), painted in white was placed inside the target quadrants of this pool, 1 cm below surface of water. The position of platform was kept unaltered throughout the training session. Each animal was subjected to four consecutive training trials on each day with inter-trial gap of 5 min. The mouse was gently placed in the water between quadrants, facing the wall of pool with drop location changing for each trial, and allowed 120 sec to locate submerged platform. Then, it was allowed to stay on the platform for 20 sec. If it failed to find the platform within 120 sec, it was guided gently onto platform and allowed to remain there for 20 sec. Day 4 escape latency time (ELT) to locate the hidden platform in water maze was noted as an index of acquisition or learning. Animal was subjected to training trials for four consecutive days, the starting poison was changed with each exposure as mentioned below and target quadrant (Q4 in the present study) remained constant throughout the training period.

Day1	Q1	Q2	Q3	Q4
Day2	Q2	Q3	Q4	Q1
Day3	Q3	Q4	Q1	Q2
Day4	Q4	Q1	Q2	Q3

On fifth day, platform was removed and each mouse was allowed to explore the pool for 120 sec. Mean time spent in all four quadrants was noted. The mean time spent by the animal in target quadrant searching for the hidden platform was noted as index of retrieval or memory.

The experimenter always stood at the same position. Care was taken that relative location of water maze with respect to other objects in the laboratory serving, as prominent visual clues were not disturbed during the total duration of study. All the trials were completed between 09.00 to 17.00 h.

Collection of samples

Animals were sacrificed by cervical dislocation; brains were removed and homogenized in phosphate buffer (pH=7.4). The homogenates were then centrifuged at 3,000 rpm for 15 min. The supernatant of homogenate was used for biochemical estimation as per the following methods.

Estimation of brain acetyl cholinesterase (AChE) activity

The whole brain AChE activity was measured by the method of Ellman, et al, (1961) and Voss and Sachsse, (1970). Change in absorbance per min of the sample was read spectrophotometrically at 420 nm.

Estimation of brain thiobarbituric acid reactive species (TBARS) level

The whole brain TBARS level was measured by the method of Ohkawa et al, (1979). The absorbance was measured spectrophotometrically (DU 640B spectrophotometer, Beckman Coulter Inc., CA, USA) at 532 nm.

Estimation of brain reduced glutathione (GSH) level

The whole brain GSH level was measured by the method of Beutler et al, (1963). The absorbance was measured spectrophotometrically at 532 nm.

Estimation of brain total protein

Total brain protein was estimated using method of Lowry et al, (1951). The absorption was read spectrophotometrically at 750 nm.

Experimental protocol

In total 70 mice were employed in the study. The animals were divided into 10 groups, each comprising 07 mice.

Group I (Control)

Mice were administered distilled water (10 ml/kg, i.p.) 30 min before acquisition trials conducted from day 1 to day 4 and 30 min before retrieval trial conducted on day 5.

Group II (Artificial cerebrospinal fluid {ACSF} control)

Mice were injected intracerebroventricularly artificial cerebrospinal fluid (i.c.v., ACSF, 25 mg/ml, 10 μ l) in two dosage schedules i.e. on first and third day followed by exposure to Morris Water Maze (MWM) test after 15 days.

Group III (scopolamine control)

Mice were administered scopolamine (0.5 mg/kg, i.p.) 30 min before acquisition trial conducted from day 1 to day 4 and then vehicle (distilled water) only 30 min prior to retrieval trial conducted on day 5.

Group IV (streptozotocin control)

Mice were injected intracerebroventricularly streptozotocin (STZ i.c.v., 3 mg/kg, 10 μ l) in two dosage schedules i.e. on first and third day followed by exposure to MWM test after 15 days.

Group V (EPO per se- Low dose)

Mice were administered EPO (500 IU/Kg, i.p.) 30 min before acquisition trial conducted from day 1 to day 4. The animals were administered vehicle (distilled water) only 30 min before retrieval trial conducted on day 5.

Group VI (EPO per se- High dose)

Mice were administered EPO (1,000 IU/Kg, i.p.) 30 min before acquisition trial conducted from day 1 to day 4. The

animals were administered vehicle (distilled water) only 30 min before retrieval trial conducted on day 5.

Group VII (EPO {Low dose}+scopolamine)

Mice were treated with EPO (500 IU/kg, i.p.) and then after 30 min with scopolamine (0.5 mg/kg, i.p.) After 30 min of scopolamine treatment these animals were then subjected to acquisition trials on MWM. The treatment of EPO+scopolamine was continued during acquisition trials from day 1 to day 4. The animals were administered vehicle (distilled water) only 30 min before retrieval trial conducted on day 5.

Group VIII (EPO {High dose}+scopolamine)

Mice were treated with EPO (1,000 IU/kg, i.p.) and then after 30 min with scopolamine (0.5 mg/kg, i.p.) After 30 min of scopolamine treatment these animals were then subjected to acquisition trials on Morris water maze. The treatment of EPO+scopolamine was continued during acquisition trials from day 1 to day 4. The animals were administered vehicle (distilled water) only 30 min before retrieval trial conducted on day 5.

Group IX (STZ+EPO {Low dose})

STZ i.c.v treated mice, were administered EPO (500 IU/kg, i.p.) starting after 2nd dose of STZ, daily for 15 days and then subjected to Morris water maze test. The administration of EPO (administered 30 min before) was also continued during acquisition trials conducted from day 1 to day 4. On day 5 animals were administered vehicle (distilled water) only, and then subjected to retrieval test after 30 min.

Group X (STZ+EPO {High dose})

STZ i.c.v treated mice, were administered EPO (1,000 IU/kg, i.p.) starting after 2nd dose of STZ, daily for 15 days and then subjected to Morris water maze test. The administration of EPO (administered 30 min before) was also continued during acquisition trials conducted from day 1 to day 4. On day 5 animals were administered vehicle (distilled water) only, and then subjected to retrieval test after 30 min.

Statistical analysis

All the results were expressed as mean \pm standard error of mean (S.E.M). The data of ELT and time spent in quadrants in various groups were statistically analyzed using two-way ANOVA followed by Bonferonni test and rests of the data was analyzed using one-way ANOVA followed by Tukey's multiple range test. $p < 0.05$ was considered to be statistically significant. Sigma Stat Statistical software version 3.5 was used for data processing.

RESULTS

Various pharmacological interventions employed in the present study did not show any significant mortality. Further, no significant difference was observed between the results obtained from mice of either sex.

Effect on escape latency time (ELT) and time spent in target quadrant (TSTQ) using Morris water maze

Control animals, showed a significant decrease in their

Table 1. Effect of erythropoietin on scopolamine/streptozotocin-induced changes in escape latency time (ELT) using Morris water-maze

Sr. No.	Group	Dose	Day 1 ELT (sec)	Day 4 ELT (sec)
I	Control	10 ml/kg, i.p.	95.5±4.11	38.5±1.04 ^a
II	ACSF control	25 mg/ml, 10 μ l, i.c.v	97.2±3.64	40.2±1.40 ^a
III	scopolamine	0.5 mg/kg, i.p.	92.8±1.01	66±0.69 ^b
IV	EPO-L	500 IU/kg, i.p.	94.1±1.5	39.4±0.97 ^a
V	EPO-H	1000 IU/kg, i.p.	95.1±1.47	41.1±1.4 ^a
VI	EPO-L+scopolamine	500 IU/kg, i.p.+0.5 mg/kg, i.p.	95.4±1.96	45.1±0.76 ^c
VII	EPO-H+scopolamine	1000 IU/kg i.p.+0.5 mg/kg, i.p.	97.7±1.04	40±1 ^c
VIII	STZ	3 mg/kg, 10 μ l, i.c.v	99.1±1.10	75.5±1.41 ^d
IX	STZ+EPO-L	3 mg/kg, 10 μ l, i.c.v+500 IU/kg, i.p.	96.5±1.67	49.1±1.03 ^e
X	STZ+EPO-H	3 mg/kg, 10 μ l, i.c.v+1,000 IU/kg, i.p.	97.5±0.78	42.5±0.89 ^e

ACSF-C: artificial cerebrospinal fluid-control; EPO-L and H: erythropoietin (low and high); STZ: streptozotocin; ELT: escape latency time. Each group (n=7) represents mean±standard errors of means. Two-way ANOVA followed by Bonferonni's post hoc test. F (3, 24)=11.100, $p < 0.001$ for evaluating the effect of days and F (9, 60)=50.740, $p < 0.001$ for evaluating the effect of treatment on ELT, ^a $p < 0.05$ Vs Day 1 ELT in control group, ^b $p < 0.05$ Vs Day 4 ELT in control, ^c $p < 0.05$ Vs Day 4 ELT in scopolamine group, ^d $p < 0.05$ Vs Day 4 ELT in ACSF group, ^e $p < 0.05$ Vs Day 4 ELT in STZ group.




Fig. 1. Effect of erythropoietin on scopolamine and streptozotocin induced memory impairments using Morris water-maze. ACSF-C: artificial cerebrospinal fluid-control; SCO-C: scopolamine-control; EPO-L and H: erythropoietin (low and high); STZ: streptozotocin. Each group (n=7) represents mean±standard errors of means. Two-way ANOVA followed by Bonferonni's post hoc test. F (3, 24)=6.455, $p < 0.0001$ for evaluating the difference in time spent in various quadrants; F (9, 60)=26.231, $p < 0.001$ for evaluating the effect of treatment on difference in time spent in target quadrant. ^a $p < 0.05$ Vs time spent in other quadrants in control group, ^b $p < 0.05$ Vs time spent in Target Quadrant (TSTQ) of control, ^c $p < 0.05$ Vs TSTQ of scopolamine group, ^d $p < 0.05$ Vs TSTQ of ACSF group, ^e $p < 0.05$ Vs TSTQ of STZ group.

day 4 ELT as compared to its value noted on day 1, reflecting acquisition (learning). Further these mice significantly spent more time in the target quadrant (Q4) in search of missing platform as compared to the time spent in other quadrants (Q1, Q2, Q3) during retrieval trial conducted on day 5, indicating memory or retrieval. Treatment of vehicle artificial cerebrospinal fluid (ACSF) did not show any significant effect on day 4 ELT (Table 1) and day 5 mean time spent in target quadrant (TSTQ) of control animals (Fig. 1). Administration of scopolamine (0.5 mg/kg, i.p.)/streptozotocin (3 mg/kg, 10 μ l, i.c.v) significantly prevented decrease in day 4 ELT of respective control group (Table 1) and markedly reduced time spent time in target quadrant (Q4) in search of missing platform during retrieval trial reflect-




Fig. 2. Effect of erythropoietin on scopolamine/streptozotocin-induced changes in brain AChE activity. ACSF-C: artificial cerebrospinal fluid-control; SCO-C: scopolamine-control; EPO-L and H: erythropoietin (low and high); STZ: streptozotocin. Each group (n=7) represents mean±standard errors of means. One-way ANOVA followed by Tukey's multiple range test; F (9, 60)=15.851, $p < 0.001$. ^a $p < 0.05$ Vs control group, ^b $p < 0.05$ Vs scopolamine, ^c $p < 0.05$ Vs ACSF control, ^d $p < 0.05$ Vs STZ.

ing impairment of both learning as well as memory (Fig. 1).

Effect of erythropoietin on scopolamine/streptozotocin-induced impairment of learning and memory using Morris water maze

Erythropoietin (500 IU/kg i.p.; 1,000 IU/kg, i.p.) *per se* did not produce any significant effect on day 4 decrease in ELT (Table 1) and day 5 increase in TSTQ of control mice (Fig. 1). However administration of EPO to scopolamine/streptozotocin treated animals, significantly attenuated day 4 rise in the ELT (Table 1) as well as day 5, decrease in TSTQ, indicating reversal of scopolamine/streptozotocin-induced memory deficits (Fig. 1).

Effect of erythropoietin on scopolamine/streptozotocin-induced changes in acetylcholinesterase (AChE) activity of brain

Scopolamine (0.5 mg/kg, i.p.)/streptozotocin (3 mg/kg,

10 μ l, i.c.v) significantly, increased the brain AChE activity when compared to control mice. EPO (500 and 1,000 IU/kg, i.p.) did not produce any significant *per se* effect on the brain AChE level as compared to control mice. Treatment of EPO significantly attenuated scopolamine as well as streptozotocin-induced rise in brain AChE activity (Fig. 2)

Effect of erythropoietin on scopolamine/streptozotocin-induced changes in oxidative stress levels of brain

Scopolamine (0.5 mg/kg, i.p.)/streptozotocin (3 mg/kg, 10 μ l, i.c.v) significantly, increased the brain thio barbituric acid reactive species (TBARS) level and reduced the brain GSH levels, compared to control group of animals, reflecting enhanced oxidative stress. EPO (500 and 1,000 IU/kg, i.p.) did not produce any significant *per se* effect on brain TBARS and GSH levels of control group animals. Treatment of EPO (500 and 1,000 IU/kg, i.p.) significantly reversed scopolamine as well as streptozotocin-induced rise in brain oxidative stress as reflected by a rise in brain TBARS levels and a fall in GSH levels (Fig. 3 and 4).

DISCUSSION

Morris Water Maze test employed in present study is one of the most widely accepted models to evaluate learning and memory of the animals (Morris, 1984; Parle and Singh, 2004). A significant decrease in day 4 escape latency time (ELT) of control animals during ongoing acquisition trials denoted normal acquisition of memory and an increase in time spent in target quadrant (TSTQ), in search of missing platform during retrieval trial indicated, retrieval of memory. These results are consistent to our earlier findings (Parle and Singh, 2007; Sharma et al, 2008a; 2008b) and reports from other laboratory (Packard et al, 1996).

In the present study scopolamine produced impairment of acquisition and retrieval of memory as reflected by significant increase in day 4 ELT and decrease in day 5 TSTQ respectively. Acetylcholine (ACh) is a classic mediator of learning and memory (Blokland, 1995). Drugs that reduce

cholinergic function such as muscarinic receptor antagonist scopolamine, cause profound memory impairments in animals and humans (Deutsch and Rocklin, 1967; Deutsch, 1971). The degeneration and dysfunction of cortical cholinergic neurons is closely associated with cognitive deficits of AD (Bartus et al, 1982; Coyle et al, 1983). These findings provided a cholinomimetic rationale for treatment of dementia closely related to AD (Becker, 1991) and support the use of animal models using muscarinic receptor antagonists (Yamazaki et al, 1995). This contention is further supported by our study, whereby a significant impairment of learning and memory in mice treated with scopolamine has been observed. Moreover, scopolamine treatment also produced a significant enhancement of brain acetylcholinesterase (AChE) activity and increase in oxidative stress as indicated by rise in brain thioarbituric acid reactive species (TBARS) and reduction in reduced glutathione (GSH) levels which is in line with previous findings (El-Sherbiny et al, 2003; Khalifa, 2004; Agrawal et al, 2008).

In the our investigation, pretreatment of erythropoietin (EPO) significantly reversed the memory deficits induced by scopolamine along with significant attenuation of scopolamine mediated rise in brain AChE activity and brain oxidative stress levels. EPO is a hematopoietic cytokine which has recently been shown to be expressed in the nervous system (Dame et al, 2000; Siren et al, 2001; Juul, 2002). The expression of EPO and its receptor in the nervous system are modulated by hypoxia and metabolic insult (Siren et al, 2001a; 2001b; 2001e). Recent studies suggest that EPO has neuroprotective effects against various insults such as glutamate-induced excitotoxicity, serum deprivation, hypoxia, and growth factor deprivation (Morishita et al, 1997; Lewczuk et al, 2000; Siren et al, 2001a). EPO has been shown to protect primary hippocampal neurons by increasing the expression of brain-derived neurotrophic factor (Viviani et al., 2005). EPO in-vitro has also been observed to exert neurotrophic effect on cholinergic and other neurons (Tabira et al, 1995). In addition erythropoietin has also been found to exert stimulatory effect on the release of dopamine and acetylcholine in rat brain slices (Yamamoto et al, 2000). Many studies have also implicated




Fig. 3. Effect of erythropoietin on scopolamine/streptozotocin-induced changes in brain thio-barbituric acid reactive species (TBARS) levels. ACSF-C: artificial cerebrospinal fluid-control; SCO-C: scopolamine-control; EPO-L and H: erythropoietin (low and high); STZ: streptozotocin. Each group (n=7) represents mean \pm standard errors of means. One-way ANOVA followed by Tukey's multiple range test; F (9, 60)=10.721, $p < 0.001$. ^a $p < 0.05$ Vs control group, ^b $p < 0.05$ Vs scopolamine, ^c $p < 0.05$ Vs ACSF control, ^d $p < 0.05$ Vs STZ.




Fig. 4. Effect of erythropoietin on scopolamine/streptozotocin-induced changes in brain reduced glutathione (GSH) levels. ACSF-C: artificial cerebrospinal fluid-control; SCO-C: scopolamine-control; EPO-L and H: erythropoietin (low and high); STZ: streptozotocin. Each group (n=7) represents mean \pm standard errors of means. One-way ANOVA followed by Tukey's multiple range test; F (9, 60)=16.522, $p < 0.001$. ^a $p < 0.05$ Vs control group, ^b $p < 0.05$ Vs scopolamine, ^c $p < 0.05$ Vs ACSF control, ^d $p < 0.05$ Vs STZ.

potential anti-oxidative (Genc et al, 2004; Signore et al, 2006; Xue et al, 2007; Wang et al, 2009) actions of EPO. Few recent reports have documented a prominent role of erythropoietin in process of long term-potentialiation (LTP) hence memory formation (Adamcio et al, 2008; El-Kordi et al, 2009). Erythropoietin probably exerts its actions via acting as a ligand for erythropoietin receptors and stimulating erythropoietin receptor mediated signaling cascade, which certainly plays a prominent role in process of LTP and memory (Adamcio et al, 2008; El-Kordi et al, 2009). Therefore it may be put forth that EPO in our investigation attenuated scopolamine -induced memory deficits and associated biochemical changes by virtue of its multiple actions including anti-oxidative, neuroprotective, and anti-cholinesterase actions. Anti-cholinesterase activity of EPO in brain tissue perhaps, being first report in this study. However it is difficult say at this point whether it is a direct or indirect effect of EPO.

Intracerebroventricular streptozotocin (STZ, i.c.v) in our study has also impaired learning and memory along with significant rise in brain oxidative stress levels and brain AChE activity of mice. The STZ i.c.v model has been described an appropriate animal model of dementia closely related to Alzheimer's disease, typically characterized by progressive impairment of learning abilities and memory capacities (Lannert and Hoyer, 1998). Cerebral glucose and energy metabolism is associated with oxidative stress. After i.c.v administration, the highest concentration of STZ (3 mg/kg) reaches the fornix and periventricular white matter at the level of 3rd ventricle, which shows the greatest damage (Shoham et al, 2003) and STZ i.c.v induced memory impairment is independent of its hyperglycemic effect (Mayer et al, 1990). Although the mechanism of action of STZ i.c.v on memory impairment is not yet known, it probably involves the induction of oxidative stress (Feillet-Coudray et al, 1999; Reagan et al, 2000) to which myelin is particularly vulnerable (Smith et al, 1999). Damage to myelin by oxidative stress is seen in disorders such as dementia of AD type with cognitive impairment (Braak et al, 2000). It causes desensitization of insulin receptors and biochemical changes similar to AD or ageing brain (Hoyer, 2000a; 2000b). In addition, reduced energy metabolism and synthesis of acetyl-CoA ultimately results in cholinergic deficiency and thereby memory deficit in STZ treated animals. Recent animal studies have demonstrated that STZ i.c.v produces brain changes that are hallmark of human AD (Salkovic-Petrisic et al, 2006; Grünblatt et al, 2007). This observation is in line with our previous reports whereby intracerebroventricular administration of streptozotocin at sub-diabetogenic dose has been shown to induce memory deficits along with increase in brain oxidative stress levels and brain AChE activity (Sharma et al, 2008a; 2008b; Kaur et al, 2009).

Administration of EPO to STZ i.c.v treated mice significantly prevented memory deficits as well as STZ induced elevation in brain oxidative stress levels and brain AChE activity. As mentioned above EPO in various studies has been reported to exert number of important actions like anti-oxidative (Genc et al, 2004; Signore et al, 2006; Xue et al, 2007; Wang et al, 2009), neuroprotective (Morishita et al, 1997; Lewczuk et al, 2000; Siren et al, 2001b), neurotrophic (Viviani et al, 2005) effects and a vital role in hippocampal LTP (Adamcio et al, 2008; El-Kordi et al, 2009). EPO has also shown to exert potential anti-inflammatory actions (Rui et al, 2005; Liu et al, 2006) and role of in-

flammation in the pathobiology of AD is well documented (Granic et al, 2009; Salminen et al, 2009). Therefore, with support from literature and data in hand it may be proposed that EPO mediated beneficial effect in STZ (i.c.v) dementia may be attributed to its multiple effects including anti-oxidative, anti-cholinesterase, anti-inflammatory and neuroprotective actions. Hence, it may be concluded that EPO has shown ameliorative effect in memory deficits and brain oxidative stress associated with experimental dementia. Nevertheless further studies including brain histopathological analysis are needed and underway to explore full potential of erythropoietin in dementia and to find out exact mechanism of erythropoietin transduction system in its memory preserving/improving effect.

REFERENCES

- Adamcio B, Sargin D, Stradomska A, Medrihan L, Gertler C, Theis F, Zhang M, Müller M, Hassouna I, Hannke K, Sperling S, Radyushkin K, El-Kordi A, Schulze L, Ronnenberg A, Wolf F, Brose N, Rhee JS, Zhang W, Ehrenreich H. Erythropoietin enhances hippocampal long-term potentiation and memory. *BMC Biol.* 2008;6:37.
- Agrawal R, Tyagi E, Shukla R, Nath C. Effect of insulin and melatonin on acetylcholinesterase activity in the brain of amnesic mice. *Behav Brain Res.* 2008;189:381-386.
- Bartus RT, Dean RT, Beer B, Lippa AS. The cholinergic hypothesis of geriatric memory dysfunction. *Science.* 1982;217:408-417.
- Becker R. Therapy of cognitive deficit of Alzheimer's disease: the cholinergic system in cholinergic basis for Alzheimer therapy. Becker R, Giacobini E, eds. Boston: Birkhauser; 1991. pp. 22.
- Beutler RG, Duron O, Kelly B. Reduced glutathion estimation. *J Lab Clinical Med.* 1963;61:82.
- Blokland A. Acetylcholine: a neurotransmitter for learning and memory? *Brain Res Rev.* 1995;21:285-300.
- Braak H, Del Tredici K, Schultz C, Braak E. Vulnerability of select neuronal types of Alzheimer's disease. *Ann N Y Acad Sci USA.* 2000;924:53-61.
- Brines ML, Ghezzi P, Keenan S, Agnello D, de Lanerolle NC, Cerami C, Itri LM, Cerami A. Erythropoietin crosses the blood-brain barrier to protect against experimental brain injury. *Proc Natl Acad Sci USA.* 2000;97:10526-10531.
- Campana WM, Misasi R, O'Brien JS. Identification of a neurotrophic sequence in erythropoietin. *Int J Mol Med.* 1998;1:235-241.
- Chandel NS, Maltepe E, Goldwasser E, Mathieu CE, Simon MC, Schumacke PT. Mitochondrial reactive oxygen species trigger hypoxia-induced transcription. *Proc Natl Acad Sci USA.* 1998;95:11715-11720.
- Coyle JT, Price DL, DeLong MR. Alzheimer's disease: a disorder of cortical cholinergic innervations. *Science.* 1983;219:1184-1190.
- Dame C, Bratmann P, Wolber EM, Fahnenstich H, Hofman D, Fandrey J. Erythropoietin gene expression in different areas of the developing human central nervous system. *Brain Res Dev Brain Res.* 2000;125:69-74.
- Deutsch JA, Rocklin KW. Amnesia induced by scopolamine and its temporal variations. *Nature.* 1967;216:89-90.
- Deutsch JA. The cholinergic synapse and the site of memory. *Science.* 1971;174:788-794.
- El-Kordi A, Radyushkin K, Ehrenreich H. Erythropoietin improves operant conditioning and stability of cognitive performance in mice. *BMC Biol.* 2009;7:37.
- Ellman GL, Courtney DK, Andres V, Feathstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol.* 1961;7:88-95.
- El-Sherbiny DA, Khalifa AE, Attia AS, Eldenshary ED. Hypericum perforatum extract demonstrates anti oxidant properties against elevated rat brain oxidative status induced by amnesic dose of scopolamine. *Pharmacol Biochem Behav.* 2003;76:525-533.

- Feillet-Coudray C, Rock E, Coudray C, Grzelkowska K, Azais-Braesco V, Dardevet D. Lipid peroxidation and anti oxidant status in experimental diabetes. *Clin Chim Acta*. 1999;284:31-43.
- Genc S, Koroglu TF, Genc K. Erythropoietin and the nervous system. *Brain Res*. 2004;1000:19-31.
- Granic I, Dolga AM, Nijholt IM, Van Dijk G, Eisel UL. Inflammation and NF-kappaB in Alzheimer's disease and diabetes. *J Alzheimers Dis*. 2009;16:809-821.
- Grünblatt E, Salkovic-Petrisic M, Osmanovic J, Riederer P, Hoyer S. Brain insulin system dysfunction in streptozotocin intracerebroventricularly treated rats generates hyperphosphorylated tau protein. *J Neurochem*. 2007;101:757-770.
- Haley TJ, McCormick WG. Pharmacological effects produced by intracerebral injection of drugs in the conscious mice. *Br J Pharmacol Chemother*. 1957;12:12-15.
- Hoyer S. The aging brain. Changes in the neuronal insulin / insulin receptor signal transduction cascade trigger late onset sporadic Alzheimer's disease (SAD). A mini review. *J Neural Transm*. 2000a;109:991-1002.
- Hoyer S. The brain insulin signal transduction system and sporadic (type II) Alzheimer's disease: an update. *J Neural Transm*. 2000b;109:341-360.
- Juul S. Erythropoietin in the central nervous system, and its use to prevent hypoxic-ischemic brain damage. *Acta Paediatr Suppl*. 2002;91:36-42.
- Juul SE, Yachnis AT, Rojiani AM, Christensen RD. Immunohistochemical localization of erythropoietin and its receptor in the developing human brain. *Pediatr Dev Pathol*. 1999;2:148-158.
- Kaur B, Singh N, Jaggi AS. Exploring mechanism of pioglitazone-induced memory restorative effect in experimental dementia. *Fundam Clin Pharmacol*. 2009;23:557-566.
- Khalifa AE. Pro oxidant activity of Zuclopenthixol in vivo: Differential effect of the drug on brain oxidative status of scopolamine treated rats. *Hum Exp Toxicol*. 2004;23:439-445.
- Konishi Y, Chui DH, Hirose H, Kunishita T. Trophic effect of erythropoietin and other hematopoietic factors on central cholinergic neurons in vitro and in vivo. *Brain Res*. 1993;609:29-35.
- Lannert H, Hoyer S. Intracerebroventricular administration of Streptozotocin causes long term diminutions in learning and memory abilities and in cerebral energy metabolism in adult rats. *Behav Neurosci*. 1998;112:1199-1208.
- Lewczuk P, Hasselblatt M, Kamrowski-Kruck H, Heyer A, Unzicker C, Siren AL. Survival of hippocampal neurons in culture upon hypoxia: effect of erythropoietin. *Neuroreport*. 2000;11:3485-3488.
- Liu X, Xie W, Liu P, Duan M, Jia Z, Li W. Mechanism of the cardioprotection of rhEPO pretreatment on suppressing the inflammatory response in ischemia-reperfusion. *Life Sci*. 2006;78:2255-2264.
- Lowry OH, Rosenbrough NJ, Farr AL, Randall RJ. Protein measurement with folin phenol reagent. *J Biol Chem*. 1951;193:265-275.
- Marti HH, Wenger RH, Rivas LA, Straumann U, Diczajloglu M, Henn V. Erythropoietin gene expression in human, monkey and murine brain. *Eur J Neurosci*. 1996;8:666-676.
- Masuda S, Okano M, Yamagishi K, Nagao M, Ueda M, Sasaki R. A novel site of erythropoietin production: oxygen-dependent production in cultured rat astrocytes. *J Biol Chem*. 1994;269:19488-19493.
- Mayer G, Nitsch R, Hoyer S. Effects of changes in peripheral and cerebral glucose metabolism on locomotor activity, learning and memory in adult male rats. *Brain Res*. 1990;532:95-100.
- Morishita E, Masuda S, Nagao M, Yasuda Y, Sasaki R. Erythropoietin receptor is expressed in rat hippocampal and cerebral cortical neurons and erythropoietin prevents in vitro glutamate-induced neuronal death. *Neuroscience*. 1997;76:105-116.
- Morris RGM. Developments of a water maze producer for studying spatial learning in the rats. *J Neurosci Meth*. 1984;11:47-60.
- Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem*. 1979;95:351-358.
- Packard MG, Teather LA, Bazan NG. Effect of intra-striated injections of platelet-activating factor and PAF antagonist BN 52021 on memory. *Neurobiol Learn Mem*. 1996;66:176-182.
- Parle M, Singh N. Animal models of testing memory. *Asia pacific J Pharmacol*. 2004;16:101-120.
- Parle M, Singh N. Reversal of memory deficits by atorvastatin and simvastatin in rats. *Yakugaku Zasshi*. 2007;127:1125-1137.
- Reagan LP, Magariños AM, Yee DK, Swzeda LI, Van Bueren A, McCall AL, McEwen BS. Oxidative stress and HNE conjugation of GLUT3 are increased in the hippocampus of diabetic rats subjected to stress. *Brain Res*. 2000;862:292-300.
- Rui T, Feng Q, Lie M, Peng T, Zhang J, Xu M. Erythropoietin prevents the acute myocardial inflammatory response induced by ischemia/reperfusion via induction of AP-1. *Cardiovasc Res*. 2005;65:719-727.
- Sakurada T, Sakurada S, Katsuyama S, Sakurada C, No KT, Terenius L. Noceptin (1-7) antagonizes noceptin induced hyperalgesia in mice. *Br J Pharmacol*. 1999;128:941-944.
- Salkovic-Petrisic M, Tribl F, Schmidt M, Hoyer S, Riederer P. Alzheimer-like changes in protein kinase B and glycogen synthase kinase-3 in rat frontal cortex and hippocampus after damage to the insulin signalling pathway. *J Neurochem*. 2006;96:1005-1015.
- Salminen A, Ojala J, Kauppinen A, Kaamiranta K, Suuronen T. Inflammation in Alzheimer's disease: amyloid-beta oligomers trigger innate immunity defence via pattern recognition receptors. *Prog Neurobiol*. 2009;87:181-194.
- Sharma B, Singh N, Singh M, Jaggi AS. Exploitation of HIV protease inhibitor indinavir as a memory restorative agent in experimental dementia. *Pharmacol Biochem Behav*. 2008b;89:535-545.
- Sharma B, Singh N, Singh M. Modulation of celecoxib and Streptozotocin induced experimental dementia of Alzheimer's disease type by pitavastatin and donepezil. *J Psychopharmacol*. 2008a;22:162-171.
- Sharma M, Gupta YK. Effect of chronic treatment of melatonin on learning, memory and oxidative deficiencies induced by intracerebroventricular Streptozotocin in rats. *Pharmacol Biochem Behav*. 2001;70:325-331.
- Shoham S, Bejar C, Kovalev E, Weinstock M. Intracerebroventricular injection of streptozotocin causes neurotoxicity to myelin that contributed to spatial memory deficits in rats. *Exp Neurol*. 2003;184:1043-1052.
- Signore AP, Weng Z, Hastings T, Van Laar AD, Liang Q, Lee YJ. Erythropoietin protects against 6-hydroxydopamine induced dopaminergic cell death. *J Neurochem*. 2006;96:428-443.
- Siren AL, Enrenreich H. Erythropoietin a novel concept for neuroprotection. *Eur Arch Psychiatry Clin Neurosci*. 2001;251:179-184.
- Siren AL, Fratelli M, Brines M, Goemans C, Casagrande S, Lewczuk P, Keenan S, Gleiter C, Pasquali C, Capobianco A, Mennini T, Heumann R, Cerami A, Ehrenreich H, Ghezzi P. Erythropoietin prevents neuronal apoptosis after cerebral ischemia and in metabolically stressed neurons. *Proc Natl Acad Sci USA*. 2001;98:4044-4049.
- Siren AL, Knerlich F, Poser W, Gleiter CH, Bruck W, Ehrenreich H. Erythropoietin and erythropoietin receptor in human ischemic/hypoxic brain. *Acta Neuropathol*. 2001;101:271-276.
- Smith KJ, Kapoor R, Felts PA. Demyelination: the role of reactive oxygen and nitrogen species. *Brain Pathol*. 1999;9:69-92.
- Tabira T, Konishi Y, Gallyas F Jr. Neurotrophic effect of hematopoietic cytokines on cholinergic and other neurons in vitro. *Int J Dev Neurosci*. 1995;13:241-252.
- Torre DLJ, Aliev G, Perry G. Drug therapy in Alzheimer's disease. *N Engl J Med*. 2004;351:1911-1913.
- Viviani B, Bartsaghi S, Corsini E, Villa P, Ghezzi P, Garau A, Galli CL, Marinovich M. Erythropoietin protects primary hippocampal neurons increasing the expression of brain-derived neurotrophic factor. *J Neurochem*. 2005;93:412-421.
- Voss G, Sachse K. Red cell and plasma cholinesterase activities in microsamples of human and animal blood determined

- simultaneously by a modified acetylcholine/DTNB procedure. *Toxicol Appl Pharmacol.* 1970;16:764-772.
- Wang ZY, Shen LJ, Tu L, Hu DN, Liu GY, Zhou ZL, Lin Y, Chen LH, Qu J.** Erythropoietin protects retinal pigment epithelial cells from oxidative damage. *Free Radic Biol Med.* 2009;46:1032-1041.
- Xue YQ, Zhao LR, Guo WP, Duan WM.** Intrastriatal administration of erythropoietin protects dopaminergic neurons and improves neurobehavioral outcome in a rat model of Parkinson's disease. *Neuroscience.* 2007;146:1245-1258.
- Yamamoto M, Koshimura K, Kawauchi M, Sohmiya M, Murakami Y, Kato Y.** Stimulating effect of Erythropoietin on the release of dopamine and acetylcholine from the rat brain slice. *Neurosci Lett.* 2000;292:131-133.
- Yamazaki M, Matsuoka N, Maeda N, Kuratani K, Ohkubo Y, Yamaguchi I.** FR121196: a potential anti dementia drug ameliorates the impaired memory of rats in Morris water maze. *J Pharmacol Exp Ther.* 1995;272:256-263.