

## Bcl-2 Knockdown Accelerates T Cell Receptor-Triggered Activation-Induced Cell Death in Jurkat T Cells

Yun-Jung Lee, Tae Joon Won, Kyeong Eun Hyung, Mi Ji Lee, Young-hye Moon, Ik Hee Lee, Byung Sung Go, and Kwang Woo Hwang

Laboratory of Host Defense Modulation, College of Pharmacy, Chung-Ang University, Seoul 156-756, Korea

Cell death and survival are tightly controlled through the highly coordinated activation/inhibition of diverse signal transduction pathways to insure normal development and physiology. Imbalance between cell death and survival often leads to autoimmune diseases and cancer. Death receptors sense extracellular signals to induce caspase-mediated apoptosis. Acting upstream of CED-3 family proteases, such as caspase-3, Bcl-2 prevents apoptosis. Using short hairpin RNAs (shRNAs), we suppressed Bcl-2 expression in Jurkat T cells, and this increased TCR-triggered AICD and enhanced *TNFR* gene expression. Also, knockdown of Bcl-2 in Jurkat T cells suppressed the gene expression of *FLIP*, *TNF receptor-associated factors 3 (TRAF3)* and *TRAF4*. Furthermore, suppressed Bcl-2 expression increased caspase-3 and diminished nuclear factor kappa B (NF- $\kappa$ B) translocation.

**Key Words:** AICD, Apoptosis, Bcl-2, shRNA, T cell

### INTRODUCTION

Apoptosis, a form of programmed cell death, plays an essential role in maintaining homeostasis of the immune system [1]. Deregulated apoptotic cell death leads to various autoimmune lymphoproliferative syndromes and cancer [2,3]. Apoptosis is triggered by the activation of death receptors, including Fas (Apo-1 or CD95), tumor necrosis factor (TNF), DR3 (TRAMP), DR4 (TRAIL-R1), and DR5 (TRAIL-R2) [4, 5]. Fas-induced apoptosis is a commonly encountered apoptotic pathway in various cell types. Engagement of Fas by FasL or type I-membrane protein [6,7] triggers apoptosis via the activation of specific caspases [8]. Interaction of Fas with Fas-L is required for the activation-induced cell death (AICD) of previously stimulated T cells [9-12].

TNF is a cytokine produced by various cell types, including macrophages, monocytes and lymphocytes, that induces apoptosis in response to inflammation, infection, injury, and other environmental challenges [13]. TNF has been shown to regulate cell growth, apoptosis, and differentiation, and is a mediator of inflammation as well as cellular immune responses [14-18]. Binding of TNF to its receptors, TNFR1, and less frequently to TNFR2, triggers a cas-

pase-like signaling cascade, subsequently leading to apoptosis or nuclear factor kappa B (NF- $\kappa$ B) activation and cell survival [13]. Caspase-3, a cysteine-aspartic acid protease in the mitochondrial pathway of apoptosis, interacts with caspase-8 and caspase-9. Cleavage and processing of caspase-3 by the action of caspases 8, 9, and 10 leads to the activation of caspases 6 and 7 [19].

Located primarily on the outer mitochondrial membrane, the anti-apoptotic protein Bcl-2 prevents apoptosis by blocking the activation of CED-3 family proteases such as caspase-3 [20-24]. To understand the role of Bcl-2, we suppressed its expression in Jurkat T cells using short hairpin RNAs (shRNAs). Here, we report that Bcl-2 knockdown in Jurkat T cells augmented TNFR gene expression and T cell receptor (TCR)-triggered AICD but, suppressed the expression of FLIP, TNF receptor (TNFR)-associated factors 3 (TRAF3), and TRAF4. In addition, it led to increased caspase-3 cleavage and decreased NF- $\kappa$ B nuclear translocation.

### METHODS

#### *shRNA construction and transfection*

For knockdown of Bcl-2 in Jurkat T cells, four shRNAs targeting human *bcl2* gene (NM\_177410) were constructed using a SureSilencing shRNA plasmid (SABiosciences). The sequences of scrambled control and Bcl2-shRNAs were 5'-TGC

Received November 18, 2013, Revised December 27, 2013,  
Accepted January 16, 2014

Corresponding to: Kwang Woo Hwang, Laboratory of Host Defense Modulation, College of Pharmacy, Chung-Ang University, 221 Heukseok-dong, Dongjak-gu, Seoul 156-756, Korea. (Tel) 82-2-820-5597, (Fax) 82-2-823-5597, (E-mail) khwang@cau.ac.kr



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**ABBREVIATIONS:** shRNAs, short hairpin RNAs; AICD, activation-induced cell death; TNF, tumor necrosis factor; TRAF, TNF receptor-associated factors; NF- $\kappa$ B, nuclear factor kappa B; TCR, T cell receptor; FLIP, FLICE (FADD-like IL-1 $\beta$ -converting enzyme)-inhibitory protein; XIAP, X-linked inhibitor of apoptosis protein.

GAC CTC TGT TTG ATT TCT-3' and 5'-GGG AGG ATT GTG GCC TTC TTT-3', respectively. Jurkat T cells were maintained in RPMI1640 containing 10% heat-inactivated fetal bovine serum (FBS), 100 U/ml penicillin, and 100  $\mu$ g/ml streptomycin. For Bcl-2 knockdown, Jurkat T cells were seeded in 24-well plates at a density of  $1 \times 10^4$  cells/well. After 24 h, cells were then transfected with Bcl2-shRNAs using FuGene 6 Transfection reagent (Roche). Efficiencies of the individual shRNAs in silencing the expression of Bcl-2 were tested, and the most effective shRNA was used in subsequent experiments. To achieve stable silencing of Bcl-2, shRNA-transfected cells were cultured in medium containing G418 for two weeks.

#### Determination of cell growth

Jurkat T cells ( $1 \times 10^5$  cells) were seeded in a 96-well plate and incubated for 24 h. After incubation, 0.4% Trypan Blue was added to the cell suspension, and cell numbers were estimated by counting under a light microscope. Cells stained blue were considered non-viable.

#### Cell viability assay

Bcl2-shRNA-transfected Jurkat T cells were incubated at a density of  $5 \times 10^4$  cells/ml for 24 h in 96-well plates coated with 0.1  $\mu$ g/ml or 1  $\mu$ g/ml anti-CD3/CD28 antibodies (eBioscience). Cells were washed once with phosphate-buffered saline (PBS), resuspended in PBS containing 5  $\mu$ g/ml propidium iodide (PI) (Sigma), and immediately analyzed by using a FACScalibur (BD biosciences) instrument.

#### Gene expression assays

Bcl-2-knockdown Jurkat T cells were incubated with plate-bound anti-CD3 (1  $\mu$ g/ml) and anti-CD28 (1  $\mu$ g/ml) antibodies for 6 h. Total RNA was isolated using TRIzol (Invitrogen) reagent and processed for first strand complementary DNA (cDNA) synthesis. Using cDNA as a template, real-time polymerase chain reaction (PCR) was performed employing SYBR Green Master Mix (Applied Biosystems) on a LightCycler system (Roche). The relative expression levels, normalized with respect to the expression of *GAPDH*, were then calculated as fold induction compared

to the scrambled control without any stimulation.

#### Western blot analysis


Bcl-2-knockdown cell lines were stimulated with plate-bound anti-CD3 antibody (1  $\mu$ g/ml) and anti-CD28 antibody (1  $\mu$ g/ml) for 24 h. Cells were lysed, and the protein present in the lysate was quantified using a BCA protein assay kit (Thermo Fisher Scientific). The proteins were then separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto a polyvinylidene fluoride (PVDF) membrane. After blocking with non-fat dried milk, the membrane was incubated with anti-Bcl-2 antibody or anti-Caspase-3 antibody (Cell Signaling) overnight, followed by horseradish peroxidase (HRP)-conjugated anti-rat IgG antibody for 2 h. Enhanced chemiluminescence (ECL) reagents were used for the detection of proteins.

#### Detection of nuclear NK- $\kappa$ B

Bcl-2-knockdown Jurkat T cells were incubated with plate-bound anti-CD3 (1  $\mu$ g/ml) and anti-CD28 (1  $\mu$ g/ml) antibodies for 30 min. Nuclear extracts were prepared using Nuclear Extraction Kit (Panomics) and incubated with biotinylated DNA probe (Panomics). The mixture was separated on polyacrylamide gel and the separated proteins were transferred onto a nylon membrane. After incubation with HRP-conjugated streptavidin, protein/DNA complexes were visualized using ECL reagents.

#### Determination of cytokine production

Scrambled- or Bcl2-shRNA-transfected Jurkat T cells were incubated with plate-bound anti-CD3 and anti-CD28 antibodies for 24 h, and the supernatants from the cultures were collected immediately. Microplates were coated overnight with purified anti-human interferon (IFN)- $\gamma$  or anti-human interleukin (IL)-2 antibodies. Individual wells of the plate were blocked with a solution of 3% bovine serum albumin in PBS, and incubated overnight with the cell culture supernatants. After washing to remove unbound substances, captured cytokines were detected by applying biotinylated antibodies followed by streptavidin-alkaline



**Fig. 1.** shRNAs knock down Bcl-2 in Jurkat T cells. The *bcl-2* gene is knocked down using shRNA in Jurkat T cells. (A) Total RNA was extracted from Bcl-2-knockdown and control Jurkat T cells and *bcl-2* gene expression was analyzed by real-time PCR using *GAPDH* as the internal control. (B) Total protein was extracted from Bcl2-knockdown and control Jurkat T cells. Bcl-2 protein expression was analyzed by western blotting using  $\beta$ -actin expression as a loading control. (C) A total of  $1 \times 10^5$  Jurkat T cells were seeded in a 96-well plate and incubated for 24 h. After incubation, 0.4% Trypan Blue was added to the cell suspension, and cell numbers were estimated by counting under a microscope. Cells stained blue were considered non-viable. Data were presented as mean  $\pm$  SD for triplicate determinations. Student's t test; \* $p < 0.05$ ; \*\* $p < 0.01$ ; and \*\*\* $p < 0.001$  vs. control sample. All data were representative of at least three individual experiments.

phosphatase. Following washes, *p*-nitrophenyl phosphate (Sigma) was added to the wells and incubated for 30 min. Color developed in each well was determined by measuring optical density at 405 nm using a microplate reader (Molecular Devices).

## RESULTS

### *Bcl-2*-shRNA suppresses *Bcl-2* mRNA and protein expression in Jurkat T cells


We employed shRNAs for stable knockdown of Bcl-2 expression in human T cell lines. Four shRNA constructs targeting human *bcl2* were separately transfected into Jurkat T cells, and their ability to suppress gene expression was analyzed by real-time PCR. The most effective shRNA suppressed *bcl2* mRNA expression by 13.13-fold (Fig. 1A), leading to significantly lower expression of Bcl-2 protein (Fig. 1B). To examine if the silencing of *bcl-2* affected cell proliferation, we cultured Jurkat T cells stably expressing Bcl-2-shRNA for 24 h and compared the cell numbers with that of control-transfected cells. In 24 h, Bcl-2-knockdown Jurkat T cells proliferated to reach 4-fold the numbers of cells initially seeded, while the numbers of control cells increased 5-fold (Fig. 1C). The ratio of non-viable cells to the total number of cells was similar for control and Bcl-2-knockdown cultures. Therefore, suppression of *bcl2* expression in Jurkat T cells likely resulted in slower cell

proliferation.


### *Bcl-2* knockdown increases TCR-triggered AICD and downregulates *FLIP* gene expression

We stimulated AICD in Bcl-2-knockdown Jurkat T cells using anti-CD3 and anti-CD28 antibodies. Less than 10% of control T lymphocytes underwent cell death following CD3/CD28 stimulation, whereas Bcl-2 knockdown led to increased cell death (14% and 36%, respectively in the presence of 0.1 and 1.0  $\mu$ g/ml antibodies) (Fig. 2A). In the absence of stimuli, both control and Bcl-2 knockdown cells showed comparable cell death rates. To investigate if Bcl-2 knockdown led to any changes in the cell death pathway, we analyzed the gene expression of Fas ligand (FasL) and Fas. Both control and Bcl-2-knockdown Jurkat T cells showed increased *FasL* gene expression in response to stimulation by CD3/CD28 agonists (Fig. 2B). However, the observed increase in *FasL* mRNA expression was lower in Bcl-2-knockdown cells than in controls. By contrast, neither Bcl-2 knockdown nor stimulation by anti-CD3/anti-CD28 antibodies altered the expression of *Fas* in Jurkat T cells (Fig. 2C).

Next, we examined the gene expression of Flice-like inhibitory protein (FLIP) and X-linked inhibitor of apoptosis protein (XIAP). Stimulation of TCR/CD28 resulted in augmented expression of FLIP mRNA in control Jurkat T cells (Fig. 2D). Bcl-2 knockdown lowered the basal levels of FLIP mRNA in Jurkat T cells by 0.08-fold, and kept suppressing



**Fig. 2.** Bcl-2 knockdown increases TCR-triggered AICD, downregulates *FLIP* gene expression, and upregulates caspase-3 cleavage. (A) Bcl-2-knockdown and control Jurkat T cells were incubated with 0.1  $\mu$ g/ml or 1  $\mu$ g/ml of plate bound anti-CD3 and anti-CD28 antibody for 24 h. Cells were washed with PBS, resuspended in PBS containing 5  $\mu$ g/ml of PI, and analyzed by flow cytometry. (B~E) Bcl2-knockdown and control Jurkat T cells were incubated with 1  $\mu$ g/ml of plate-bound anti-CD3 and anti-CD28 antibodies for 6 h. Total RNA was isolated, reverse transcribed, and gene expression was analyzed by real-time PCR. Relative gene expression levels were normalized with respect to those of *GAPDH*. (F) Bcl2-knockdown and control Jurkat T cells were incubated with 1  $\mu$ g/ml of plate-bound anti-CD3 and anti-CD28 antibodies for 24 h. Total protein was extracted, and procaspase-3 and cleaved caspase-3 protein expression was analyzed by western blot.  $\beta$ -actin protein expression was used as a loading control. Data were presented as mean $\pm$ SD for triplicate determinations. Student's *t* test; \**p*<0.05; \*\**p*<0.01; and \*\*\**p*<0.001 vs. control sample. All data were representative of at least three individual experiments.



**Fig. 3.** Bcl-2 knockdown enhances the expression of *TNFR* and suppresses *TRAF* gene expression. Bcl-2-knockdown and control Jurkat T cells were incubated with 1  $\mu$ g/ml of plate-bound anti-CD3 and anti-CD28 antibodies for 6 h. Total RNA was extracted and reverse transcribed, and gene expression was analyzed by real-time PCR. Relative gene expression levels of (A) *TNFR1*, (B) *TNFR2*, (C) *TRAF3*, (D) *TRAF4* were normalized to those of *GAPDH* used as the internal control. Data were presented as mean $\pm$ SD for triplicate determinations. Student's t test; \* $p < 0.05$ ; \*\* $p < 0.01$ ; and \*\*\* $p < 0.001$  vs. control sample. All data were representative of at least three individual experiments.

strongly, at a 0.76-fold-change level relative to unstimulated control cells, even under activation with anti-CD3/CD28 antibodies. Neither Bcl-2 knockdown nor TCR stimulation resulted in any significant changes in caspase inhibitor XIAP mRNA expression (Fig. 2E). We confirmed that the augmented cell death observed in CD3/CD28-activated Bcl-2-knockdown Jurkat T cells was due to increased caspase-3 cleavage (Fig. 2F). Therefore, the Bcl-2 knockdown induced a TCR-mediated AICD that was accompanied by attenuated FLIP mRNA expression.

#### ***Bcl-2 knockdown enhances the expression of TNFR and suppresses TRAF gene expression***

Despite the reduced levels of FasL mRNA expression, Bcl-2-knockdown Jurkat T cells showed augmented cell death compared to control cells following TCR/CD28 stimulation. To determine the mechanisms responsible for the enhanced AICD observed in Bcl-2-knockdown cells, we analyzed the mRNA expression levels of TNFR1, TNFR2, TRAF3, and TRAF4. Following CD3 and CD28 stimulation, mRNA expression of the TNF- $\alpha$  receptors, TNFR1 and TNFR2, increased in Bcl-2-knockdown Jurkat T cells (Fig. 3A and B). By contrast, both basal- and agonist-stimulated expression of the inhibitors of apoptosis, TRAF3 and TRAF4, were lower in Bcl-2-knockdown cells than in control cells (Fig. 3C and D). Therefore, Bcl-2 knockdown in Jurkat T cells led to augmented expression of *TNFR* and suppressed *TRAF3*, *TRAF4* gene expression.

#### ***Bcl-2 knockdown suppresses the nuclear translocation of NF- $\kappa$ B***


The transcription factor NF- $\kappa$ B regulates both cell death and survival. We examined if Bcl-2 knockdown interfered

with the activation of the NF- $\kappa$ B pathway in Jurkat T lymphocytes. Nuclear translocation of NF- $\kappa$ B was observed in TCR/CD28-stimulated control cells. Bcl-2 knockdown led to impaired NF- $\kappa$ B nuclear translocation (Fig. 4A). To confirm that Bcl-2 knockdown resulted in impaired NF- $\kappa$ B activation, we measured mRNA and secreted protein levels of IFN- $\gamma$  as well as IL-2. Expression of IFN- $\gamma$  transcripts and protein increased in agonist-stimulated control-transfected cells (Fig. 4B). Bcl-2 knockdown led to attenuated IFN- $\gamma$  mRNA expression and lower IFN- $\gamma$  secretion even in the presence of anti-CD3/anti-CD28 antibodies. Expression of IL-2 mRNA and protein were also found to be lower in Bcl-2-knockdown Jurkat T cells than in control-transfected cells (Fig. 4C).

## **DISCUSSION**

Apoptosis is a developmentally and physiologically critical process that is tightly regulated by the coordinated action of diverse extracellular cues and intracellular signaling molecules [25-28]. Bcl-2 is an anti-apoptotic protein that blocks the release of cytochrome c from mitochondria. Since this molecule keeps the apoptosis pathway in check, numerous studies have focused on the roles of Bcl-2 in development, physiology, and pathophysiology. Overexpression of Bcl-2 prevents the initiation of apoptosis by blocking the efflux of cytochrome c from the mitochondria [20]. Bcl-2 overexpression in lymphoid cells suppresses apoptosis and promotes T and B cell development [29]. Furthermore, it has been shown that overexpression of Bcl-2 delays caspase-3 activation and rescues cerebellar degeneration in prion-deficient mice [30].

In our study, we knocked down Bcl-2 expression in Jurkat T cells using shRNA. Stable knockdown of Bcl-2 in



**Fig. 4.** Bcl-2 shRNA suppresses the nuclear translocation of NF- $\kappa$ B. (A) Bcl-2-knockdown and control Jurkat T cells were incubated with 1  $\mu$ g/ml of plate-bound anti-CD3 and anti-CD28 antibodies for 30 min. Nuclear extracts were prepared and analyzed by using an electrophoretic mobility shift assay (EMSA). (B, C) Bcl-2 knockdown and control Jurkat T cells were incubated with 1  $\mu$ g/ml of plate-bound anti-CD3 and anti-CD28 antibodies. After 24 h, supernatants were collected and analyzed for cytokines by using the enzyme-linked immunosorbent assay (ELISA) method. After 6 h, total RNA was isolated and reverse transcribed, and gene expression was analyzed by real-time PCR. Relative gene expression levels were normalized to those of *GAPDH* used as the internal control.

Jurkat T cells led to moderate retardation of cell growth. Compared to control-transfected cells, Bcl-2 knockdown Jurkat T cells underwent significantly higher AICD following stimulation by anti-CD3 and anti-CD28 antibodies.

Engagement of Fas by FasL induces AICD in T cells. Although Bcl-2 knockdown led to enhanced agonist-induced cell death in Jurkat T cells compared to control-transfected cells, the relative expression levels of FasL mRNA followed a reversed trend. Thus, the agonist-induced increase in FasL mRNA expression was lower in Bcl-2 knockdown cells than in controls. No differences in the relative expression of Fas mRNA were observed between these cells. Therefore, we conclude that the increased apoptosis observed in Bcl-2 knockdown cells was not mediated by the Fas or FasL-related apoptosis pathways.

Caspases are crucial mediators of programmed cell death (apoptosis). Caspase-3 catalyzes the cleavage of several key proteins [31]. FLIP and XIAP inhibit programmed cell death by inhibiting caspase-3 activation. Bcl-2 knockdown in Jurkat T cells led to reduced mRNA expression of FLIP both in the absence and presence of agonists, whereas mRNA expression of XIAP remained unchanged. In addition, CD3/CD28 activation in Bcl-2-knockdown Jurkat T cells resulted in increased caspase-3 cleavage. Therefore, we conclude that knockdown of Bcl-2 induced TCR-mediated AICD, down-regulated FLIP mRNA levels, and up-regulated caspase-3 cleavage in Jurkat T cells.

TNF binding to its receptors induces apoptosis by triggering a caspase-like cascade. TNFR activation initiates signal transduction by recruiting one or more adaptor molecules known as TRAFs [32,33]. TRAF2, TRAF5, and TRAF6 induce activation of the NF- $\kappa$ B pathway. However, unlike other TRAFs, overexpression of TRAF3 fails to activate NF- $\kappa$ B [34]. Instead, TRAF3 promotes the ubiquitin-mediated degradation of NF- $\kappa$ B [35]. TNFR-induced apoptosis is negatively regulated by TRAF4 [36].

Compared to control-transfected cells, Bcl-2-knockdown

Jurkat T cells expressed increased levels of TNFR1 and TNFR2 mRNAs, and reduced levels of TRAF3 and TRAF4 transcripts. Therefore, Bcl-2 knockdown initiated cell death by augmenting *TNFR* gene expression and suppressing the expression of TRAF genes.

NF- $\kappa$ B plays key roles in regulating both cell death and survival. We found that Bcl-2 knockdown led to altered mRNA expression of TRAF3 and TRAF4. Since it was reported that TRAFs regulate NF- $\kappa$ B activation [32,33,35,36], we sought to examine the possible involvement of NF- $\kappa$ B in the increased cell death observed in Bcl-2 knockdown cells. Compared to control-transfected cells, Bcl-2-knockdown Jurkat T cells showed impaired nuclear translocation of NF- $\kappa$ B following TCR/CD28-stimulation. Furthermore, Bcl-2 knockdown cells expressed reduced levels of IFN- $\gamma$  and IL-2 compared to control cells under the same conditions. Taken together, Bcl-2 silencing in Jurkat T cells suppressed nuclear translocation of NF- $\kappa$ B and exerted an inhibitory effect on the transcription of anti-apoptotic molecules, including FLIP, TRAF3, and TRAF4 amidst unexpectedly lowering cellular FasL mRNA levels.

In conclusion, knockdown of Bcl-2 in Jurkat T cells led to elevated agonist-induced AICD, likely via a mechanism that involves enhanced *TNFR1* and *TNFR2* gene expression, attenuated expression of *FLIP* and *TRAF3*, increased caspase-3 cleavage, and reduced NF- $\kappa$ B nuclear translocation.

## ACKNOWLEDGEMENTS

This research was supported by the Chung-Ang University Excellent Student Scholarship and Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2010-0012877).

## REFERENCES

1. Van Parijs L, Biuckians A, Abbas AK. Functional roles of Fas and Bcl-2-regulated apoptosis of T lymphocytes. *J Immunol.* 1998;160:2065-2071.
2. Rieux-Laucat F, Le Deist F, Fischer A. Autoimmune lymphoproliferative syndromes: genetic defects of apoptosis pathways. *Cell Death Differ.* 2003;10:124-133.
3. Igney FH, Krammer PH. Death and anti-death: tumour resistance to apoptosis. *Nat Rev Cancer.* 2002;2:277-288.
4. Ashkenazi A, Dixit VM. Death receptors: signaling and modulation. *Science.* 1998;281:1305-1308.
5. Schmitz I, Kirchhoff S, Krammer PH. Regulation of death receptor-mediated apoptosis pathways. *Int J Biochem Cell Biol.* 2000;32:1123-1136.
6. Itoh N, Yonehara S, Ishii A, Yonehara M, Mizushima S, Sameshima M, Hase A, Seto Y, Nagata S. The polypeptide encoded by the cDNA for human cell surface antigen Fas can mediate apoptosis. *Cell.* 1991;66:233-243.
7. Trauth BC, Klas C, Peters AM, Matzku S, Möller P, Falk W, Debatin KM, Krammer PH. Monoclonal antibody-mediated tumor regression by induction of apoptosis. *Science.* 1989;245:301-305.
8. Fraser A, Evan G. A license to kill. *Cell.* 1996;85:781-784.
9. Russell JH, Wang R. Autoimmune gld mutation uncouples suicide and cytokine/proliferation pathways in activated, mature T cells. *Eur J Immunol.* 1993;23:2379-2382.
10. Russell JH, Rush B, Weaver C, Wang R. Mature T cells of autoimmune lpr/lpr mice have a defect in antigen-stimulated suicide. *Proc Natl Acad Sci U S A.* 1993;90:4409-4413.
11. Bossu P, Singer GG, Andres P, Ettinger R, Marshak-Rothstein A, Abbas AK. Mature CD4+ T lymphocytes from MRL/lpr mice are resistant to receptor-mediated tolerance and apoptosis. *J Immunol.* 1993;151:7233-7239.
12. Alderson MR, Tough TW, Davis-Smith T, Braddy S, Falk B, Schooley KA, Goodwin RG, Smith CA, Ramsdell F, Lynch DH. Fas ligand mediates activation-induced cell death in human T lymphocytes. *J Exp Med.* 1995;181:71-77.
13. Bau D, Karin M. Signal transduction by tumor necrosis factor and its relatives. *Trends Cell Biol.* 2001;11:372-377.
14. Beutler B, Cerami A. Tumor necrosis, cachexia, shock, and inflammation: a common mediator. *Annu Rev Biochem.* 1988;57:505-518.
15. Beutler B, Cerami A. The biology of cachectin/TNF- $\alpha$  primary mediator of the host response. *Annu Rev Immunol.* 1989;7:625-655.
16. Paul NL, Ruddle NH. Lymphotoxin. *Annu Rev Immunol.* 1988;6:407-438.
17. Vassalli P. The pathophysiology of tumor necrosis factors. *Annu Rev Immunol.* 1992;10:411-452.
18. Laster SM, Wood JC, Gooding LR. Tumor necrosis factor can induce both apoptotic and necrotic forms of cell lysis. *J Immunol.* 1988;141:2629-2634.
19. Alnemri ES, Livingston DJ, Nicholson DW, Salvesen G, Thornberry NA, Wong WW, Yuan J. Human ICE/CED-3 protease nomenclature. *Cell.* 1996;87:171.
20. Yang J, Liu X, Bhalla K, Kim CN, Ibrado AM, Cai J, Peng TI, Jones DP, Wang X. Prevention of apoptosis by Bcl-2: release of cytochrome c from mitochondria blocked. *Science.* 1997;275:1129-1132.
21. Chinnaiyan AM, Orth K, O'Rourke K, Duan H, Poirier GG, Dixit VM. Molecular ordering of the cell death pathway. Bcl-2 and Bcl-xL function upstream of the CED-3-like apoptotic proteases. *J Biol Chem.* 1996;271:4573-4576.
22. Erhardt P, Cooper GM. Activation of the CPP32 apoptotic protease by distinct signaling pathways with differential sensitivity to Bcl-xL. *J Biol Chem.* 1996;271:17601-17604.
23. Armstrong RC, Aja T, Xiang J, Gaur S, Krebs JF, Hoang K, Bai X, Korsmeyer SJ, Karanewsky DS, Fritz LC, Tomaselli KJ. Fas-induced activation of the cell death-related protease CPP32 is inhibited by Bcl-2 and by ICE family protease inhibitors. *J Biol Chem.* 1996;271:16850-16855.
24. Shimizu S, Eguchi Y, Kamiike W, Matsuda H, Tsujimoto Y. Bcl-2 expression prevents activation of the ICE protease cascade. *Oncogene.* 1996;12:2251-2257.
25. Vaux DL, Weissman IL, Kim SK. Prevention of programmed cell death in *Caenorhabditis elegans* by human bcl-2. *Science.* 1992;258:1955-1957.
26. Gaumer S, Guénel I, Brun S, Théodore L, Mignotte B. Bcl-2 and Bax mammalian regulators of apoptosis are functional in *Drosophila*. *Cell Death Differ.* 2000;7:804-814.
27. Hengartner MO, Ellis RE, Horvitz HR. *Caenorhabditis elegans* gene *ced-9* protects cells from programmed cell death. *Nature.* 1992;356:494-499.
28. Hengartner MO, Horvitz HR. *C. elegans* cell survival gene *ced-9* encodes a functional homolog of the mammalian proto-oncogene *bcl-2*. *Cell.* 1994;76:665-676.
29. Langenau DM, Jette C, Berghmans S, Palomero T, Kanki JP, Kutok JL, Look AT. Suppression of apoptosis by bcl-2 overexpression in lymphoid cells of transgenic zebrafish. *Blood.* 2005;105:3278-3285.
30. Nicolas O, Gavín R, Braun N, Ureña JM, Fontana X, Soriano E, Aguzzi A, del Río JA. Bcl-2 overexpression delays caspase-3 activation and rescues cerebellar degeneration in prion-deficient mice that overexpress amino-terminally truncated prion. *FASEB J.* 2007;21:3107-3117.
31. Porter AG, Jänicke RU. Emerging roles of caspase-3 in apoptosis. *Cell Death Differ.* 1999;6:99-104.
32. Li Q, Ching AK, Chan BC, Chow SK, Lim PL, Ho TC, Ip WK, Wong CK, Lam CW, Lee KK, Chan JY, Chui YL. A death receptor-associated anti-apoptotic protein, BRE, inhibits mitochondrial apoptotic pathway. *J Biol Chem.* 2004;279:52106-52116.
33. Cui X, Hawari F, Alsaaty S, Lawrence M, Combs CA, Geng W, Rouhani FN, Miskinis D, Levine SJ. Identification of ARTS-1 as a novel TNFR1-binding protein that promotes TNFR1 ectodomain shedding. *J Clin Invest.* 2002;110:515-526.
34. Xu Y, Cheng G, Baltimore D. Targeted disruption of TRAF3 leads to postnatal lethality and defective T-dependent immune responses. *Immunity.* 1996;5:407-415.
35. Liao G, Zhang M, Harhaj EW, Sun SC. Regulation of the NF- $\kappa$ B-inducing kinase by tumor necrosis factor receptor-associated factor 3-induced degradation. *J Biol Chem.* 2004;279:26243-26250.
36. Fleckenstein DS, Dirks WG, Drexler HG, Quentmeier H. Tumor necrosis factor receptor-associated factor (TRAF) 4 is a new binding partner for the p70S6 serine/threonine kinase. *Leuk Res.* 2003;27:687-694.