

Original Article



The Versatility of Framework Regions of Chicken V_H and V_L to Mutations

Jung Won Shin^{1,2}, Sang Il Kim^{2,3}, Aerin Yoon^{1,2}, Junyeong Jin^{1,2}, Hyung Bae Park^{1,4}, Hyori Kim^{5,*}, Junho Chung^{1,2,*}

¹Department of Biochemistry and Molecular Biology, Seoul National University College of Medicine, Seoul 03080, Korea

²Cancer Research Institute, Seoul National University College of Medicine, Seoul 03080, Korea

³Department of Cancer Biology, Seoul National University College of Medicine, Seoul 03080, Korea

⁴Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul 03080, Korea

⁵Biomedical Research Center, Asan Institute for Life Sciences, Asan Medical Center, Seoul 05505, Korea



Received: Oct 3, 2017
Revised: Jan 4, 2018
Accepted: Jan 13, 2018

*Correspondence to

Junho Chung

Department of Biochemistry and Molecular Biology, Seoul National University College of Medicine, Samsung Cancer Research Building 510, 103 Daehak-ro, Jongno-gu, Seoul 03080, Korea.

E-mail: jhchung@snu.ac.kr

Hyori Kim

Biomedical Research Center, Asan Institute for Life Sciences, Asan Medical Center, 88 Olympic-ro 43-gil, Songpa-gu, Seoul 05505, Korea.

E-mail: hyorikim@amc.seoul.kr

Current address: Aerin Yoon, Mogam Institute, Yongin 16924, Korea.

Copyright © 2018. The Korean Association of Immunologists

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Conflict of Interest

The authors declare no potential conflict of interest.

ABSTRACT

To identify the interchangeability of V_H and V_L framework region (FR) residues, we artificially introduced random mutations at all residue positions in a chicken monoclonal antibody, which has only one functional V_H and V_L gene. When we classified the amino acids into 5 groups by their physicochemical properties, all FR residues could be replaced by another group except L23 (C), H36 (W), H86 (D), H104 (G), and H106 (G). Eighty-two (50.9%), 48 (29.8%), 17 (10.6%), and 9 FR residues (5.6%) could be replaced by 4, 3, 2, and 1 group(s), individually, without significant loss of reactivity. We also confirmed a similar level of versatility with 2 different chicken antibodies. This high level of versatility on FR residues has not been predicted because it has not been observed in the 150 chicken antibodies that we previously generated or in the 1,269 naïve chicken V_H sequences publicly available. In conclusion, chicken antibody FR residues are highly interchangeable and this property can be applied for improving the physicochemical property of antibody including thermal stability, solubility and viscosity.

Keywords: Chicken; Ig Framework region; Immunoglobulin somatic hypermutation; Mutation

INTRODUCTION

B cell Ig diversity is achieved in 2 stages (1). The initial diversity is obtained by V(D)J rearrangement of the heavy chain and light chain, which involves Tdt-induced nucleotide addition and exonuclease-mediated deletion at the segment junctions (1). The second stage of diversification is achieved by somatic hypermutation (SHM) (1). SHM is initiated by activation-induced cytidine deaminase to form U/G mismatches followed by several possible fates, including C to T transition mutation, removal of the U-leading C→A/C/T mutation, or excision of neighboring bases and DNA synthesis (1). SHM preferentially targets C bases in the WRCY hot-spot motif (W=A or T, R=G or A, and Y=T or C), its reverse complementary sequence RGYW, or its variants, such as WRCH or DGYW. Since these hot-spot motifs are abundant within complementarity determining regions (CDRs) of Igs, SHM occurs more frequently in CDRs (1) and plays a critical role in the affinity maturation of B cell Igs.

Abbreviations

CDR, complementarity determining region; *E. coli*, *Escherichia coli*; FR, framework region; PSA, prostate specific antigen; scFv, single chain Fv; SHM, somatic hypermutation

Author Contributions

Conceptualization: Shin JW, Yoon A; Data curation: Shin JW, Yoon A; Formal analysis: Kim SI, Jin J; Investigation: Kim SI, Jin J; Project administration: Park HB, Kim H; Supervision: Kim H, Chung J; Writing - original draft: Yoon A; Writing - review & editing: Shin JW, Kim SI.

Following the onset of recombinant antibody technology, artificial mutations were frequently introduced into CDRs to increase the affinity of an antibody (2,3). Development of experimentally-derived high affinity antibodies revealed that mutations of greater diversity can be introduced into CDRs to increase antibody affinity. Meanwhile, artificial mutation of the immunoglobulin framework region (FR) has been attempted for back-to-mouse mutations to restore affinity after grafting mouse CDRs into human Ig FR residues (4). However, if FR residues could be mutated without loss in affinity, improvements in physicochemical properties could be achieved more easily.

In this study, we investigated whether diverse mutations could be introduced into Ig FR residues while preserving antibody affinity. We selected a chicken antibody for this study because the chicken possesses only one functional V_H and V_L gene to make a functional antibody repertoire, even though shuffling with pseudogenes occurs (5). The FR residues of these pseudogenes have been reported to be homologous to the functional genes (5). Therefore, the chicken antibody is an ideal model system for comparing the FR of a natural repertoire to that of a synthetic one. This comparison is relatively less complicated to analyze in chicken than in mouse, rabbit, or human systems, all of which possess multiple V_H and V_L genes.

MATERIALS AND METHODS

Randomization of FR codons

A total of 161 PCRs were performed using degenerate primers to produce gene fragments encoding a portion of PC24 single chain Fv (scFv), an anti-prostate specific antigen (PSA) antibody (Genbank accession numbers KP764766 and KP764767), with an NNK degenerate codon (N=any nucleotide; K=G or T) at each of the 84 V_H and 77 V_L FR residues. The PCR products were fused to gene fragments encoding the remainder of the scFv gene with linker PCR to genes encoding a whole scFv. After restriction digestion with *Sfi I*, the genes were cloned into the pComb3XSS phagemid vector. The ligation products were used to transform competent ER2738 *Escherichia coli* (*E. coli*) to generate 161 scFv libraries as described previously (6). PC17 (another anti-PSA chicken antibody, unpublished) and VCM2 (an anti-VCAM-1 antibody, unpublished) were randomized at 16 framework residues to produce 16 scFv libraries in the same manner. All the amino acid of FR residues were numbered using Kabat numbering system (7).

Phage enzyme immunoassay

The scFv-displaying phages were rescued from titer plates after transformation and then used for phage enzyme immunoassay as described previously (6). Microtiter plates (Corning Inc., NY, USA) were coated with PSA (Fitzgerald Industries International, Acton, MA, USA) or recombinant human VCAM-1 (R&D Systems, Minneapolis, MN, USA) by adding 20 μ l of protein in PBS (5 μ g/ml) to each well followed by incubation overnight at 4°C. After blocking with 3% bovine serum albumin in PBS, the plate was then sequentially incubated with scFv-displaying phages in the culture supernatant, horseradish peroxidase (HRP)-conjugated mouse anti-M13 monoclonal antibody (GE Healthcare, Pittsburgh, PA, USA), and 2,2'-Azinobis [3-ethylbenzothiazoline-6-sulfonic acid]-diammonium salt (ABTS) substrate solution (Amresco, Solon, OH, USA) with intermittent washing using 0.05% Tween-20 in PBS. After incubating the plates at 37°C for 10 min, the optical density was measured at 405 nm using a microtiter plate reader (Labsystems, Barcelona, Spain).

Sequence analysis

Sequence analysis of phagemid clones was performed as described previously (6). Plasmid DNA was prepared using a small-scale plasmid preparation kit (Qiagen, Hilden, Germany). The pComb-F primer (ACA CTT TAT GCT TCC GGC TC) and pComb-R primer (CAA AAT CAC CGG AAC CAG AG) were used to determine the nucleotide sequence of V_L and V_H , respectively. The Accelrys Gene software program (Accelrys, San Diego, CA, USA) was used to analyze the sequence.

Generation of scFv-displaying phages with mutated H82, H82A, and H82B residues

PCR primers were designed to produce 70 mutants with different amino acids at these specific residues. PCR was performed to produce gene fragments encoding a portion of scFv, which was linked to the remainder of the *scFv* gene fragment by linker PCR as described above. The resulting *scFv* genes were digested with *Sfi I* and ligated into the pComb3X phagemid vector prior to transformation into ER2738 *E. coli*. The scFv-displaying phages were rescued from titer plates after transformation and used for phage enzyme immunoassay as described previously (6).

Enzyme immunoassay of chicken-human chimeric IgG1 with mutated H78 and L85 residues

Genes encoding 8 V_H and 16 V_L mutants were synthesized and cloned into an expression vector as described previously (6, 7). The IgG1 molecules were purified via protein A gel affinity chromatography following the instructions provided by the supplier (Repligen, Waltham, MA, USA). To check the affinity of the mutant IgG1s, enzyme immunoassay was performed using PSA-coated microtiter plates and HRP-conjugated anti-human IgG1 antibody (Thermo Fisher Scientific, Waltham, MA, USA) as described previously (6).

Statistical analysis

The statistical analysis was performed using Student's *t*-tests employing SigmaPlot 10.0 software (SPSS, Chicago, IL, USA).

RESULTS

Versatility of the framework residues in chicken scFv

We constructed 161 sets of a scFv phage display library with randomly introduced amino acids at every FR residue within a chicken antibody (V_H and V_L consist of 84 and 77 residues, respectively) using degenerate oligomers. Each set contained a scFv randomized at a single FR residue. Subsequently, we randomly selected 92 clones from each library, rescued the scFv-displaying phage, and performed phage enzyme immunoassay. We defined positive clones as those exhibiting a 70% or greater signal over that of the original scFv phage rescued in parallel. Because in this setting, the absorbance value of positive clones and that of negative clones was different with statistical significance ($p < 0.05$) among all sets of FR mutant library (**Supplementary Fig. 1**).

When the number of positive clones exceeded 20, only 20 clones were selected randomly for sequencing. Next, we classified the 20 amino acids into 5 groups based on their physicochemical properties as described previously (8): [G, A, P, V, L, I, M – nonpolar and

aliphatic R groups], blue; [F, Y, W – aromatic R groups], purple; [S, T, C, N, Q – polar and uncharged R groups], green; [K, R, H – positively charged R groups], red; [D, E – negatively charged R groups], orange. Overall, 82 (50.9%), 48 (29.8%), 17 (10.6%), and 9 (5.6%) out of 161 residues could be replaced by a further 4, 3, 2, and 1 group of amino acid, respectively. Five FR residues (3.1%) of L23 (C), H36 (W), H86 (D), H104 (G), and H106 (G) were found to be replaced only with amino acids in the same group while maintaining its reactivity.

For comparison, we collected the amino acid sequences of 150 antigen-reactive chicken antibodies that we previously generated. Of these naturally-occurring chicken antibody FR, 1 (0.6%), 11 (6.8%), and 52 (32.3%) out of 161 FR residues were replaced by further 3, 2, and 1 group of amino acid, respectively. In case of 97 (60.3%) out of 161 residues, replacement with an amino acid from a different group was not found suggesting that naïve chicken antibody FR residues have limited sequence variations (**Fig. 1**).

Effect of mutating 3 consecutive framework residues

Next, we investigated the effect of mutating 3 consecutive FR residues on the reactivity of a chicken antibody. We hypothesized that this arrangement of mutations would affect the antibody structure more significantly than mutation on 3 sporadic residues. Therefore, we mutated H82 residue, which is located at the end of the beta chain, along with H82A residue and H82B residue in the loop of the chicken anti-phospho-tau antibody (PDB ID: 4GLR), which is one of two chicken antibodies with a published 3-dimensional structure (9). Based on our previous data, the following substitutions could be made: H82 residue could be replaced with A, V, L, F, W, S, K, R, and H, which belong to 4 groups; H82A residue could be replaced with G, A, P, V, L, I, W, S, T, C, N, K, R, and E, which are members of 5 groups; and H82B residue could be substituted with G, P, V, L, I, F, W, S, T, C, N, Q, K, R, H, and D, which are also members of 5 groups.

A total of 70 mutants were created by replacing H82A residue with a different amino acid from the same group, substituting H82 residue with A, V, L, F, and W, and altering H82B residue to G, P, V, L, F, W, S, T, C, N, Q, K, R, and D. From these combinatorial mutants, we obtained 19 positive clones (**Fig. 2**). When both H82 residue and H82B residue were replaced with residues from different groups, only 6.3% (1 out of 16 mutants) were positive. Substitution of either H82 residue or H82B residue with an amino acid from a different group produced 19.4% (7 out of 36 mutants) positive clones. However, when both residues were mutated to amino acids from the same group, 61.1% (11 out of 18 mutants) were positive.

Versatility of the framework residues in 2 other chicken scFvs

To determine whether other chicken antibodies can tolerate mutations as effectively as the PC24 antibody, we tested 2 other antigen-reactive chicken antibodies. To accomplish this, we arbitrarily categorized the FR residues of PC24 into 2 classes: flexible and non-flexible. When a residue harbored 4 or 5 amino acid groups, we defined it as a flexible residue. Otherwise, the residues were classified as non-flexible residues. We selected 8 flexible residues (L5 [T], L6 [Q], L7 [P], L43 [A], H13 [T], H16 [G], H43 [K], and H82A [N]) and 8 non-flexible residues (L23 [C], L88 [C], H22 [C], H36 [W], H90 [Y], H92 [C], H103 [W], and H106 [G]). Subsequently, we performed mutational analysis on these residues in the two chicken antibodies. All 8 flexible residues were also found to be flexible in both chicken antibodies. Furthermore, 3 and 1 non-flexible residues were found to be flexible in the PC17 and VCM2 antibodies, respectively (**Fig. 3**). With these data, we concluded that the redundant nature of FR residues toward mutations is widely shared among different chicken antibodies.

The Versatility of Chicken Antibody Framework Regions




Figure 1. Comparing the interchangeability of 161 FR residues in PC24 scFv with the natural mutations found in naturally-occurring chicken antibodies. Amino acids were classified into 5 group of amino acid based on their physicochemical properties: [GAPVLM], blue; [FYW], purple; [STCNQ], green; [KRH], red; [DE], orange. Positive rate was defined as the percentage of positive clones among 92 clones tested in phage enzyme immunoassay. The secondary structure of chicken antibody FR was adopted from the anti-phospho-tau antibody (PDB ID: 4GLR) (18). The sequences of 150 antigen-reactive chicken antibodies were collected, analyzed in the same manner, and presented as the natural mutation.




Figure 4. Versatility of FR residues in the IgG1 format. H78 (A) and L85 (B) FR residue mutants retaining antigen-reactivity in scFv format were expressed as IgG1. Their reactivity was evaluated in enzyme immunoassay using PSA-coated microtiter plates. The assays were performed in triplicate.

(H78: G, A, L, M, F, S, Q, and H; L85: G, A, P, L, M, F, Y, W, S, T, N, Q, K, R, D, and E). All of the mutant IgG1 molecules bound to PSA in a dose-dependent manner (**Fig. 4**). These data demonstrated that the versatility of the chicken framework is not confined within the context of a scFv but could be extended to IgG1 molecules.

DISCUSSION

The extent to which SHM impairs FR structure and function has not been studied thoroughly. Previously, one article studied the effect of the random mutations on V_H FR2 of a mouse monoclonal antibody (12). Only one out of 16 single residue mutants and 2 out of 8 double residue mutants had impaired antigen binding. When considering the fact that in the case of V_H CDR2, 5 out of 20 single residue mutants and 6 out of 7 double residue mutants lost their reactivity, the possibility was proposed that V_H FRs could accommodate many non-conservative changes. In this study, we showed that among 161 chicken antibody FR residues, 82 (50.9%), 48 (29.8%), 17 (10.6%), and 9 FR residues (5.6%) could be replaced by 4, 3, 2, and 1 group(s) of amino acids, individually, without significant loss of reactivity.

We evaluated the amino acid sequences of 150 naturally-occurring antigen-reactive chicken antibodies. Among these sequences, only 1 (0.6%), 11 (6.8%), and 52 (32.3%) FR residues were replaced by 3, 2, and 1 group(s) of amino acids, respectively. Furthermore, in a recent report analyzing the SHM of 1,269 unique naïve chicken V_H FR residues, only 4 (4.9%), 3 (3.7%), 26 (32.1%), and 27 (33.3%) residues out of 81 residues were replaced by 4, 3, 2, and 1 groups, respectively (5). These results clearly demonstrate that the chicken antibody FR could tolerate much more mutational diversity than that observed in natural chicken antibodies.

One of the main reasons why the mutations in FR residues of natural chicken antibodies covers only a fraction of possible mutations is that the sequence composition of FRs is intrinsically less mutable than that of CDRs in human and mouse V_H (11, 12). Moreover, the actual observed discrepancy in regional mutability in human V_H is much larger; although the mechanism contributing to regional mutability is still not clear (12). Because the frequency

of SHM in CDRs is limited to maintain the antigen reactivity *in vivo*, it is presumed that hypermutation in FRs would also be limited.

Another mechanism that could limit hypermutation of FRs is where mutations result in decreased stability and increased aggregation that would negatively affect secretion by B cells. The mutations in V_H FR2 caused secretion defects in 24% (8 out of 33), a frequency twice that of V_H CDR2 mutation (10). Interestingly, there are a few reports, in which stabilization of the variable region, especially in more unstable scFv or domain antibodies, could be achieved by introducing mutations in FRs. Substitution of hydrophobic FR residues with hydrophilic residues at surface-exposed positions increased the secretion of the scFv molecule in *E. coli* by reducing its aggregation potential (13). A single mutation in the V_H FR2 (E43G) of a human scFv increased its stability (14). Furthermore, the stability of a human single domain V_H antibody was dramatically increased by introducing 5 mutations on FRs (15). Because we tested the reactivity of the antibody in a phage-displayed format that should be less affected by the antibody stability, some unstable antibody clones could not be screened out.

Currently, we do not know if the versatility of FRs to accept mutations is a unique phenomenon of chicken antibody, which has only one functional V_H and V_λ gene. However, our rather unexpected finding that chicken antibody FR residues can accommodate much more diverse sequence variations than those found in nature, including mutations of consecutive residues, is of significant benefit to the field of antibody engineering. Chicken FRs could be de-immunized with great versatility by introducing mutations (16). In addition, many FR residues could be substituted with cysteine or artificial amino acids to provide a point of conjugation with chemical entities including drugs, toxins, peptides, aptamers, isotopes or enzymes. In this case, the intrinsic characteristics of the antibody including *in vivo* pharmacokinetics, distribution, and effector functions would be less affected than antibodies conjugated through the Fc domain, which interacts with Fc receptors and complements.

ACKNOWLEDGEMENTS

This research was supported by the Bio&Medical Technology Development Program through the National Research Foundation of Korea (NRF) funded by the Korea government (MSIP) (grant number: 2016M3A9B5941738).

SUPPLEMENTARY MATERIAL

Supplementary Figure 1

Phage enzyme immunoassay of L36 (A) and H82C (B) residue mutants.

[Click here to view](#)

REFERENCES

1. Cohen RM, Kleinstein SH, Louzoun Y. Somatic hypermutation targeting is influenced by location within the immunoglobulin V region. *Mol Immunol* 2011;48:1477-1483.

[PUBMED](#) | [CROSSREF](#)

2. Balint RF, Larrick JW. Antibody engineering by parsimonious mutagenesis. *Gene* 1993;137:109-118.
[PUBMED](#) | [CROSSREF](#)
3. Barbas CF 3rd, Burton DR. Selection and evolution of high-affinity human anti-viral antibodies. *Trends Biotechnol* 1996;14:230-234.
[PUBMED](#) | [CROSSREF](#)
4. Haidar JN, Yuan QA, Zeng L, Snively M, Luna X, Zhang H, Zhu W, Ludwig DL, Zhu Z. A universal combinatorial design of antibody framework to graft distinct CDR sequences: a bioinformatics approach. *Proteins* 2012;80:896-912.
[PUBMED](#) | [CROSSREF](#)
5. Wu L, Oficjalska K, Lambert M, Fennell BJ, Darmanin-Sheehan A, Ní Shúilleabháin D, Autin B, Cummins E, Tchistiakova L, Bloom L, et al. Fundamental characteristics of the immunoglobulin VH repertoire of chickens in comparison with those of humans, mice, and camelids. *J Immunol* 2012;188:322-333.
[PUBMED](#) | [CROSSREF](#)
6. Barbas CF. Phage Display: a Laboratory Manual. Cold Spring Harbor, NY; Cold Spring Harbor Laboratory Press; 2001.
7. Kabat EA, Wu TT, Perry HM, Gottesman KS, Foeller C. Sequences of Proteins of Immunological Interest, 5th ed. Bethesda, MD; U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health; 1991.
8. Kim HY, Tsai S, Lo SC, Wear DJ, Izadjoo MJ. Production and characterization of chimeric monoclonal antibodies against Burkholderia pseudomallei and B. mallei using the DHFR expression system. *PLoS One* 2011;6:e19867.
[PUBMED](#) | [CROSSREF](#)
9. Tsurushita N, Park M, Pakabunto K, Ong K, Avdalovic A, Fu H, Jia A, Vásquez M, Kumar S. Humanization of a chicken anti-IL-12 monoclonal antibody. *J Immunol Methods* 2004;295:9-19.
[PUBMED](#) | [CROSSREF](#)
10. Murray F, Darzentas N, Hadzidimitriou A, Tobin G, Boudjogra M, Scielzo C, Laoutaris N, Karlsson K, Baran-Marzszak F, Tsiftaris A, et al. Stereotyped patterns of somatic hypermutation in subsets of patients with chronic lymphocytic leukemia: implications for the role of antigen selection in leukemogenesis. *Blood* 2008;111:1524-1533.
[PUBMED](#) | [CROSSREF](#)
11. Wiens GD, Heldwein KA, Stenzel-Poore MP, Rittenberg MB. Somatic mutation in VH complementarity-determining region 2 and framework region 2: differential effects on antigen binding and Ig secretion. *J Immunol* 1997;159:1293-1302.
[PUBMED](#)
12. Wagner SD, Milstein C, Neuberger MS. Codon bias targets mutation. *Nature* 1995;376:732.
[PUBMED](#) | [CROSSREF](#)
13. Shapiro GS, Aviszus K, Ikle D, Wysocki LJ. Predicting regional mutability in antibody V genes based solely on di- and trinucleotide sequence composition. *J Immunol* 1999;163:259-268.
[PUBMED](#)
14. Yang J, Jia Z, Xiang J. Substitution of surface-exposed framework residues alters secretion of recombinant fusion protein Fv/tumor necrosis factor in *Escherichia coli*. *IUBMB Life* 1999;48:327-332.
[PUBMED](#) | [CROSSREF](#)
15. Rodríguez-Rodríguez ER, Ledezma-Candanoza LM, Contreras-Ferrat LG, Olamendi-Portugal T, Possani LD, Becerril B, Riaño-Umbarila L. A single mutation in framework 2 of the heavy variable domain improves the properties of a diabody and a related single-chain antibody. *J Mol Biol* 2012;423:337-350.
[PUBMED](#) | [CROSSREF](#)
16. Tanha J, Nguyen TD, Ng A, Ryan S, Ni F, Mackenzie R. Improving solubility and refolding efficiency of human V(H)s by a novel mutational approach. *Protein Eng Des Sel* 2006;19:503-509.
[PUBMED](#) | [CROSSREF](#)
17. Tangri S, LiCalsi C, Sidney J, Sette A. Rationally engineered proteins or antibodies with absent or reduced immunogenicity. *Curr Med Chem* 2002;9:2191-2199.
[PUBMED](#) | [CROSSREF](#)
18. Shih HH, Tu C, Cao W, Klein A, Ramsey R, Fennell BJ, Lambert M, Ní Shúilleabháin D, Autin B, Kouranova E, et al. An ultra-specific avian antibody to phosphorylated tau protein reveals a unique mechanism for phosphoepitope recognition. *J Biol Chem* 2012;287:44425-44434.
[PUBMED](#) | [CROSSREF](#)