

Mechanical Thrombectomy in Acute Stroke Patients with Moderate to Severe Pre-Stroke Disability

Marek Sykora,^{a,b} Patrik Michel,^c Davide Strambo,^c Stefan Krebs,^a Julia Ferrari,^a Alexandra Posekany,^{d,e} Dominika Mikšová,^d Konstantin Hermann,^b Thomas Gattringer,^{f,g} Elke Gizewski,^{h,i} Hannes Deutschmann,^g Christian Neumann,^j Wilfried Lang^{a,b}

^aDepartment of Neurology, St. John's Hospital, Vienna, Austria

^bMedical Faculty, Sigmund Freud University Vienna, Vienna, Austria

^cStroke Center, Neurology Service, Department of Clinical Neurosciences, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland

^dAustrian National Public Health Institute (Gesundheit Österreich GmbH)/The Austrian National Institute for Quality in Health Care (Bundesinstitut für Qualität im Gesundheitswesen, BIQG), Vienna, Austria

^eResearch Unit of Computational Statistics, University of Technology, Vienna, Austria

^fDepartment of Neurology, Medical University of Graz, Graz, Austria

^gDivision of Neuroradiology, Vascular and Interventional Radiology, Department of Radiology, Medical University of Graz, Graz, Austria

^hDepartment of Neuroradiology, Medical University of Innsbruck, Innsbruck, Austria

ⁱNeuroimaging Research Core Facility, Medical University of Innsbruck, Innsbruck, Austria

^jDepartment of Radiology, St. John's Hospital, Vienna, Austria

Background and Purpose Studies on mechanical thrombectomy (MT) in acute ischemic stroke (AIS) patients with preexisting disability are limited. We aimed to compare the outcomes of MT versus best medical treatment (BMT) in these patients.

Methods In the nationwide Austrian registry and Swiss monocentric registry, we identified 462 AIS patients with pre-stroke disability (modified Rankin Scale [mRS] score ≥ 3) and acute large vessel occlusion. The primary outcome was returning to pre-stroke mRS or better at 3 months. Secondary outcomes were early neurological improvement (National Institutes of Health Stroke Scale score improvement ≥ 8 at 24 to 48 hours), 3-month mortality, and symptomatic intracerebral hemorrhage (sICH). Multivariable regression models and propensity score matching (PSM) were used for statistical analyses.

Results Compared with the BMT group (n=175), the MT group (n=175) had younger age, more severe strokes, and lower pre-stroke mRS, but similar proportion of receiving intravenous thrombolysis. MT was associated with higher odds of returning to baseline mRS or better at 3 months (adjusted odds ratio [aOR], 2.5; 95% confidence interval [CI], 1.39 to 4.47), early neurological improvement (aOR, 2.62; 95% CI, 1.41 to 4.88), and lower risk of 3-month mortality (aOR, 0.29; 95% CI, 0.18 to 0.49). PSM analysis showed similar findings. MT was not associated with an increased risk of sICH (4.0% vs. 2.1% in all patients; 4.2% vs. 2.4% in the PSM cohort).

Conclusions MT in patients with pre-stroke mRS ≥ 3 might improve the 3-month outcomes and short-term neurological impairment, suggesting that pre-stroke disability alone should not be a reason to withhold MT, but that individual case-by-case decisions may be more appropriate.

Keywords Thrombectomy; Stroke; Pre-stroke; Disability; Handicapped; Outcome study

Correspondence: Marek Sykora
Department of Neurology, St. John's Hospital, Johannes von Gott Platz 1, 1020 Vienna, Austria
Tel: +43-1-21121-5183
Fax: +43-1-21121-3245
E-mail: marek.sykora@bbwien.at
<https://orcid.org/0000-0003-3508-2176>

Received: March 3, 2022

Revised: June 12, 2022

Accepted: June 13, 2022

Introduction

Several randomized trials have demonstrated the effectiveness of mechanical thrombectomy (MT) in acute ischemic stroke (AIS) patients with large vessel occlusion (LVO) and pre-morbid modified Rankin Scale (mRS) score <2 .¹⁻³ Based on this, the recent American (AHA/ASA) guidelines recommend MT for patients with previous disability up to mRS 1.⁴ While approximately 30% of patients with LVO have pre-stroke mRS ≥ 2 ,^{5,6} randomized data on efficacy and safety of MT in these patients are unavailable. Previous observational studies suggested a similar benefit of MT in patients with and without pre-stroke disability.⁷⁻⁹ These studies mainly compared MT-treated patients with moderate pre-stroke disability to those without any or only minimal previous disability.⁷⁻⁹ However, studies comparing MT with best medical treatment (BMT) including intravenous thrombolysis (IVT) in previously disabled patients, allowing examination of the effect size of MT, are scarce.¹⁰

Using retrospective data from two large European stroke registries, we aimed to investigate the efficacy and safety of MT as compared to medical treatment, including IVT, in consecutive stroke patients with moderate to severe pre-stroke disability.

Methods

Patients

The study population consisted of consecutive AIS patients registered in the Austrian Stroke Unit Registry (ASUR) and the Acute Stroke Registry and Analysis of Lausanne (ASTRAL) database. ASUR is a nationwide prospective registry of the Austrian stroke unit network founded by the Federal Ministry of Health. Anonymized data on baseline characteristics, risk factors and etiology, acute management, and functional outcome at discharge and at 3 months are registered for all patients admitted to one of currently 38 stroke units in Austria. Data collection and clinical ratings are performed by experienced stroke neurologists using standardized definitions of variables and scores. To ensure high data quality, immediate electronic data entry is performed. The web-based database includes online plausibility checks and a help function. Biannual educational meetings are also held to guarantee uniform data documentation. A detailed methodological description has been published previously.¹¹ On the other hand, ASTRAL is a single center-based cohort of all ischemic stroke patients admitted within 24 hours of stroke onset at the stroke center and/or intensive care unit of the Centre Hospitalier Universitaire Vaudois (CHUV).¹² This registry is approved by its institution as a clinical and research registry and follows institutional regulations. Data on pre-stroke condi-

tions, cerebrovascular risk factors, imaging and laboratory findings, stroke mechanism, acute management, functional outcome, and recurrences are collected up to 12 months. Data are collected online in a prespecified format by experienced and formed medical personnel. The web-based database also includes online plausibility checks and a help function. There is no uniform policy in ASUR-participating centers on whether patients with preexisting disability mRS ≥ 2 should be offered MT. At the CHUV, MT was offered standardly when pre-stroke mRS was ≤ 2 until 2017, and ≤ 3 since 2018. However, the final decision to offer MT remained at the discretion of the treating stroke unit physician at the time of stroke.

For the current analysis, we extracted from both databases those patients with AIS, treated between January 1, 2015 and December 31, 2019, aged ≥ 18 years, who had a pre-stroke mRS¹³ ≥ 3 , and potentially treatable LVO. Demographic and clinical data collection included age, sex, pre-stroke mRS, stroke severity, history of atrial fibrillation, arterial hypertension, and diabetes mellitus. Stroke severity on admission was measured using the National Institutes of Health Stroke Scale (NIHSS).¹⁴ LVO was categorized based on treatable occlusions in vessel segments M1 and M2 of the middle cerebral artery, extracranial and/or intracranial internal carotid artery, P1 segment of the posterior cerebral artery, and basilar artery. Patients were further categorized based on treatment received: MT, IVT, both, or none. Patients not receiving MT were considered as BMT.

Primary efficacy endpoint was defined with regard to pre-stroke disability as no or negative difference between pre-stroke mRS and mRS at follow-up at 3 months. Secondary efficacy endpoint was defined as a NIHSS difference of ≥ 8 points (early neurological improvement) between admission and at 24–48 hours. Further endpoint included mortality at 3 months. Safety endpoint was defined as a symptomatic intracerebral hemorrhage (sICH) according to the Thrombolysis with Alteplase 3 to 4.5 Hours after Acute Ischemic Stroke (ECASS3) study criteria.¹⁵

Statistics

We summarized the categorical variables as absolute numbers and percentages and continuous variables as means and standard deviation; quantitatively skewed variables were summarized as medians and interquartile range. Categorical variables were compared using the chi-square test, Fisher's exact test, or Kruskal-Wallis test, whereas continuous variables were compared using the Student's t-test or Mann-Whitney U test. Multivariable regression models were used to examine the adjusted effects of MT on outcomes. Standard demographic and clinical

factors known to influence outcome, including age, sex, admission NIHSS, pre-stroke mRS, hypertension, diabetes mellitus, atrial fibrillation, occlusion site, IVT use, and center, were used as covariates. We performed propensity score matching based on a logistic regression model controlling for the covariates as mentioned above, which matched the patients with most similar regression scores against each other, as implemented in the R package "MatchIt." Statistical analyses were performed using the IBM SPSS statistical software version 27 (IBM Co., Armonk, NY, USA), as well as the R statistical software version 4.0.2 (R Foundation for Statistical Computing, Vienna, Austria), with GLM function from the MASS package and "rpart" function from the "rpart" package.

Ethics

The ASUR is part of a government quality assessment program for nationwide stroke care and is financed by the Federal Ministry of Health. Data entry is mandatory for stroke units. All data are anonymized and centrally administered by the Gesundheit Österreich GmbH, which is the national research and planning institute for health care and a competence and fund-

ing center of health promotion. All scientific analyses included in this study were approved and supervised by a national academic review board. Based on this setting, informed consent for patients enrolled in the registry was waived. In ASTRAL, all data came from routine clinical and radiological management and were anonymized before analysis following the principles of the Swiss Human Research Ordinance. Therefore, there was no need for ethical commission approval or patient consent according to the Swiss Human Research Act and the applicable data protection legislation.

Data availability

Anonymized patient data are administered by the Gesundheit Österreich GmbH; inquiries on data availability are reviewed and granted by an academic review board. The anonymized data from ASTRAL supporting the findings of this study are available from the authors upon reasonable request and after signing a data transfer and use agreement.

Table 1. Demographic and clinical characteristics of the study population categorized by MT and best medical treatment

Characteristic	MT (n=175)	Medical (n=287)	P*
Age (yr)	79.4±12.2 (25–99)	85.2±8.8 (24–102)	<0.001
Female sex	109 (62.3)	187 (65.2)	0.550
Admission NIHSS	18 (1–36) (13–21)	15 (0–36) (8–21)	0.010
Pre-stroke mRS			0.010
3	136 (77.7)	185 (64.5)	
4	33 (18.9)	83 (28.9)	
5	6 (3.4)	19 (6.6)	
Hypertension	140 (80)	248 (86.4)	0.100
Diabetes mellitus	46 (27.2)	75 (26.2)	0.800
Atrial fibrillation	120 (71)	156 (58.2)	0.010
Occlusion site			0.070
M1 [†]	99 (56.6)	136 (47.4)	
M2 [†]	30 (17.1)	82 (28.6)	
ICA	26 (14.9)	35 (12.2)	
PCA P1	11 (6.3)	21 (7.3)	
BA	9 (5.1)	13 (4.5)	
Intravenous thrombolysis	75 (42.9)	100 (34.8)	0.100
ODT	96 (29–886) (52–121)	67.5 (24–544) (67–135)	0.900
DTN	36 (15–89) (27–59)	41 (20–88) (28–49)	0.200

Values are presented as mean±standard deviation (range), number (%), or median (range) (interquartile range).

MT, mechanical thrombectomy; NIHSS, National Institutes of Health Stroke Scale Score; mRS, modified Ranking Scale; ICA, internal carotid artery; PCA, posterior cerebral artery (P1–P1 segment); BA, basilar artery; ODT, onset-to-door time; DTN, door-to-needle time.

*P-value using the Student's t-test, chi-square test, Fisher exact test, and Kruskal-Wallis test, as appropriate; [†]M1 and M2, M1 and M2 segment of middle cerebral artery.

Results

Population

Within the study period, 2,111 patients with AIS were entered in the ASTRAL registry. Of those, 347 has pre-stroke mRS 3–5; 170 patients with potentially treatable LVO and pre-stroke mRS 3–5 were included in the study. In the same period, 8,664 AIS patients with pre-stroke mRS 3–5 were included in the ASUR registry. Of those, 292 with potentially treatable LVO and pre-stroke mRS 3–5 were included in the study. Overall, we included 462 patients in the analysis: 292 from the ASUR registry (111 MT, 181 BMT) and 170 from the ASTRAL registry (64 MT, 106 BMT). Altogether, 175 patients underwent MT and 287 received BMT. Patients undergoing MT were younger, had slightly higher admission stroke severity, and lower pre-stroke disability. Thrombolysis in cerebral infarction scores of 2b/3 in patients undergoing MT were achieved in 142 (81.1%) patients. IVT was administered in 75 (42.9%) patients in the MT group and 100 (34.8%) patients in the BMT group ($P=0.100$). Onset-to-door times and door-to-needle delays did not differ significantly between the two groups (Table 1).

Efficacy outcomes

At 3 months after stroke, 47 (26.9%) patients in the MT group returned to their pre-stroke mRS, compared to 58 (20.2%) in the BMT group ($P=0.100$) (Table 2). After adjusting for age, sex, admission NIHSS, pre-stroke mRS, hypertension, diabetes mellitus, atrial fibrillation, occlusion site, IVT, and center, MT was associated with higher odds of returning to the pre-stroke mRS at 3 months (adjusted odds ratio [aOR], 2.5; 95% confidence

interval [CI], 1.39 to 4.47; $P=0.002$) as compared to BMT (Table 3). Early neurological improvement NIHSS ≥ 8 points occurred in 50 (30.3%) patients undergoing MT compared to 26 (11.8%) receiving BMT ($P<0.001$) (Table 2). After adjustment, MT was associated with early neurological improvement NIHSS ≥ 8 points (aOR, 2.62; 95% CI, 1.41 to 4.88; $P=0.002$) (Table 3). Death at 3 months occurred in 76 (43.4%) patients in the MT group compared to 180 (62.7%) in BMT group ($P<0.001$) (Table 2). After adjustment, MT was inversely associated with death at 3 months (aOR, 0.29; 95% CI, 0.18 to 0.49; $P<0.001$) (Table 3).

Safety outcome

As defined by the ECASS3 criteria, sICH was present in seven (4%) patients in the MT group and in six (2.1%) patients in the BMT group ($P=0.250$) (Table 2).

Results after propensity score matching

Using propensity score matching, 168 MT and 168 BMT patients were selected. Except for age, there were no significant differences between the groups (Table 4).

Efficacy outcomes after propensity score matching

At 3 months after stroke, 47 (28.0%) patients in the MT group returned to their pre-stroke mRS, compared to 32 (19.0%) in the BMT group ($P=0.040$) (Table 5). Detailed distribution of mRS at 3 months categorized by MT versus BMT is shown in Figure 1. After adjustment, MT was associated with higher odds of returning to the pre-stroke mRS at 3 months (aOR, 2.54; 95% CI, 1.16 to 5.57; $P=0.020$) compared with the BMT

Table 2. Univariate analysis of the efficacy and safety of MT versus best medical treatment in stroke patients with pre-stroke mRS ≥ 3

Endpoint	MT (n=175)	Medical (n=287)	<i>P</i> *
Return to at least pre-stroke mRS at 3 months	47 (26.9)	58 (20.2)	0.100
Improvement NIHSS ≥ 8	50 (28.6)	26 (9.1)	<0.001
Mortality at 3 months	76 (43.4)	180 (62.7)	<0.001
Symptomatic ICH ECASS3	7 (4)	6 (2.1)	0.250

Values are presented as number (%).

MT, mechanical thrombectomy; mRS, modified Ranking Scale; NIHSS, National Institutes of Health Stroke Scale Score; ICH ECASS3, intracerebral hemorrhage according to ECASS3 definition.

**P*-value using the Mann-Whitney test, chi-square test, or Fisher exact test, as appropriate.

Table 3. Multivariate analysis of the efficacy of MT versus best medical treatment in stroke patients with pre-stroke mRS ≥ 3

Endpoint	Adjusted OR	95% CI	<i>P</i>
Return to at least pre-stroke mRS at 3 months	2.50	1.39–4.47	0.002
Early neurological improvement NIHSS ≥ 8 points	2.62	1.41–4.88	0.002
Mortality at 3 months	0.29	0.18–0.49	<0.001

MT, mechanical thrombectomy; OR, odds ratio; CI, confidence interval; mRS, modified Ranking Scale; NIHSS, National Institutes of Health Stroke Scale Score.

(Table 5). Early neurological improvement NIHSS ≥ 8 points occurred in 50 (29.8%) patients undergoing MT compared to 18 (10.7%) receiving BMT ($P < 0.001$) (Table 5). After adjustment, MT was associated with early neurological improvement NIHSS ≥ 8 points (aOR, 2.72; 95% CI, 1.26 to 5.92; $P = 0.010$) (Table 6). Death at 3 months occurred in 71 (42.3%) patients in the MT group compared to 107 (63.7%) in BMT group ($P < 0.001$) (Table 5). After adjustment, MT was inversely associated with death at 3 months (aOR, 0.27; 95% CI, 0.15 to 0.51; $P < 0.001$) (Table 6).

Safety outcome after propensity score matching

As defined by the ECASS3 criteria, sICH was present in seven (4.2%) patients in the MT group and in four (2.4%) patients in the BMT group ($P = 0.270$) (Table 5).

Discussion

In this retrospective multicenter observational study of AIS patients with pre-stroke disability, MT was significantly associated with higher odds of 3-month return to baseline mRS or bet-

Table 4. Demographic and clinical characteristics of the study population categorized by MT and best medical treatment after propensity score matching

Characteristic	MT (n=168)	Medical (n=168)	P*
Age (yr)	79±12.2 (25–97)	83±9.5 (24–99)	0.002
Female sex	108 (64.3)	105 (62.5)	0.820
Admission NIHSS	17 (1–36) (13–19)	17 (0–36) (13–21)	0.360
Pre-stroke mRS			0.360
3	132 (78.6)	123 (73.2)	
4	32 (19)	37 (22)	
5	4 (2.4)	8 (4.8)	
Hypertension	134 (79.8)	136 (81.5)	0.780
Diabetes mellitus	46 (27.4)	41 (24.4)	0.620
Atrial fibrillation	119 (70.8)	121 (72)	0.900
Occlusion site			0.750
M1 [†]	95 (56.5)	86 (51.2)	
M2 [†]	30 (17.9)	39 (23.2)	
ICA	25 (14.9)	23 (13.7)	
PCA P1	6 (3)	10 (6)	
BA	8 (4.8)	10 (6)	
Intravenous thrombolysis	72 (42.9)	65 (38.7)	0.500
ODT	70 (24–544) (53–125)	97 (35–886) (73–130)	0.940
DTN	40 (20–88) (27–56)	36 (15–89) (28–49)	0.230

Values are presented as mean±standard deviation (range), number (%), or median (range) (interquartile range).

MT, mechanical thrombectomy; NIHSS, National Institutes of Health Stroke Scale Score; mRS, modified Ranking Scale; ICA, internal carotid artery; PCA, posterior cerebral artery (P1–P1 segment); BA, basilar artery; ODT, onset-to-door time; DTN, door-to-needle time.

*P-value using the Student's t-test, chi-square test, Fisher exact, test and Kruskal-Wallis test, as appropriate; [†]M1 and M2, M1 and M2 segment of middle cerebral artery.

Table 5. Univariate analysis of the efficacy and safety of MT versus best medical treatment in stroke patients with pre-stroke mRS ≥ 3 after propensity score matching

Endpoint	MT (n=168)	Medical (n=168)	P*
Return to at least pre-stroke mRS at 3 months	47 (28.0)	32 (19.0)	0.040
Improvement NIHSS ≥ 8	50 (29.8)	18 (10.7)	<0.001
Mortality at 3 months	71 (42.3)	107 (63.7)	<0.001
Symptomatic ICH ECASS3	7 (4.2)	4 (2.4)	0.270

Values are presented as number (%).

MT, mechanical thrombectomy; mRS, modified Ranking Scale; NIHSS, National Institutes of Health Stroke Scale Score; ICH ECASS3, intracerebral hemorrhage according to ECASS3 definition.

*P-value using the Mann-Whitney test, chi-square test, or Fisher exact test, as appropriate.

ter, short-term NIHSS improvement, and lower odds of mortality, without increased risk for sICH.

Patients with relevant pre-stroke disability represent a special subgroup of stroke patients. Due to the preexisting handicap, usual outcome measures, such as excellent or favorable functional outcome defined as mRS 0–1 or mRS 0–2 at 3 months are not appropriate. For a patient with moderate pre-stroke disability, any mRS increase may represent a clinically unfavorable outcome. On the other hand, a stable mRS may represent a favorable outcome.

Our observation is consistent with that of previous studies. Several single center studies and registries have reported a probably beneficial effect of MT in pre-stroke dependent patients.^{7–9,16} However, only two studies used a control group. A monocentric study by Kastrup et al.¹⁷ (MT, n=142) showed that MT in patients with pre-stroke mRS 2–4 increased the odds of avoiding a poor outcome as compared with IVT controls. A multicenter study by Tanaka et al.¹⁰ suggested that in 175 patients with pre-stroke mRS 2–4, MT increased the odds of return to at least the pre-stroke mRS score as compared to medical management including IVT. Moreover, 28% of patients in the MT group returning to the pre-stroke mRS at 3 months and the aOR of 3.0 (95% CI, 1.4 to 6.6) for MT to return to the pre-stroke mRS at 3 months are fully consistent with our findings

(27% and aOR 2.5; 95% CI, 1.4 to 4.5). Interestingly, we observed, in both arms, higher mortality rates (43.4% and 62.7%) as compared to the similarly designed study by Tanaka et al.¹⁰ (17.7% and 26.8%). This may partially be explained by the fact that our population had cumulatively higher pre-stroke disability and more comorbidities and that we also included patients with posterior circulation stroke.

As in previous studies, the MT and medical treatment groups displayed baseline differences, mirroring probable treatment indication biases. Patients undergoing MT were younger, had a lower degree of pre-stroke disability, and had more favorable occlusion sites. Meanwhile, the medical treatment group had lower admission NIHSS and fewer atrial fibrillation, counterbalancing at least hypothetically the predicted better outcome in the MT group. Selection bias is a recognized limitation of the observational study design. Standard logistic regression models and propensity score matching have been used to account for the group's asymmetry, although it has to be considered a major limitation. The indication for MT was made individually by the treating physician, based on patient's cumulative comorbidity, cognitive impairment, frailty, center's preferences, and ischemic core volumes. Unfortunately, we were unable to include data on infarct core sizes (e.g., The Alberta Stroke Program Early CT Score [ASPECTS]¹⁸ scores) as the ASUR does not contain data on ASPECTS scores in patients not treated by MT. Altogether, these factors represent possible unmeasured confounders and account for additional limitations. Thus, our results should be interpreted with caution and with regard to the abovementioned limitations.

Meanwhile, the strength of our study is the rigorously collected large prospective multicenter dataset reflecting closely the real-world setting in hospitalized AIS patients.

Plausibly, patients with pre-stroke disability show increased rates of poor outcome and mortality despite treatment efforts. Nevertheless, MT seems to be clinically effective and prevent further functional deterioration and mortality in our series. Pre-stroke quality of life and regaining the previous quality of life may be crucial in decision-making processes. Pre-stroke mRS may not fully reflect the quality of life and adaptation to disability of individuals. Some patients with mRS 3, 4, or even 5 may




Figure 1. Distribution of modified Ranking Scale (mRS) score at 3 months for mechanical thrombectomy (MT) versus best medical treatment in stroke patients with pre-stroke mRS ≥ 3 after propensity score matching.

Table 6. Multivariate analysis of the efficacy of MT versus best medical treatment in stroke patients with pre-stroke mRS ≥ 3 after propensity score matching

Endpoint	Adjusted OR	95% CI	P*
Return to at least pre-stroke mRS at 3 months	2.54	1.16–5.57	0.020
Early neurological improvement NIHSS ≥ 8 points	2.72	1.26–5.92	0.010
Mortality at 3 months	0.27	0.15–0.51	<0.001

MT, mechanical thrombectomy; mRS, modified Ranking Scale; OR, odds ratio; CI, confidence interval; NIHSS, National Institutes of Health Stroke Scale Score.

eventually have, with supportive care, a meaningful quality of life. Therefore, pre-stroke mRS alone should not constitute a contraindication for endovascular treatment. Stroke-derived neurological impairment could potentially exacerbate suffering and decrease the quality of life. For patients with pre-stroke disability due to other serious health conditions or even life-limiting conditions, MT may represent a measure to decrease their sum of harm and alleviate further suffering. Basic ethical principles of beneficence and nonmaleficence make these considerations of offering stroke interventions in those previously disabled or even in a palliative setting justifiable.¹⁹

It would be very unlikely that this specific patient subgroup will be in focus of future randomized control trials examining the efficacy of MT. Therefore, based on results from this study and those of others,^{10,17} we suggest that MT should not be routinely withheld based solely on pre-stroke mRS. Careful and individualized decision-making, including quality of life considerations, ethical issues, and patients' declared or putative will, seems to be justified in this situation.

Conclusions

MT seems to be effective and safe in reducing neurological deficits and preventing further functional deterioration in patients with relevant pre-stroke disability. Careful and individual case-by-case decisions instead of unselected withholding of treatment based on pre-stroke mRS may be appropriate.

Disclosure

The authors have no financial conflicts of interest.

Acknowledgments

Patrik Michel has received, within the last 2 years, research grants from the Swiss National Science Foundation and the Swiss Heart Foundation. The other authors do not have any relevant disclosures.

We thank all Austrian Stroke Unit Registry collaborators.

References

1. Goyal M, Menon BK, van Zwam WH, Dippel DW, Mitchell PJ, Demchuk AM, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016;387:1723-1731.
2. Albers GW, Marks MP, Kemp S, Christensen S, Tsai JP, Orte-

- ga-Gutierrez S, et al. Thrombectomy for stroke at 6 to 16 hours with selection by perfusion imaging. *N Engl J Med* 2018;378:708-718.
3. Nogueira RG, Jadhav AP, Haussen DC, Bonafe A, Budzik RF, Bhuva P, et al. Thrombectomy 6 to 24 hours after stroke with a mismatch between deficit and infarct. *N Engl J Med* 2018;378:11-21.
4. Powers WJ, Rabinstein AA, Ackerson T, Adeoye OM, Bambakidis NC, Becker K, et al. Guidelines for the early management of patients with acute ischemic stroke: 2019 update to the 2018 guidelines for the early management of acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2019;50:e344-e418.
5. Karlinski M, Kobayashi A, Czlonkowska A, Mikulik R, Vaclavik D, Brozman M, et al. Role of preexisting disability in patients treated with intravenous thrombolysis for ischemic stroke. *Stroke* 2014;45:770-775.
6. Ganesh A, Luengo-Fernandez R, Pendlebury ST, Rothwell PM. Long-term consequences of worsened poststroke status in patients with premorbid disability. *Stroke* 2018;49:2430-2436.
7. Goldhoorn RB, Verhagen M, Dippel DW, van der Lugt A, Lingsma HF, Roos YB, et al. Safety and outcome of endovascular treatment in prestroke-dependent patients. *Stroke* 2018;49:2406-2414.
8. Salwi S, Cutting S, Salgado AD, Espallat K, Fusco MR, Froehler MT, et al. Mechanical thrombectomy in patients with ischemic stroke with prestroke disability. *Stroke* 2020;51:1539-1545.
9. de Havenon A, Castonguay A, Nogueira R, Nguyen TN, English J, Satti SR, et al. Prestroke disability and outcome after thrombectomy for emergent anterior circulation large vessel occlusion stroke. *Neurology* 2021;97:e1914-e1919.
10. Tanaka K, Yamagami H, Yoshimoto T, Uchida K, Morimoto T, Toyoda K, et al. Endovascular therapy for acute ischemic stroke in patients with prestroke disability. *J Am Heart Assoc* 2021;10:e020783.
11. Steiner MM, Brainin M; Austrian Stroke Registry for Acute Stroke Units. The quality of acute stroke units on a nation-wide level: the Austrian Stroke Registry for acute stroke units. *Eur J Neurol* 2003;10:353-360.
12. Michel P, Odier C, Rutgers M, Reichhart M, Maeder P, Meuli R, et al. The Acute STroke Registry and Analysis of Lausanne (ASTRAL): design and baseline analysis of an ischemic stroke registry including acute multimodal imaging. *Stroke* 2010;41:2491-2498.
13. Bonita R, Beaglehole R. Recovery of motor function after

- stroke. *Stroke* 1988;19:1497–1500.
14. Brott T, Adams HP Jr, Olinger CP, Marler JR, Barsan WG, Biller J, et al. Measurements of acute cerebral infarction: a clinical examination scale. *Stroke* 1989;20:864–870.
 15. Hacke W, Kaste M, Bluhmki E, Brozman M, Dávalos A, Guidetti D, et al. Thrombolysis with alteplase 3 to 4.5 hours after acute ischemic stroke. *N Engl J Med* 2008;359:1317–1329.
 16. Seker F, Pfaff J, Schönenberger S, Herweh C, Nagel S, Ringleb PA, et al. Clinical outcome after thrombectomy in patients with stroke with premorbid modified Rankin Scale scores of 3 and 4: a cohort study with 136 patients. *AJNR Am J Neuroradiol* 2019;40:283–286.
 17. Kastrup A, Roth C, Politi M, Alexandrou M, Hildebrandt H, Schröter A, et al. Endovascular therapy vs. thrombolysis in pre-stroke dependent patients with large vessel occlusions within the anterior circulation. *Front Neurol* 2021;12:666596.
 18. Barber PA, Demchuk AM, Zhang J, Buchan AM. Validity and reliability of a quantitative computed tomography score in predicting outcome of hyperacute stroke before thrombolytic therapy. ASPECTS Study Group. Alberta Stroke Programme Early CT Score. *Lancet* 2000;355:1670–1674.
 19. Young MJ, Regenhardt RW, Sokol LL, Leslie-Mazwi TM. When should neuroendovascular care for patients with acute stroke be palliative? *AMA J Ethics* 2021;23:E783–E793.