

## Original Article



# Definitive treatment of primary vaginal cancer with radiotherapy: multi-institutional retrospective study of the Korean Radiation Oncology Group (KROG 12-09)

Ji Hyun Chang,<sup>1,\*</sup> Won Il Jang,<sup>2,\*</sup> Yong Bae Kim,<sup>3</sup> Jin Hee Kim,<sup>4</sup> Young Seok Kim,<sup>5</sup>  
Yeon Sil Kim,<sup>1</sup> Won Park,<sup>6</sup> Juree Kim,<sup>7</sup> Won Sup Yoon,<sup>8</sup> Joo-Young Kim,<sup>9</sup>  
Hak Jae Kim<sup>10</sup>

## OPEN ACCESS

**Received:** Sep 2, 2015

**Revised:** Sep 29, 2015

**Accepted:** Oct 7, 2015

### Correspondence to

**Hak Jae Kim**

Department of Radiation Oncology, Seoul  
National University College of Medicine, 101  
Daehak-ro, Jongno-gu, Seoul 03080, Korea.  
E-mail: khjae@snu.ac.kr

\*The first two authors contributed equally to  
this study.

**Copyright** © 2016. Asian Society of  
Gynecologic Oncology, Korean Society of  
Gynecologic Oncology

This is an Open Access article distributed under  
the terms of the Creative Commons Attribution  
Non-Commercial License ([http://  
creativecommons.org/licenses/by-nc/4.0/](http://creativecommons.org/licenses/by-nc/4.0/))  
which permits unrestricted non-commercial  
use, distribution, and reproduction in any  
medium, provided the original work is properly  
cited.

### ORCID

Ji Hyun Chang  
<http://orcid.org/0000-0001-5921-5522>  
Won Il Jang  
<http://orcid.org/0000-0002-5279-3087>  
Yong Bae Kim  
<http://orcid.org/0000-0003-4884-2330>  
Yeon Sil Kim  
<http://orcid.org/0000-0003-1291-0821>

<sup>1</sup>Department of Radiation Oncology, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Korea

<sup>2</sup>Department of Radiation Oncology, Korea Institute of Radiological & Medical Science, Seoul, Korea

<sup>3</sup>Department of Radiation Oncology, Yonsei Cancer Center, Yonsei University College of Medicine, Seoul, Korea

<sup>4</sup>Department of Radiation Oncology, Keimyung University Dongsan Medical Center, Keimyung University School of Medicine, Daegu, Korea

<sup>5</sup>Department of Radiation Oncology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

<sup>6</sup>Department of Radiation Oncology, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

<sup>7</sup>Department of Radiation Oncology, Cheil General Hospital & Women's Healthcare Center, Dankook University College of Medicine, Seoul, Korea

<sup>8</sup>Department of Radiation Oncology, Korea University Ansan Hospital, Korea University College of Medicine, Ansan, Korea

<sup>9</sup>Center for Uterine Cancer, Research Institute and Hospital, National Cancer Center, Goyang, Korea

<sup>10</sup>Department of Radiation Oncology, Seoul National University College of Medicine, Seoul, Korea

## ABSTRACT

**Objective:** To assess the outcome of the treatment of primary vaginal cancer using definitive radiotherapy (RT) and to evaluate the prognostic factors of survival.

**Methods:** The medical records of nine institutions were retrospectively reviewed to find the patients with vaginal cancer treated with definitive RT with or without chemotherapy. A total of 138 patients met the inclusion criteria. None had undergone curative excision.

**Results:** The median follow-up time of the survivors was 77.6 months and the median survival time was 46.9 months. The 5-year overall survival, cancer-specific survival (CSS), and progression-free survival (PFS) rates were 68%, 80%, and 68.7%, respectively. In the survival analysis, the multivariate analysis showed that a lower the International Federation of Gynecology and Obstetrics (FIGO) stage and prior hysterectomy were favorable prognostic factors of CSS, and a lower FIGO stage and diagnosed prior to year 2000 were favorable prognostic factors of PFS. In the subgroup analysis of the patients with available human papillomavirus (HPV) results (n=27), no statistically significant relationship between the HPV status and recurrence or survival was found. Grade 3 or 4 acute and late toxicity were present in 16 and 9 patients, respectively. The FIGO stage and the tumor size were predictors of severe late toxicity.

Won Sup Yoon

<http://orcid.org/0000-0002-6098-1993>

Hak Jae Kim

<http://orcid.org/0000-0003-3602-2263>

#### Conflict of Interest

No potential conflict of interest relevant to this article was reported.

**Conclusion:** The data clearly showed that a higher FIGO stage was correlated with a worse survival outcome and higher severe late toxicity. Therefore, precise RT and careful observation are crucial in advanced vaginal cancer. In this study, the HPV status was not related to the survival outcome, but its further investigation is needed.

**Keywords:** Radiotherapy; Stage; Vaginal Neoplasms

## INTRODUCTION

Primary vaginal cancer is a rare malignancy, accounting for 1% to 2% of all gynecologic malignancies. According to the annual report on cancer statistics in South Korea in 2012, the annual incidence of vaginal cancer accounts for 0.1% of the all-cancer registry, with a 5-year incidence of approximately 240 patients, which is lower than the reported data [1]. Therefore, most studies of primary vaginal cancer are small due to the rarity of the disease.

To treat vaginal cancer, surgical options can be considered for highly selected small and localized early-stage patients. Radiotherapy (RT) with or without chemotherapy is a treatment of choice for most primary vaginal cancer patients. Additional chemotherapy is often used by extrapolating cervical cancer data, yet no randomized trial has compared RT alone and concurrent chemoradiotherapy (CCRT). Recently, the National Cancer Data Base of the United States has reported the results of the study of the largest population of primary vaginal cancer patients and provided survival data and prognostic factors of data for over 8,000 patients, which revealed the superiority of CCRT over RT alone in the treatment of primary vaginal cancer [2]. However, detailed data were significantly missing, such as the purpose of chemotherapy or RT (inductive or definitive) and the toxicity profile. Moreover, though this study was designed to deal with definitive RT, patients who had undergone lymphadenectomy were included.

To overcome the limitations of previous studies of primary vaginal cancer due to the rarity of the disease, the Korean Radiation Oncology Group (KROG) performed a multi-institutional retrospective study, the results of which are reported herein.

## MATERIALS AND METHODS

### 1. Patients and tumor characteristics

This multicenter retrospective study was conducted on primary vaginal cancer patients treated with definitive RT with or without chemotherapy from June 1976 to November 2011 in nine institutions approved by KROG (KROG, 12-09). Patients with distant metastasis were excluded, except patients who had para-aortic lymph node metastasis, as those could be covered by the extended pelvic RT field. Patients with a history of curative excision, who were in the Tis stage, or who had a histology of vaginal melanoma were also excluded. Patients were diagnosed as vaginal cancer in case where there was no evidence of recurrence of cervical/endometrial cancer for more than 10 years or when histologic subtypes were different from the two diseases. This study was approved by the Institutional Review Board of each of the included institutions.

**Table 1.** Patient, tumor, and treatment characteristics

Characteristic	No. of patients (%)
Age (yr), median (range)	61 (30–84)
Previous history of cervical cancer	
Yes	7 (5)
No	131 (95)
Previous history of CIS	
Yes	13 (9)
No	125 (91)
Prior hysterectomy	
Yes	55 (40)
No	83 (60)
Histology	
SqCC	119 (86)
Adenocarcinoma	6 (4)
Others	13 (10)
Size (cm)	
≤4	105 (76)
>4	33 (24)
Year of initial diagnosis	
<2000	57 (41)
≥2000	81 (59)
FIGO stage	
I	59 (43)
II	31 (22)
III	26 (19)
IVA	19 (14)
IVB (with para-aortic metastasis)	3 (2)
HPV status	
Positive	17 (12)
Negative	10 (7)
Unknown	111 (81)
Treatment	
RT alone	81 (59)
CCRT	46 (33)
Neoadjuvant chemotherapy+RT or CCRT	11 (8)
RT technique	
EBRT alone	22 (16)
Brachytherapy alone	4 (3)
EBRT+brachytherapy	112 (81)
Total dose (Gy)	
<80	87 (63)
≥80	51 (37)

CCRT, chemoradiotherapy; CIS, carcinoma *in situ*; EBRT, external beam radiotherapy; FIGO, International Federation of Gynecology and Obstetrics; HPV, human papillomavirus; RT, radiotherapy; SqCC, squamous cell carcinoma.

A total of 152 patients in nine institutions who satisfied the inclusion criteria were identified. Of these, 11 patients with incomplete RT or an RT dose ≤ 50.4 Gy to the vagina were excluded. Three patients whose follow-up period was less than four months were also excluded. Finally, 138 patients were evaluated for the survival analysis. The patient, tumor, and treatment characteristics are shown in **Table 1**. The median age was 61 years (range, 30 to 84 years). Seven patients had a history of cervical cancer, and 13 had carcinoma *in situ* (CIS) before vaginal cancer. Fifty-five patients had undergone hysterectomy, 22 of them for malignancy (seven for cervical ca, one for endometrial cancer, one for malignant mixed Müllerian tumor, and 13

**Table 2.** Patients characteristics according to the treatment (n=127)

Characteristic	RT alone (n=81)	CCRT (n=46)	p-value
Age (yr)			0.065
<60	33 (41)	27 (59)	
≥60	48 (59)	19 (41)	
History of cervical cancer or CIS			0.010
Yes	17 (21)	2 (4)	
No	64 (79)	44 (96)	
Prior hysterectomy			0.852
Yes	34 (42)	18 (39)	
No	47 (58)	28 (61)	
Histology			0.056
SqCC	74 (91)	36 (78)	
Non-SqCC	7 (9)	10 (22)	
Size (cm)			0.284
≤4	64 (79)	32 (70)	
>4	17 (21)	14 (30)	
Year of initial diagnosis			<0.001
<2000	46 (57)	4 (9)	
≥2000	35 (43)	42 (91)	
FIGO stage			0.124
I–II	56 (69)	25 (54)	
III–IVB	25 (31)	21 (46)	
RT technique			0.300
EBRT alone	13 (16)	7 (15)	
Brachytherapy alone	4 (5)	0	
EBRT+brachytherapy	64 (79)	39 (85)	
Total dose (Gy)			0.342
<80	49 (61)	32 (70)	
≥80	32 (39)	14 (30)	

Values are presented as number (%).

CCRT, concurrent chemoradiotherapy; CIS, carcinoma in situ; EBRT, external beam radiotherapy; FIGO, International Federation of Gynecology and Obstetrics; RT, radiotherapy; SqCC, squamous cell carcinoma.

for CIS) and 33 for various benign diseases such as uterine fibroids or pelvic inflammatory disease. One patient had a pelvic RT history because of cervical cancer. All the patients underwent a thorough gynecological examination at the time of their diagnosis, with chest X-ray for 90% of the patients, cystoscopy for 59%, colonoscopy for 70%, a pelvic computed tomography (CT) scan for 54%, a pelvic magnetic resonance imaging scan for 62%, and a positron emission tomography-CT scan for 28%. Generally, the size was measured by imaging studies if available, or by gynecological exams.

Eighty-six percent of the patients were diagnosed with squamous cell carcinoma (SqCC), and 24% had tumors larger than 4 cm. The human papillomavirus (HPV) status of the tumor was available in 27 patients, and 17 (63%) of those were HPV-positive. The HPV-positive tumors were all SqCC.

When the patients who had undergone RT alone were compared with the CCRT group (n=127), the RT alone group showed a stronger history of cervical cancer or CIS, and more patients received the treatment before the year 2000 (Table 2). Moreover, in the comparison of the patients treated before the year 2000 and after 2000, there were more older patients, more large tumors, and more CCRT procedures performed on the patients treated after 2000, whose total dose was also lower (Table 3).

**Table 3.** Patients characteristics according to the year of diagnosis

Characteristic	Before 2000 (n=57)	After 2000 (n=81)	p-value
Age (yr)			0.010
<60	34 (60)	31 (38)	
≥60	23 (40)	50 (62)	
History of cervical cancer or CIS			0.628
Yes	7 (12)	13 (16)	
No	50 (88)	68 (84)	
Previous history of hysterectomy			0.113
Yes	18 (32)	37 (46)	
No	39 (68)	44 (54)	
Histology			0.455
SqCC	51 (90)	68 (84)	
Non-SqCC	6 (10)	13 (16)	
Size (cm)			0.026
≤4	49 (86)	56 (69)	
>4	8 (14)	25 (31)	
FIGO stage			0.587
I–II	39 (68)	51 (63)	
III–IVB	18 (32)	30 (37)	
Treatment			<0.001
RT alone	46 (81)	35 (43)	
CCRT	4 (7)	42 (52)	
ICT+ other Tx	7 (12)	4 (5)	
RT technique			0.111
EBRT alone	5 (9)	17 (21)	
Brachytherapy alone	1 (2)	3 (4)	
EBRT+brachytherapy	51 (89)	61 (75)	
Total dose (Gy)			0.007
<80	28 (49)	59 (73)	
≥80	29 (51)	22 (27)	

Values are presented as number (%).

CCRT, chemoradiotherapy; CIS, carcinoma in situ; EBRT, external beam radiotherapy; FIGO, International Federation of Gynecology and Obstetrics; ICT, induction chemotherapy; SqCC, squamous cell carcinoma; RT, radiotherapy; Tx, treatment.

## 2. Treatment

Eighty-one percent of the patients had undergone both external beam radiotherapy (EBRT) and brachytherapy; 16%, EBRT alone; and 3%, brachytherapy alone. The pelvic RT field of the 134 patients who had undergone EBRT basically included their entire vagina and their pelvic lymph nodes, and 53 of them were selected to undergo extended field pelvic RT that included their inguinal lymph node or para-aortic lymph node regions. The median dose for the pelvis was 46 Gy (range, 39.6 to 54 Gy). Boost doses were given to 49 patients for gross primary tumor, the parametrial region, or lymph nodes. Among the node-positive patients (n=34), 27 underwent boost irradiation of their grossly enlarged lymph nodes.

For the intracavitary brachytherapy, low-dose rate from a cesium-137 radioactive source (median dose, 40.2 Gy; range, 30 to 65.2 Gy) was used for 35 patients in two hospitals, and high-dose rate via iridium-192 (median dose, 30 Gy; range, 10 to 50 Gy), for 80 patients in seven hospitals. Vaginal cylinders or Fletcher-Suit-Delclos colpostats were used for brachytherapy. The dose prescription point was 0.5 or 1.0 cm in depth to the surface of the vaginal mucosa to cover the gross residual tumor after EBRT, which was made upon clinician's decision.

### 3. Statistical analysis

The survival durations were calculated from the first day of the initial treatment. Overall survival (OS) and cancer-specific survival (CSS) were defined as the time to death from any cause and the time to death from vaginal cancer. Loco-regional recurrence-free survival (LRFS) and progression-free survival (PFS) were defined as the time to the loco-regional and overall recurrence, respectively. The Kaplan-Meier survival graphs and statistical differences were presented using the log-rank test. A Cox proportional hazard model was used for the multivariate analysis, and the variables showed a p-value <0.2 in the log-rank test. To find predictors of grade 3 or 4 late toxicity, logistic regression analysis was used. All the statistical analyses were carried out using SPSS ver. 22 (IBM Co., Armonk, NY, USA). A p<0.05 was considered statistically significant.

## RESULTS

### 1. Treatment outcome

The median follow-up time of the survivors was 77.6 months (range, 4.4 to 426.1 months). The median survival time was 46.9 months (range, 4.3 to 426.1 months). Twenty-seven patients (20%) died from vaginal cancer during the follow-up period, and 25 (18%) died from other causes. The 5-year OS, CSS, and PFS rates were 68%, 80%, and 68.7%, respectively. The 5-year LRFS rate was 74%. The 5-year OS rate in stage I of the International Federation of Gynecology and Obstetrics (FIGO) staging system was 75.4%; in stage II, 68.8%; in stage III, 68%; and in stage IV, 47%. The 5-year CSS rate in stage I was 85%; in stage II, 83.6%; in stage III, 74.5%; and in stage IV, 69.2%. The 5-year PFS rate in stage I was 78.9%; in stage II, 66%; in stage III, 52.5%; and in stage IV, 65.2%.

The prognostic factors of the OS and the CSS, as well as of the 5-year survival rate, significantly differed. In the multivariate analysis, an age above 60 and treatment after the year 2000 were statistically significant for the OS. In contrast, the FIGO stage and prior hysterectomy were associated with the CSS (Table 4). In the same analysis, the year of the initial diagnosis and the FIGO stage were found to be related to the PFS, and the year of the initial diagnosis, to the LRFS.

Distant metastasis occurred in 19 patients (13.7%). The most common sites of the metastases were the lung (in five patients) and the para-aortic lymph node (in five patients), followed by the supraclavicular lymph node (in four patients). Two patients each had liver, cervical lymph node, peritoneum, and brain metastases, and one patient had bone, abdominal wall, and mediastinal lymph nodes.

### 2. Subgroup analysis of the HPV-tested patients

In the patients whose tumor HPV status was available (n=27), no statistically significant difference was found in all the outcomes. However, the patients with HPV-positive tumors all showed trends toward better survival outcomes, especially in the LRFS (5-year OS: 68.8% for the HPV-positive patients vs. 44.4% for the HPV-negative patients, p=0.406; 5-year CSS: 82.4% vs. 74.1%, p=0.748; 5-year PFS: 81.6% vs. 64.8%, p=0.386; and 5-year LRFS: 93.3% vs. 77.8%, p=0.219).

**Table 4.** Prognostic factors for survival outcome of primary vaginal cancer

Factor	No. of patients (%)	5-yr LRFS			5-yr PFS			5-yr CSS			5-yr OS		
		5-yr survival rate	Univariate*	Multivariate†	5-yr survival rate	Univariate*	Multivariate†	5-yr survival rate	Univariate*	Multivariate†	5-yr survival rate	Univariate*	Multivariate†
Age (yr)													
<60	65 (47)	76.5	0.430		70.3	0.663		82.2	0.309		77.0	0.02	0.009
≥60	73 (53)	71.3			66.8			78.0			57.4		
History of cervical cancer or CIS													
Yes	20 (14)	94.7	0.081	0.226	94.7	0.035	0.167	93.8	0.226		78.7	0.178	0.142
No	118 (86)	70.5			64.3			77.7			66.2		
Prior hysterectomy													
Yes	55 (40)	80.8	0.117	0.267	77.7	0.032	0.096	87.7	0.105	0.036	68.0	0.591	
No	83 (60)	69.3			62.2			74.8			67.4		
Histology													
SqCC	119 (86)	74.6	0.894		69.2	0.811		82.2	0.125	0.252	71.0	0.202	
Non-SqCC	19 (14)	66.8			60.2			64.3			46.1		
Size (cm)													
≤4	105 (76)	76.7	0.04	0.191	72.1	0.029	0.169	81.7	0.109	0.620	71.0	0.112	0.704
>4	33 (24)	66.0			57.7			75.1			57.4		
Year of initial diagnosis													
<2000	57 (41)	83.0	0.051	0.048	79.3	0.035	0.041	88.2	0.027	0.076	88.2	<0.001	0.014
≥2000	81 (59)	66.9			60.1			73.3			53.6		
FIGO stage													
I-II	90 (65)	76.3	0.185	0.204	74.4	0.009	0.01	84.4	0.015	0.034	73.0	0.037	0.140
III-IVB	48 (35)	69.9			58.1			71.8			59.0		
Treatment													
RT alone	81 (59)	78.7			75.1			88.6			79.7		
CCRT	46 (33)	63.7	0.394		53.7	0.153	0.966	64.8	0.021	0.457	47.8	0.010	0.239
ICT+RT or CCRT	11 (8)	78.8			80.0			77.1			68.6		
RT technique													
EBRT alone	22 (16)	70.8			71.1			84.1			68.2		
Brachytherapy alone	4 (3)	75.0	0.751		75.0	0.948		75.0	0.970		37.5	0.820	
EBRT+brachytherapy	112 (81)	74.9			68.3			79.5			69.0		
Total dose (Gy)													
<80	87 (63)	76.1	0.279		71.9	0.252		79.1	0.379		64.4	0.269	
≥80	51 (37)	69.1			62.5			81.3			73.3		

CCRT, concurrent chemoradiotherapy; CIS, carcinoma in situ; CSS, cancer-specific survival; EBRT, external beam radiotherapy; FIGO, International Federation of Gynecology and Obstetrics; ICT, induction chemotherapy; LRFS, loco-regional recurrence-free survival; OS, overall survival; PFS, progression free survival; RT, radiotherapy; SqCC, squamous cell carcinoma. \*Log-rank test. †Cox proportional hazard model.

**Table 5.** Toxicity profiles in patients treated for primary vaginal cancer

Variable	Acute				Chronic			
	Grade 1	Grade 2	Grade 3	Grade 4	Grade 1	Grade 2	Grade 3	Grade 4
Skin	18 (13)	17 (12)	10 (7)	2 (1)	3 (2)	4 (3)	2 (1)	1 (1)
Gastrointestinal	34 (25)	22 (16)	1 (1)	0	5 (4)	10 (7)	7 (5)	2 (1)
Genitourinary	21 (15)	12 (9)	3 (2)	0	4 (3)	5 (4)	6 (4)	2 (1)
Others	7 (5)	4 (3)	0	0	1 (1)	8 (6)	0	0

Values are presented as number (%).

### 3. Complications

The toxicity profiles of the treated patients are listed in **Table 5** and were recommended to be graded according to the Common Terminology Criteria for Adverse Events ver. 4.0. Sixteen patients (12%) had grade 3 or 4 acute toxicity; 12 (9%), grade 3 or 4 acute skin toxicity; one (1%), grade 3 acute gastrointestinal (GI) toxicity; and three (2%), grade 3 acute genitourinary toxicity. Twelve patients (8%) showed late toxicity. Grade 3 or 4 late GI toxicity occurred in nine patients (7%), and grade 3 or 4 genitourinary complications, in eight patients (6%). Grade 3 or 4 late skin toxicity developed in three patients (2%). Prognostic factors for severe toxicity (age, prior hysterectomy, histology, tumor size, year of initial diagnosis, FIGO stage, treatment, RT technique, and total dose) were evaluated by logistic regression. The result showed that a higher FIGO stage (I or II vs. III or IV) (odds ratio [OR], 4.153; 95% CI, 1.145 to 15.056;  $p=0.03$ ), and an over 4 cm tumor size (OR, 4.044; 95% CI, 1.189 to 13.756;  $p=0.025$ ) were significantly related to the risk of grade 3 or 4 late toxicity.

## DISCUSSION

This study presents the results of treatments of primary vaginal cancer with definitive RT in multiple institutions in Korea for 35 years. The presented CSS outcomes were comparable to or higher than those in recently reported data [3-5]. The results of this study showed that the prognostic factors of the OS (the age and the year of the initial diagnosis) and the CSS (prior hysterectomy and the FIGO stage) significantly differed. Because many deaths were unrelated to vaginal cancer, the CSS rather than the OS represented the survival outcome after the treatment of primary vaginal cancer. Therefore, the authors focused on the CSS rather than the OS in the interpretation of the results.

In this study, the FIGO stage was strongly associated with the CSS and the PFS in the multivariate analysis, as expected [3,6]. Additionally, prior hysterectomy was related to better CSS. It has been acknowledged that a number of vaginal neoplasms develop after hysterectomy [7,8]. In the recent study of Daling et al. [9], they also found that prior hysterectomy was a risk factor of vaginal cancer in women with no history of anogenital cancer. The data are consistent with the results of Chyle et al. [10] and Tran et al. [4], which showed the relationship between prior hysterectomy and the vaginal cancer survival outcome. Although the relationship was inconclusive, the researchers attributed the better survival in the prior hysterectomy group to early diagnosis and the smaller tumors due to the absence of the uterus and the lower amount of vaginal substrates.

Although the tumor size is known as an important prognostic factor of the survival of vaginal cancer patients, its accurate measurement is difficult without surgery due to the anatomical difficulties [11]. Therefore, it is difficult to apply an accurate tumor size to the prediction of



the survival outcome in a clinical setting without surgical resection in the clinical setting. The data from the univariate analysis in this study also showed better disease control in small tumors ( $\leq 4$  cm), as in other studies [2,3], but they were insignificant in the multivariate analysis in this study.

The results of this study that were difficult to interpret were that the patients treated after the year 2000 had much worse LRFS, PFS, and OS rates and tended to have worse CSS rates. Although such results remained significant after they were adjusted based on other factors in the multivariate analysis, increased surgery in the treatment of early-stage vaginal cancer and the use of neo-adjuvant chemotherapy were speculated to have facilitated curative resection in the previously unresectable cases. The reported results of the National Cancer Data Base and Surveillance, Epidemiology, and End Results data analysis after the 1990s showed better survival outcomes with surgery than with definitive RT, but also inherent selection biases [11,12]. Therefore, there could have been trends toward increased surgical resection before considering definitive RT in early-stage vaginal cancer. As seen in **Table 3**, there were older patients and patients with larger primary tumors in the group treated after the year 2000. Several surgical series showed biases in the selection of surgical candidates and more unfavorable cases treated with RT [13-15].

Another result of this study that differed from that of other studies was that the CCRT did not show a survival benefit and instead, showed an inferior survival outcome in the CCRT group [2]. Therefore, the RT-alone group and the CCRT group were compared, as shown in **Table 2**, and the comparison revealed trends toward more non-SqCC in the CCRT group and the treatment of more patients after the year 2000. Therefore, selection biases were speculated in the decision-making on the vaginal cancer treatment modality in Korea.

Research on the effect of the tumor HPV status on vaginal cancer survival is still in its early stage mainly due to the rarity of vaginal cancer, but there is a presumption that the tumor HPV status in vaginal cancer would also contribute to the cancer development or the treatment response, based on the studies of head and neck cancer as well as of cervical cancer [16,17]. Sinno et al. [18] examined the HPV status in 60 cases of vaginal cancer, and found that 75% were HPV-positive and 25% were HPV-negative. The unadjusted OS rates in the HPV-positive patients were 57.4% and 35.7% for the HPV-negative patients, but they were statistically insignificant ( $p=0.243$ ) [18]. This study included 27 patients with HPV test results, 63% of whom were HPV-positive, and their survival rates showed no statistically significant difference. That might have been due to the small number of patients in the performed studies, because the data showed a trend toward a better survival outcome in HPV-positive patients. Therefore, a further study with a sufficient number of patients is needed.

Regarding acute toxicity, 12% of the patients experienced a severe (grade 3 or 4) acute complication, lower than in other studies (24% to 40%). However, the late toxicity result was comparable to that of other studies, in that 8% of the patients had a grade 3 or 4 late complication [5,10,19,20]. However, as Hegemann et al. [19] stated, clinicians grade toxicities differently, which makes it difficult to directly compare the results of various studies. However, the toxicities are still considerable. Therefore, based on these data, toxicities should be considered during treatment, especially late toxicities, and special attention should be paid to the characteristics of each tumor. The results of this study showed that a higher FIGO stage and a larger tumor could significantly predict severe late toxicity, as

mentioned in some previous studies [3,4]. Other researchers have reported the predictors of complications as a higher RT dose [21,22], pelvic lymphadenectomy [10], a higher FIGO stage, and a smoking history [3]. As mentioned, the FIGO stage is also a critical risk factor of the disease recurrence and worse CSS, as well as a predictor of severe late toxicity. Therefore, especially in the advanced stage, more precise RT such as intensity-modulated RT or brachytherapy planning with three-dimensional imaging should replace conventional RT to enhance the treatment outcome and to reduce the toxicity.

This multicenter study revealed that the institutions had heterogeneous treatment policies. Furthermore, the obtainable data were limited, such as the location of the tumor, tumor markers, detailed RT technique, and salvage treatment after recurrence, because they were lacking in some institutions. However, a multicenter study on this rare malignancy was inevitable to accrue analyzable data, because a single-institution study could not provide useful prognostic factors of survival due to the small number of patients, except for large institutions, and the study of the national database provided few of the important data. An attempt was made to enhance the homogeneity of the data, especially on the treatment schemes, by strictly excluding cases of prior excision before RT and Tis lesions, which were included in previous studies [2,10].

In conclusion, a survival outcome comparable to or higher than the historical data and moderate toxicities in the patients with primary vaginal cancer were shown, and revealed that a lower FIGO stage and prior hysterectomy are favorable prognostic factors. The FIGO stage and the tumor size were related to higher severe toxicity. Therefore, individualized precise treatment of advanced-stage vaginal cancer is imperative. In the patients whose tumor HPV status was available, 63% had an HPV-positive tumor. No relationship was found between the patients with HPV-positive and -negative tumors, but there was a trend toward better local control in the HPV-positive patients. Therefore, a further study on HPV in vaginal cancer is needed.

## REFERENCES

1. Korea Central Cancer Registry, National Cancer Center. Annual report of cancer statistics in Korea in 2012, Ministry of Health and Welfare. Goyang: National Cancer Center; 2014.
2. Rajagopalan MS, Xu KM, Lin JF, Sukumvanich P, Krivak TC, Beriwal S. Adoption and impact of concurrent chemoradiation therapy for vaginal cancer: a National Cancer Data Base (NCDB) study. *Gynecol Oncol* 2014;135:495-502.  
[PUBMED](#) | [CROSSREF](#)
3. Frank SJ, Jhingran A, Levenback C, Eifel PJ. Definitive radiation therapy for squamous cell carcinoma of the vagina. *Int J Radiat Oncol Biol Phys* 2005;62:138-47.  
[PUBMED](#) | [CROSSREF](#)
4. Tran PT, Su Z, Lee P, Lavori P, Husain A, Teng N, et al. Prognostic factors for outcomes and complications for primary squamous cell carcinoma of the vagina treated with radiation. *Gynecol Oncol* 2007;105:641-9.  
[PUBMED](#) | [CROSSREF](#)
5. Mock U, Kucera H, Fellner C, Knoke TH, Pötter R. High-dose-rate (HDR) brachytherapy with or without external beam radiotherapy in the treatment of primary vaginal carcinoma: long-term results and side effects. *Int J Radiat Oncol Biol Phys* 2003;56:950-7.  
[PUBMED](#) | [CROSSREF](#)
6. de Crevoisier R, Sanfilippo N, Gerbaulet A, Morice P, Pomel C, Castaigne D, et al. Exclusive radiotherapy for primary squamous cell carcinoma of the vagina. *Radiother Oncol* 2007;85:362-70.  
[PUBMED](#) | [CROSSREF](#)

7. Stuart GC, Allen HH, Anderson RJ. Squamous cell carcinoma of the vagina following hysterectomy. *Am J Obstet Gynecol* 1981;139:311-5.  
[PUBMED](#) | [CROSSREF](#)
8. Bell J, Sevin BU, Averette H, Nadji M. Vaginal cancer after hysterectomy for benign disease: value of cytologic screening. *Obstet Gynecol* 1984;64:699-702.  
[PUBMED](#)
9. Daling JR, Madeleine MM, Schwartz SM, Shera KA, Carter JJ, McKnight B, et al. A population-based study of squamous cell vaginal cancer: HPV and cofactors. *Gynecol Oncol* 2002;84:263-70.  
[PUBMED](#) | [CROSSREF](#)
10. Chyle V, Zagars GK, Wheeler JA, Wharton JT, Delclos L. Definitive radiotherapy for carcinoma of the vagina: outcome and prognostic factors. *Int J Radiat Oncol Biol Phys* 1996;35:891-905.  
[PUBMED](#) | [CROSSREF](#)
11. Shah CA, Goff BA, Lowe K, Peters WA, Li CI. Factors affecting risk of mortality in women with vaginal cancer. *Obstet Gynecol* 2009;113:1038-45.  
[PUBMED](#) | [CROSSREF](#)
12. Creasman WT, Phillips JL, Menck HR. The National Cancer Data Base report on cancer of the vagina. *Cancer* 1998;83:1033-40.  
[PUBMED](#) | [CROSSREF](#)
13. Stock RG, Chen AS, Seski J. A 30-year experience in the management of primary carcinoma of the vagina: analysis of prognostic factors and treatment modalities. *Gynecol Oncol* 1995;56:45-52.  
[PUBMED](#) | [CROSSREF](#)
14. Lee LJ, Jhingran A, Kidd E, Cardenes HR, Elshaikh MA, Erickson B, et al. ACR appropriateness Criteria management of vaginal cancer. *Oncology (Williston Park)* 2013;27:1166-73.  
[PUBMED](#)
15. Davis KP, Stanhope CR, Garton GR, Atkinson EJ, O'Brien PC. Invasive vaginal carcinoma: analysis of early-stage disease. *Gynecol Oncol* 1991;42:131-6.  
[PUBMED](#) | [CROSSREF](#)
16. de Jong MC, Pramana J, Kneijens JL, Balm AJ, van den Brekel MW, Hauptmann M, et al. HPV and high-risk gene expression profiles predict response to chemoradiotherapy in head and neck cancer, independent of clinical factors. *Radiother Oncol* 2010;95:365-70.  
[PUBMED](#) | [CROSSREF](#)
17. Walboomers JM, Jacobs MV, Manos MM, Bosch FX, Kummer JA, Shah KV, et al. Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. *J Pathol* 1999;189:12-9.  
[PUBMED](#) | [CROSSREF](#)
18. Sinno AK, Saraiya M, Thompson TD, Hernandez BY, Goodman MT, Steinau M, et al. Human papillomavirus genotype prevalence in invasive vaginal cancer from a registry-based population. *Obstet Gynecol* 2014;123:817-21.  
[PUBMED](#) | [CROSSREF](#)
19. Hegemann S, Schäfer U, Lellé R, Willich N, Micke O. Long-term results of radiotherapy in primary carcinoma of the vagina. *Strahlenther Onkol* 2009;185:184-9.  
[PUBMED](#) | [CROSSREF](#)
20. Kanayama N, Isohashi F, Yoshioka Y, Baek S, Chatani M, Kotsuma T, et al. Definitive radiotherapy for primary vaginal cancer: correlation between treatment patterns and recurrence rate. *J Radiat Res* 2015;56:346-53.  
[PUBMED](#) | [CROSSREF](#)
21. Lee WR, Marcus RB, Sombeck MD, Mendenhall WM, Morgan LS, Freeman DE, et al. Radiotherapy alone for carcinoma of the vagina: the importance of overall treatment time. *Int J Radiat Oncol Biol Phys* 1994;29:983-8.  
[PUBMED](#) | [CROSSREF](#)
22. Eddy GL, Jenrette JM, Creasman WT. Effect of radiotherapeutic technique on local control in primary vaginal carcinoma. *Int J Gynecol Cancer* 1993;3:399-404.  
[PUBMED](#) | [CROSSREF](#)