

Original Article
Urology



CDC6 mRNA Expression Is Associated with the Aggressiveness of Prostate Cancer

Ye-Hwan Kim ,¹ Young Joon Byun ,¹ Won Tae Kim ,¹ Pildu Jeong ,¹ Chunri Yan ,^{1,2} Ho Won Kang ,¹ Yong-June Kim ,¹ Sang-Cheol Lee ,¹ Sung-Kwon Moon ,³ Yung-Hyun Choi ,⁴ Seok Joong Yun ,¹ and Wun-Jae Kim ¹

¹Department of Urology, Chungbuk National University, College of Medicine, Cheongju, Korea

²Department of Preventive Medicine, Medical College, Yanbian University, Yanji, China

³Department of Food and Nutrition, Chung-Ang University, Anseong, Korea

⁴Department of Biochemistry, Donggeui University College of Oriental Medicine, Busan, Korea

OPEN ACCESS

Received: Dec 29, 2017

Accepted: Aug 28, 2018

Address for Correspondence:

Wun-Jae Kim, MD, PhD

Department of Urology, Chungbuk National University, College of Medicine and Institute for Tumor Research, 776 1-Sunhwan-ro, Seowon-gu, Cheongju 28644, Republic of Korea.
E-mail: wjkim@chungbuk.ac.kr

*Ye-Hwan Kim and Young Joon Byun contributed equally to this work.

© 2018 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Ye-Hwan Kim
<https://orcid.org/0000-0002-8676-7119>
Young Joon Byun
<https://orcid.org/0000-0003-0528-1502>
Won Tae Kim
<https://orcid.org/0000-0002-7482-4550>
Pildu Jeong
<https://orcid.org/0000-0002-5602-5376>
Chunri Yan
<https://orcid.org/0000-0002-3043-6687>
Ho Won Kang
<https://orcid.org/0000-0002-8164-4427>

ABSTRACT

Background: Cell division cycle 6 (CDC6) is an essential regulator of DNA replication and plays important roles in the activation and maintenance of the checkpoint mechanisms in the cell cycle. CDC6 has been associated with oncogenic activities in human cancers; however, the clinical significance of CDC6 in prostate cancer (PCa) remains unclear. Therefore, we investigated whether the *CDC6* mRNA expression level is a diagnostic and prognostic marker in PCa.

Methods: The study subjects included 121 PCa patients and 66 age-matched benign prostatic hyperplasia (BPH) patients. *CDC6* expression was evaluated using real-time polymerase chain reaction and immunohistochemical (IH) staining, and then compared according to the clinicopathological characteristics of PCa.

Results: *CDC6* mRNA expression was significantly higher in PCa tissues than in BPH control tissues ($P = 0.005$). In addition, *CDC6* expression was significantly higher in patients with elevated prostate-specific antigen (PSA) levels (> 20 ng/mL), a high Gleason score, and advanced stage than in those with low PSA levels, a low Gleason score, and earlier stage, respectively. Multivariate logistic regression analysis showed that high expression of *CDC6* was significantly associated with advanced stage ($\geq T3b$) (odds ratio [OR], 3.005; confidence interval [CI], 1.212–7.450; $P = 0.018$) and metastasis (OR, 4.192; CI, 1.079–16.286; $P = 0.038$). Intense IH staining for CDC6 was significantly associated with a high Gleason score and advanced tumor stage including lymph node metastasis stage (linear-by-linear association, $P = 0.044$ and $P = 0.003$, respectively).

Conclusion: *CDC6* expression is associated with aggressive clinicopathological characteristics in PCa. CDC6 may be a potential diagnostic and prognostic marker in PCa patients.

Keywords: Prostatic Neoplasms; Gene Expression; Immunohistochemical Staining; *CDC6*

INTRODUCTION

Prostate cancer (PCa) has an extremely heterogeneous clinical course, ranging from indolent and organ-confined to aggressive, metastatic lethal disease.¹ Consequently, the tumor characteristics of PCa need to be accurately assessed so that suitable therapeutic options can

Yong-June Kim 
<https://orcid.org/0000-0001-7638-7174>
 Sang-Cheol Lee 
<https://orcid.org/0000-0002-4163-2210>
 Sung-Kwon Moon 
<https://orcid.org/0000-0002-4514-3457>
 Yung-Hyun Choi 
<https://orcid.org/0000-0002-1454-3124>
 Seok Joong Yun 
<https://orcid.org/0000-0001-7737-4746>
 Wun-Jae Kim 
<https://orcid.org/0000-0002-8060-8926>

Funding

This research was supported by the International Science and Business Belt Program through the Ministry of Science, ICT and Future Planning (2017K000490), Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (No. NRF-2016R1A6A3A11930957) and by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2017R1A6A3A11028303). This work was supported by a research grant from Chungbuk National University, 2012.

Disclosure

The authors have no potential conflict of interest to disclose.

Author Contributions

Conceptualization: Yun SJ, Kim WJ. Data curation: Kim YH, Byun YJ, Kim WT, Jeong P, Yan C, Kang HW, Kim YJ, Lee SC, Moon SK, Choi YH, Kim WJ. Formal analysis: Kim YH, Byun YJ, Kim WT, Jeong P, Yan C, Kang HW, Kim YJ, Lee SC, Moon SK, Choi YH, Kim WJ. Investigation: Kim YH, Kang HW, Yun SJ, Kim WJ. Validation: Byun YJ, Kim WT, Kang HW, Kim WJ. Writing - original draft: Kim YH, Kang HW, Kim WT. Writing - review & editing: Lee SC, Choi YH, Moon SK, Yun SJ, Kim WJ.

be considered. Currently, serum prostate-specific antigen (PSA) levels and histopathological analysis, including the tumor grade and tumor stage as measured by the Gleason score, are the primary determinants of therapeutic decisions. However, none of the bio-markers or histological criteria reported to date have appropriate sensitivity and specificity for detecting, monitoring, or determining the prognosis of PCa.^{2,3} Thus, there are crucial methods capable of predicting clinical outcomes and responses to appropriate treatment in patients with PCa. The abnormal expression of certain genes in cancer cells is closely related to various aspects of tumor progression, including tumor growth, invasion, and metastasis. Proper cell division requires the precise coordination and execution of several events in the cell cycle, including centrosome duplication, DNA replication, mitotic spindle assembly, chromosome segregation, and cytokinesis.⁴ The failed execution or mistiming of any of these events can lead to chromosome segregation defects, resulting in aneuploidy or polyploidy.⁵⁻⁷

Cell division cycle 6 (CDC6), a multi-functional molecular switch, is an important regulator of DNA replication and plays essential roles in the activation and maintenance of the checkpoint mechanisms in the cell cycle.⁸ CDC6 belongs to the AAA+ (ATPases associated with diverse cellular activities) family of ATPases with chaperone-like activities and was initially identified in a genetic screen aimed at detecting mutations that arrest the budding yeast cell cycle.⁹ The human *CDC6* gene is mapped to chromosome 17q21.3 and its expression is regulated by the E2F family of transcription factors that control S phase-promoting genes.¹⁰⁻¹² CDC6 is also a component of a pre-replication complex that forms at the origins of DNA replication in early G1 phase and initiates DNA replication during S phase. CDC6 is involved in checkpoint mechanisms that coordinate S phase of the cell cycle and mitotic entry. By coupling DNA replication and the cell cycle S-M phase checkpoint, CDC6 ensures the entire genome is replicated only once per cell division.⁸ Many previous studies indicate that abnormal expression of CDC6 plays an important role in several human malignancies such as brain tumors,¹³ hepatocellular carcinoma,¹⁴ lung cancer,¹⁵ and ovarian cancer.^{16,17} High expression of CDC6 detected by immunohistochemical (IH) staining and Western blotting is associated with a higher tumor grade and a more advanced stage. Although protein expression of CDC6 is elevated around S phase in the LNCaP PCa cell line,¹⁸ the clinical significance of CDC6 in PCa remains unclear to the best of our knowledge.

The aim of the current study was to assess the clinical significance of CDC6 in PCa using real-time quantitative polymerase chain reaction (RT-qPCR) and IH staining.

METHODS

Study population

This case-control study included 121 cases of newly diagnosed PCa and 66 age-matched benign prostatic hyperplasia (BPH) controls. The study cases were recruited from among patients with histologically confirmed primary adenocarcinoma of the prostate at our institution. Controls were selected from a database of BPH patients who underwent transurethral resection of the prostate (TURP) and were matched according to age and date of blood sampling. Controls with serum PSA levels > 2.5 ng/mL underwent transrectal prostate biopsy before TURP to rule out the presence of cancer, and those with PSA levels > 10 ng/mL were excluded from the study. Subjects with a suspicious history of previous management for PCa or incomplete medical records were also excluded. The Gleason score and 2002 tumor stage, lymph nodes, metastasis (TNM) stage were used as prognostic factors. The

Gleason score was measured from 12-core transrectal biopsy, TURP, or radical prostatectomy specimens. Tumor stage was estimated from radical prostatectomy specimens or from computed tomography, magnetic resonance imaging, or bone scan results.

RNA extraction and construction of cDNA

Total RNA was separated from tissue homogenized in a 5 mL glass tube in 1 mL TRIzol (Invitrogen, Carlsbad, CA, USA). The homogenate was transferred to a 1.5 mL tube and mixed with 200 μ L chloroform. After incubation for 5 minutes at 4°C, the homogenate was centrifuged for 13 minutes at 13,000 g and 4°C. The upper aqueous phase was transferred to a clean tube, 500 μ L isopropanol was added, and the mixture was incubated for 60 minutes at 4°C. The sample was then centrifuged for 8 minutes at 13,000 g and 4°C. Then, the upper aqueous phase was removed, mixed with 500 μ L of 75% ethanol, and centrifuged for 5 minutes at 13,000 g and 4°C. After the upper aqueous layer was discarded, the pellet was dried at room temperature, dissolved in diethylpyrocarbonate-treated water, and stored at -80°C. The quality and integrity of the RNA were confirmed by agarose gel electrophoresis and ethidium bromide staining. cDNA was then prepared from 1 μ g total RNA using the First-Strand cDNA Synthesis kit (Clontech, TAKARA, Otsu, Japan) according to the manufacturer's protocol.

RT-qPCR

To quantify the expression of *CDC6*, real-time PCR was performed using a Rotor Gene 6000 instrument (Corbett Research, Mortlake, Australia). Real-time PCR assays using SYBR Premix Ex Taq (Takara Bio Inc., Otsu, Japan) were carried out in micro-reaction tubes (Corbett Research) using *CDC6* sense (5'-CTC CAG TGA TGC CAA ACT AG-3') and antisense (5'-TGG TGA ACT TTG GCT AGT TC-3') primers. PCR was performed in a final volume of 10 μ L consisting of 5 μ L 2 \times SYBR Premix Ex Taq buffer, 0.5 μ L of each primer (10 pmol/ μ L), and 2 μ L cDNA. The product was purified with a QIAquick Extraction Kit (QIAGEN, Hilden, Germany), quantified with a spectrometer (Perkin Elmer MBA-2000; Perkin Elmer, Fremont, CA, USA), and sequenced with an automated laser fluorescence sequencer (ABI PRISM 3100 Genetic Analyzer; Applied Biosystems, Foster City, CA, USA). The product was serially diluted from 100 pg/ μ L to 0.1 pg/ μ L to establish a standard curve. The real-time PCR conditions were as follows: 1 cycle of 20 seconds at 96°C, followed by 45 cycles of 2 seconds at 96°C for denaturation, 20 seconds at 52°C for annealing, and 20 seconds at 72°C for extension. The melting program was performed at 72°C–99°C, with a heating rate of 1°C per 45 seconds. Spectral data were captured and analyzed using Rotor Gene Real-Time Analysis Software 6.0 Build 14 (Corbett Research). All samples were run in triplicate. Glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) was used as an endogenous reference gene. *CDC6* expression was normalized to the expression of *GAPDH*.

IH staining of CDC6 in PCa and BPH tissues

Paraffin blocks from ten CaP cases and five BPH controls were available for IH analysis. Tissue sections were cut and placed on SuperfrostPlus microscope slides (Fisher Scientific, Cheongju, Korea). Using the Benchmark XT automated immunohistochemistry stainer (Ventana Medical Systems, Tucson, AZ, USA), tissues were stained as follows. Detection was performed using a Ventana Ultraview DAB Kit (Ventana Medical Systems). Sections were deparaffinized using EZ Prep solution. CC1 standard (Tris/Borate/EDTA buffer, pH 8.4) was used for antigen retrieval. DAB inhibitor (3% H₂O₂, endogenous peroxidase) was blocked for 4 minutes at 37°C. Sections were incubated with an anti-CDC6 primary antibody (Abcam Inc., San Diego, CA, USA) for 40 minutes at 37°C and then with the Universal HRP Multimer secondary antibody for 8 minutes at 37°C. Slides were stained with DAB/H₂O₂ substrate for 8 minutes followed by hematoxylin and

bluing reagent counterstain at 37°C. Reaction buffer (Tris buffer, pH 7.6) was used for washing. The staining intensity and proportion of positively stained epithelial cells were evaluated. CDC6 localized primarily in the cytoplasm. The staining intensity was classified as follows: none (0), weak (1), moderate (2), and strong (3). Each specimen was examined and scored separately by three investigators, and discrepancies were discussed until agreement was reached.

Statistical analysis

Clinical variables such as age and PSA levels were compared using the Mann-Whitney U test. Spearman's rank correlation coefficient was calculated to evaluate the association between CDC6 mRNA and protein expression levels. Receiver operating characteristic (ROC) curves were used to identify the optimal cutoff point for each risk score that yielded the highest combined sensitivity and specificity. To evaluate tumor characteristics, Gleason scores were classified as ≤ 7 and ≥ 8 , and the clinical stage was categorized as T2, T3a, T3b, and T4. Multivariate logistic regression analysis was performed to assess the association between CDC6 tissue mRNA expression and clinicopathological parameters in patients with advanced disease or metastasis (odds ratio [OR] and 95% confidence interval [CI]). Statistical analysis was performed using IBM SPSS 23.0 software (IBM Co., Armonk, NY, USA), and a P value < 0.05 was considered statistically significant.

Ethics statement

The Ethics Committee of Chungbuk National University approved the protocol, and written informed consent was obtained from each subject. The collection and analyses of all samples were approved by the Institutional Review Board (IRB) of Chungbuk National University (IRB approval No. 2010-12-010), and informed consent was obtained from each subject.

RESULTS

Baseline characteristics

Table 1 lists the baseline clinical and pathological characteristics of the 66 BPH controls and 121 PCa patients. The median age of the PCa patients was 68.3 years (range, 42–86 years). Among the 121 PCa patients, 103 (85.1%) patients underwent radical prostatectomy and the other 18 (14.9%) patients underwent palliative TURP. At the time of diagnosis, 39 (32.2%) patients had T2 stage, 30 (24.8%) had T3a stage, 30 (24.8%) had T3b stage, and 22 (18.2%) had T4 stage (metastasis). In total, 70 (57.9%), 12 (9.9%), 38 (31.4%), and 1 (0.8%) patients had a Gleason score of 7, 8, 9, and 10, respectively.

Expression levels of CDC6 mRNA in BPH and PCa tissues

The CDC6 mRNA expression level was significantly higher in PCa tissues (median, 253.3×10^5 copies per μg ; interquartile range [IQR], 181.2–317.5) than in BPH tissues (median, 183.5×10^5 copies per μg ; IQR, 131.1–317.6) ($P = 0.005$, **Table 1** and **Fig. 1**).

Relationship between CDC6 tissue mRNA expression levels and clinicopathological features

CDC6 mRNA expression was significantly higher in patients with elevated PSA levels (> 20 ng/mL) than in those with low PSA levels ($P < 0.001$) (**Table 2**). In addition, CDC6 expression was significantly higher in cancer tissue specimens from patients with a high Gleason score and advanced stage than in samples from those with a low Gleason score and earlier stage, respectively ($P < 0.005$ in each case) (**Table 2** and **Fig. 2**).

Table 1. Comparison of clinicopathologic characteristics between PCa cases and controls

Characteristic	Controls	Cases	P value ^a
No.	66	121	
Age, yr (range)	68.8 (46–85)	68.3 (42–86)	0.648
PSA \pm SD, ng/mL	4.0 \pm 4.5	200.6 \pm 809.2	< 0.001
Operation, %			
TURP	66 (100)	18 (14.9)	
Radical prostatectomy		103 (85.1)	
Gleason score, %			
7 (3 + 4)		36 (29.8)	
7 (4 + 3)		34 (28.1)	
8		12 (9.9)	
9		38 (31.4)	
10		1 (0.8)	
Stage, %			
T2		39 (32.2)	
T3a		30 (24.8)	
T3b		30 (24.8)	
T4		22 (18.2)	
CDC6 expression, median (IQR, $\times 10^5$ copies per μ g)	183.5 (131.1–317.6)	253.3 (181.2–371.5)	0.005

PCa = prostate cancer, PSA = prostate-specific antigen, SD = standard deviation, TURP = transurethral resection of the prostate, CDC6 = cell division cycle 6, IQR = interquartile range.

^aThe Mann-Whitney U test was used to compare the expression levels according to clinical variables.




Fig. 1. CDC6 mRNA expression levels in BPH and PCa tissues.

CDC6 = cell division cycle 6, BPH = benign prostatic hyperplasia, PCa = prostate cancer.

Table 2. CDC6 mRNA expression levels according to the PSA level, Gleason score, and PCa stage

Characteristics	No. of patients	CDC6 expression, median (IQR, $\times 10^5$ copies per μ g)	P value ^a
PSA, ng/mL			< 0.001
≤ 20	71	221.6 (155.7–317.7)	
> 20	50	377.7 (226.7–468.2)	
Gleason score			0.002
7 (3 + 4)	36	205.3 (143.8–301.7)	
7 (4 + 3)	34	220.0 (165.7–309.1)	
8	12	290.5 (162.0–454.4)	
≥ 9	39	350.5 (244.2–552.4)	
Stage			< 0.001
T2	39	194.0 (138.2–294.2)	
T3a	30	226.3 (182.6–354.5)	
T3b	30	329.1 (224.6–426.4)	
T4	22	398.1 (281.8–491.4)	

CDC6 = cell division cycle 6, PSA = prostate-specific antigen, PCa = prostate cancer, IQR = interquartile range.

^aThe Mann-Whitney U test was used to compare the expression levels according to clinical variables.




Fig. 2. Relationship between *CDC6* mRNA expression levels and clinicopathological features of Pca. (A) *CDC6* mRNA expression according to Gleason scores. (B) *CDC6* mRNA expression according to the T stage. *CDC6* = cell division cycle 6, Pca = prostate cancer.

Multivariate logistic regression analysis between *CDC6* tissue mRNA expression and advanced disease and metastasis

The area under the ROC curve of *CDC6* expression in advanced stage ($\geq T3b$) was 0.735 (Fig. 3). The highest combined sensitivity of *CDC6* expression in advanced stage (T3b and T4) was 73.1%, and specificity was 66.7%. High *CDC6* tissue expression was significantly associated with advanced stage (OR, 3.005; CI, 1.212–7.450; $P = 0.018$) and metastasis (OR, 4.192; CI, 1.079–16.286; $P = 0.038$) (Table 3).




Fig. 3. ROC curve analysis of *CDC6* expression according to the T stage ($\leq T3a$ and $\geq T3b$). ROC = receiver operating characteristic, *CDC6* = cell division cycle 6, AUC = area under the curve.

Table 3. Multivariate logistic regression analysis of *CDC6* expression as a prognostic marker of advanced disease and metastasis

Parameters	Advanced disease (T stage, $\geq T3b$)		Metastasis	
	OR (95% CI)	P value	OR (95% CI)	P value
Age, yr	1.025 (0.953–1.025)	0.500	0.970 (0.892–1.055)	0.476
PSA (< 50 ng/mL, ≥ 50 ng/mL)	15.246 (1.734–134.046)	0.014	39.949 (7.090–225.083)	< 0.001
Gleason score (≤ 7 , > 8)	5.359 (2.141–13.412)	< 0.001	3.406 (0.968–11.982)	0.056
<i>CDC6</i> expression (low, high)	3.005 (1.212–7.450)	0.018	4.192 (1.079–16.286)	0.038

CDC6 = cell division cycle 6, OR = odds ratio, CI = confidence interval, PSA = prostate-specific antigen.




Fig. 4. The IH staining pattern of CDC6 differs between BPH and PCa tissues (200× magnification). **(A)** Weak cytoplasmic staining intensity of CDC6 in BPH epithelial tissue. Note the intense staining in stromal muscle cells. **(B)** Moderate staining intensity of CDC6 in localized PCa with a Gleason score of 7 (3 + 4). **(C)** Strong staining intensity of CDC6 in advanced PCa with a Gleason score of 8 (4 + 4).
CDC6 = cell division cycle 6, BPH = benign prostatic hyperplasia, PCa = prostate cancer.

IH staining of CDC6 in PCa and BPH tissues

The presence of CDC6 protein was assessed in ten PCa and five BPH samples by IH staining (Fig. 4). CDC6 localized primarily in the cytoplasm in cancer tissues. Cytoplasmic staining of CDC6 correlated well with *CDC6* mRNA expression (Spearman's correlation, $r = 0.576$, $P = 0.025$). The pattern of CDC6 staining was different in BPH tissue; there was intense staining in stromal muscle cells, but the cytoplasm and nuclei of glandular epithelial cells were weakly stained. The staining intensity of CDC6 was significantly higher in PCa tissue than in BPH tissue ($P = 0.015$). Interestingly, intense staining of CDC6 was significantly associated with a high Gleason score and advanced TNM stage (linear-by-linear association, $P = 0.044$ and $P = 0.003$, respectively).

DISCUSSION

In this study, we identified the clinical significance of CDC6 in PCa by performing RT-qPCR and IH staining. The level of *CDC6* mRNA expression was higher in PCa tissues than in BPH tissues, and the increased expression of this gene was associated with a high Gleason score. Additionally, *CDC6* mRNA expression was significantly higher in patients with elevated PSA levels (> 20 ng/mL) and advanced stage than in those with low PSA levels and earlier stage, respectively. IH staining for CDC6 yielded similar results. These results indicate that *CDC6* expression is a novel diagnostic and prognostic marker of PCa.

The human *CDC6* gene is mapped to chromosome 17q21.3¹⁹ and CDC6 is an important component of pre-replication complexes. CDC6 is essential for DNA replication, and CDC6 silencing inhibits DNA replication and arrests cells in G1/S phase, resulting in an inability to enter S phase.^{20,21} CDC6 silencing not only inhibits DNA replication but also leads to cell apoptosis.²² Jin and Fondell²³ reported that CDC6 is a key regulatory target for the androgen receptor and it is a new mechanisms of PCa cell proliferation as an essential regulatory of G1-S phase progression. Aberrant CDC6 expression may cause malignant proliferation of cells and poses a serious risk of carcinogenesis. A few studies have probed the relationship between CDC6 and many cancers, such as cervical cancer,^{24,25} lung cancer,¹⁵ oral squamous cell carcinoma,²⁶ and ovarian cancer,^{16,17} and found that the level of *CDC6* expression is correlated with the extent of malignancy. Sun et al.¹⁷ revealed that *CDC6* is highly expressed in ovarian tissues in contrast with normal ovarian cancer tissues and suggested that *CDC6* is related to the stage of ovarian cancer. Deng et al.¹⁶ demonstrated that elevated levels of *CDC6*

are related to cellular proliferation and poor prognosis in epithelial ovarian cancer. Similarly, Zhang et al.¹⁵ found that the expression level of *CDC6* is significantly higher in lung cancer tissue specimens than in normal lung tissues, and overexpression of *CDC6* is associated with poor overall survival of lung cancer patients.

To the best of our knowledge, no study has addressed the prognostic implications of *CDC6* in PCa. Our study focused on *CDC6* mRNA and protein expression with respect to pathological phenotypes and oncological outcomes. *CDC6* expression was significantly higher in PCa patients than in BPH controls. More importantly, the characteristics of PCa were more aggressive as *CDC6* expression increased. Our result revealed that increased expression of *CDC6* was associated with a high Gleason score, advanced stage, and metastasis. In addition to significant differences of expression of *CDC6* expression in high Gleason score prostate tumors, we also found differential expression of *CDC6* between Gleason score 7 tumors with primary Gleason grades 3 and 4. Because numerous studies suggest that Gleason 3 + 4 tumors have a better prognosis than Gleason 4 + 3 tumors, thus, it is important to distinguish Gleason 3 + 4 vs. 4 + 3 = 7 tumors.^{27,28} On the other hand, we failed to find a significant expression of *CDC6* between those with a total pretreatment serum PSA ≤ 4 vs. 4–10 vs. 10–20 ng/mL (data not shown). The only significant difference in *CDC6* expression was observed between those with a total pretreatment serum PSA ≤ 20 vs. < 20 ng/mL. Clinically, the PSA level alone does not facilitate precise staging on an individual basis, although advanced stage tends to correlate with an increased PSA level.²⁹ Therefore, the data presented here suggest that *CDC6* expression could be a novel indicator of tumor aggressiveness in patients with PCa.

This study has a possible weakness. IH analysis was performed on a small number of samples. Nevertheless, this analysis is sufficient to demonstrate that cytoplasmic *CDC6* staining correlates with *CDC6* mRNA expression and that the pattern of *CDC6* staining differs between BPH and PCa tissues and varies according to the aggressiveness of PCa.

In conclusion, the *CDC6* mRNA level was significantly higher in PCa patients than in BPH controls. Increased PSA levels, higher Gleason scores, and advanced stage disease were closely associated with the level of *CDC6* expression. Our findings indicate that expression of *CDC6* is a novel diagnostic and prognostic marker of PCa.

ACKNOWLEDGMENTS

The specimens used in this study were provided by Chungbuk National University Hospital, a member of the National Biobank of Korea, which is supported by the Ministry of Health, Welfare and Family Affairs. All samples from the National Biobank of Korea were obtained with informed consent under institutional review board-approved protocols. The authors wish to thank Ms. Eun-Ju Shim from the National Biobank of Korea at Chungbuk National University Hospital for sample preparation and technical assistance.

REFERENCES

1. Seo WI, Kang PM, Chung JI. Predictive value of the cancer of the prostate risk assessment score for recurrence-free survival after radical prostatectomy in Korea: a single-surgeon series. *Korean J Urol* 2014;55(5):321-6.
PUBMED | CROSSREF

2. Kattan MW, Eastham JA, Stapleton AM, Wheeler TM, Scardino PT. A preoperative nomogram for disease recurrence following radical prostatectomy for prostate cancer. *J Natl Cancer Inst* 1998;90(10):766-71.
[PUBMED](#) | [CROSSREF](#)
3. Partin AW, Kattan MW, Subong EN, Walsh PC, Wojno KJ, Oesterling JE, et al. Combination of prostate-specific antigen, clinical stage, and Gleason score to predict pathological stage of localized prostate cancer. A multi-institutional update. *JAMA* 1997;277(18):1445-51.
[PUBMED](#) | [CROSSREF](#)
4. Negrini S, Gorgoulis VG, Halazonetis TD. Genomic instability--an evolving hallmark of cancer. *Nat Rev Mol Cell Biol* 2010;11(3):220-8.
[PUBMED](#) | [CROSSREF](#)
5. Charames GS, Bapat B. Genomic instability and cancer. *Curr Mol Med* 2003;3(7):589-96.
[PUBMED](#) | [CROSSREF](#)
6. Romanowski P, Madine MA. Mechanisms restricting DNA replication to once per cell cycle: MCMs, pre-replicative complexes and kinases. *Trends Cell Biol* 1996;6(5):184-8.
[PUBMED](#) | [CROSSREF](#)
7. Sasaki H, Kawano O, Endo K, Suzuki E, Yukiue H, Kobayashi Y, et al. Human MOB1 expression in non-small-cell lung cancer. *Clin Lung Cancer* 2007;8(4):273-6.
[PUBMED](#) | [CROSSREF](#)
8. Borlado LR, Méndez J. CDC6: from DNA replication to cell cycle checkpoints and oncogenesis. *Carcinogenesis* 2008;29(2):237-43.
[PUBMED](#) | [CROSSREF](#)
9. Hartwell LH. Sequential function of gene products relative to DNA synthesis in the yeast cell cycle. *J Mol Biol* 1976;104(4):803-17.
[PUBMED](#) | [CROSSREF](#)
10. Hateboer G, Wobst A, Petersen BO, Le Cam L, Vigo E, Sardet C, et al. Cell cycle-regulated expression of mammalian CDC6 is dependent on E2F. *Mol Cell Biol* 1998;18(11):6679-97.
[PUBMED](#) | [CROSSREF](#)
11. Ohtani K, Tsujimoto A, Ikeda M, Nakamura M. Regulation of cell growth-dependent expression of mammalian CDC6 gene by the cell cycle transcription factor E2F. *Oncogene* 1998;17(14):1777-85.
[PUBMED](#) | [CROSSREF](#)
12. Yan Z, DeGregori J, Shohet R, Leone G, Stillman B, Nevins JR, et al. CDC6 is regulated by E2F and is essential for DNA replication in mammalian cells. *Proc Natl Acad Sci U S A* 1998;95(7):3603-8.
[PUBMED](#) | [CROSSREF](#)
13. Ohta S, Koide M, Tokuyama T, Yokota N, Nishizawa S, Namba H. CDC6 expression as a marker of proliferative activity in brain tumors. *Oncol Rep* 2001;8(5):1063-6.
[PUBMED](#)
14. Xiong XD, Fang JH, Qiu FE, Zhao J, Cheng J, Yuan Y, et al. A novel functional polymorphism in the CDC6 promoter is associated with the risk for hepatocellular carcinoma. *Mutat Res* 2008;643(1-2):70-4.
[PUBMED](#) | [CROSSREF](#)
15. Zhang X, Xiao D, Wang Z, Zou Y, Huang L, Lin W, et al. MicroRNA-26a/b regulate DNA replication licensing, tumorigenesis, and prognosis by targeting CDC6 in lung cancer. *Mol Cancer Res* 2014;12(11):1535-46.
[PUBMED](#) | [CROSSREF](#)
16. Deng Y, Jiang L, Wang Y, Xi Q, Zhong J, Liu J, et al. High expression of CDC6 is associated with accelerated cell proliferation and poor prognosis of epithelial ovarian cancer. *Pathol Res Pract* 2016;212(4):239-46.
[PUBMED](#) | [CROSSREF](#)
17. Sun TY, Xie HJ, He H, Li Z, Kong LF. miR-26a inhibits the proliferation of ovarian cancer cells via regulating CDC6 expression. *Am J Transl Res* 2016;8(2):1037-46.
[PUBMED](#)
18. Liu Y, Gong Z, Sun L, Li X. FOXM1 and androgen receptor co-regulate CDC6 gene transcription and DNA replication in prostate cancer cells. *Biochim Biophys Acta* 2014;1839(4):297-305.
[PUBMED](#) | [CROSSREF](#)
19. Williams RS, Shohet RV, Stillman B. A human protein related to yeast Cdc6p. *Proc Natl Acad Sci U S A* 1997;94(1):142-7.
[PUBMED](#) | [CROSSREF](#)
20. Niimi S, Arakawa-Takeuchi S, Uranbileg B, Park JH, Jinno S, Okayama H. CDC6 protein obstructs apoptosome assembly and consequent cell death by forming stable complexes with activated Apaf-1 molecules. *J Biol Chem* 2012;287(22):18573-83.
[PUBMED](#) | [CROSSREF](#)

21. Piatti S, Lengauer C, Nasmyth K. CDC6 is an unstable protein whose de novo synthesis in G1 is important for the onset of S phase and for preventing a 'reductional' anaphase in the budding yeast *Saccharomyces cerevisiae*. *EMBO J* 1995;14(15):3788-99.
[PUBMED](#) | [CROSSREF](#)
22. Feng CJ, Lu XW, Luo DY, Li HJ, Guo JB. Knockdown of CDC6 inhibits proliferation of tongue squamous cell carcinoma Tca8113 cells. *Technol Cancer Res Treat* 2013;12(2):173-81.
[PUBMED](#) | [CROSSREF](#)
23. Jin F, Fondell JD. A novel androgen receptor-binding element modulates CDC6 transcription in prostate cancer cells during cell-cycle progression. *Nucleic Acids Res* 2009;37(14):4826-38.
[PUBMED](#) | [CROSSREF](#)
24. Xiong XD, Zeng LQ, Xiong QY, Lu SX, Zhang ZZ, Luo XP, et al. Association between the CDC6 G1321A polymorphism and the risk of cervical cancer. *Int J Gynecol Cancer* 2010;20(5):856-61.
[PUBMED](#) | [CROSSREF](#)
25. Martin CM, Astbury K, McEvoy L, O'Toole S, Sheils O, O'Leary JJ. Gene expression profiling in cervical cancer: identification of novel markers for disease diagnosis and therapy. *Methods Mol Biol* 2009;511:333-59.
[PUBMED](#) | [CROSSREF](#)
26. Feng CJ, Li HJ, Li JN, Lu YJ, Liao GQ. Expression of Mcm7 and CDC6 in oral squamous cell carcinoma and precancerous lesions. *Anticancer Res* 2008;28(6A):3763-9.
[PUBMED](#)
27. Makarov DV, Sanderson H, Partin AW, Epstein JI. Gleason score 7 prostate cancer on needle biopsy: is the prognostic difference in Gleason scores 4 + 3 and 3 + 4 independent of the number of involved cores? *J Urol* 2002;167(6):2440-2.
[PUBMED](#) | [CROSSREF](#)
28. Kang HW, Jung HD, Lee JY, Kwon JK, Jeh SU, Cho KS, et al. The Within-Group Discrimination Ability of the Cancer of the Prostate Risk Assessment Score for Men with Intermediate-Risk Prostate Cancer. *J Korean Med Sci* 2018;33(5):e36.
[PUBMED](#) | [CROSSREF](#)
29. Ruckle HC, Klee GG, Oesterling JE. Prostate-specific antigen: concepts for staging prostate cancer and monitoring response to therapy. *Mayo Clin Proc* 1994;69(1):69-79.
[PUBMED](#) | [CROSSREF](#)