

- 1 -

· · · ·

---

[ ]

1/3 , 가 가  
, 가  
·

: , ,

---

가 가  
가

, 0.3%  
4,8,9)

, 48 5  
가 1-3) 8 1/3

---

: 302-241, 가 685

: +82.42-600-6905, Fax: +82.42-545-2373  
e-mail: wslecos@kyuh.co.kr

\* 2001

가

(Fig. 1).

가

, , , , , .  
,  
,  
,  
,

가  
(Fig. 2).

가 가

가 가

가 .

, 가  
2  
, 1 5  
,

## REFERENCES

- 1) **Barbier O, Malghem J, Delaere O, Vande Berg B**

- 
- and Rombouts JJ:** Injury to the brachial plexus by a fragment of bone after fracture of the clavicle. *J Bone Joint Surg*, 79(4)-B: 534-536, 1997.
- 2) **Chen CE and Liu HC:** Delayed brachial plexus neuropraxia complicating malunion of the clavicle. *Am J Orthop*, 29(4): 321-322, 2000.
- 3) **Connolly JF and Dhne R:** Nonunion of the clavicle and thoracic outlet syndrome. *J Trauma*, 29(8): 1127-1132, 1989.
- 4) **Hill JM, McGuire MH and Crosby LA:** Closed treatment of displaced middle-third fractures of clavicle gives poor results. *J Bone Joint Surg*, 79(4)-B: 537-539, 1997.
- 5) **Holloway RG, Courtright CE, Rahilly CR, Tottenham SM and Shrier D:** Magnetic resonance imaging in delayed brachial plexopathy following a clavicle fracture. *Eur Neurol*, 40(2): 105-106, 1998.
- 6) **Kay SP and Eckardt JJ:** Brachial plexus palsy secondary to clavicular nonunion. Case report and literature survey. *Clin Orthop*, 206: 219-222, 1986.
- 7) **Lee SE, Shin YS, Yang JH and Lee JH:** Brachial plexus injury to exuberant callus formation of fracture of clavicle. *J Korean Society Fractures*, 11(2): 427-434, 1998.
- 8) **Marz SO, Welliver PS and Welliver DI:** Brachial plexus neuropraxia complicating a comminuted clavicle fracture in a college football player. Case report and review of the literature. *Am J Sports Med*, 17(4): 581-583, 1989.
- 9) **Rumball KM, Da Silva VF, Preston DN and Carruthers CC:** Brachial-plexus injury after clavicular fracture: case report and literature review. *Can J Surg*, 34(3): 264-266, 1991.

**Abstract****Delayed Brachial Plexus Palsy due to Clavicular Fracture  
- A Case Report -**

**Woo-Suk Lee, M.D., Whan-Yong Chung, M.D., Taek-Soo Jeon, M.D.,  
Yong-Sang Kim, M.D., Nam-Hyun Kim, M.D.**

*Department of Orthopedic Surgery, Konyang University College of Medicine, Daejeon, Korea*

The brachial plexus palsies secondary to nonunion of the clavicle fracture are extremely rare. The nonunions are hypertrophic and usually in the middle third of the clavicle. Hypertrophic callus produced during healing process will cause a compression of the neurovascular bundle. This lesion requires operative treatment for decompression of the brachial plexus and internal fixation of nonunion. We present a case of delayed brachial plexus palsy due to nonunion and excessive callus formation of a clavicular fracture.

**Key Words:** Clavicle, Fracture, Brachial plexus palsy

**Address reprint requests to** \_\_\_\_\_

Woo-Suk Lee

685 Gasuwon-dong, Seo-gu, Daejeon 302-241, Korea

Department of Orthopedic Surgery, Konyang University Hospital

Tel : +82.42-600-6905, Fax : +82.42-545-2373

E-mail : wslees@kyuh.co.kr