

# Early Detection of Pulmonary Hypertension in Connective Tissue Disease

Chan Hong Jeon

Division of Rheumatology, Department of Internal Medicine, Soonchunhyang University Bucheon Hospital, Bucheon, Korea

Pulmonary hypertension (PH) is a hemodynamic and pathophysiological state that is defined by an increase in mean pulmonary arterial pressure  $\geq 25$  mmHg at rest measured with right heart catheterization (RHC) [1].

PH can be classified into 5 main subgroups by the clinical and pathophysiological characteristics: group 1 - pulmonary arterial hypertension (PAH); group 2 - PH due to left heart disease; group 3 - PH due to chronic lung disease and/or hypoxemia; group 4 - PH due to chronic thromboembolism or other pulmonary artery obstruction; and group 5 - PH due to uncertain or multifactorial mechanisms [2].

Group 1 PH, PAH, is characterized by a pre-capillary PH in the absence of other causes, such as lung disease, left heart disease, chronic thromboembolism, etc. Connective tissue disease (CTD)-associated PAH belongs to group 1 PH [3]. The European Society of Cardiology (ESC) and European Respiratory Society (ERS) define PAH as a mean pulmonary artery pressure of  $\geq 25$  mmHg at rest and pulmonary artery wedge pressure/left ventricular end diastolic pressure of  $\leq 15$  mmHg and pulmonary vascular resistance of more than 3 wood units by RHC [4].

The reported prevalence of PAH is 5~25 cases/million with an incidence of 2~5 cases/million [5]. The most common etiology of PAH in adults is idiopathic, followed by CTD, congenital heart disease or portal hypertension [6].

In a UK study in 2005, the prevalence of CTD-PAH was 4.23/million and the incidence was 1.55/million [7]. The frequency of PAH among patients with CTD is increasing, which may be due to the increased awareness of the subtypes of the disease and the use of non-invasive screening

diagnostic tests [6,8]. Among CTDs, systemic sclerosis (SSc) is the most prevalent cause of CTD-PAH, comprising 75% and followed by systemic lupus erythematosus (8%~19%), mixed connective tissue disease (8%~9%), rheumatoid arthritis (3%~5%), dermatomyositis/polymyositis (4%), unclassified CTD (2%), and Sjögren's syndrome (1%). The prevalence of PAH in SSc patients has been estimated to be 8%~12% [9].

Regarding the survival of subgroups of PAH, patients with CTD have more severe symptoms and a poorer prognosis than other groups in the absence of therapy [10]. The survival of PAH is improved significantly after early detection of the disease, the application of vasodilator therapy, and improved therapeutic strategies [5]. The data from PAH trials shows that earlier intervention may improve the treatment efficacy and delay the time to clinical deterioration [3].

Unfortunately, most patients with PH are diagnosed late. The initial phase of PAH is clinically silent and detectable signs and symptoms usually appear when the pathological changes are fully developed. The length of this preclinical phase of the disease is currently unknown [3]. Furthermore, nonspecific symptoms of PH-CTD usually overlap with other common medical conditions, such as musculoskeletal disease or anemia, and contribute to delayed diagnosis [3,11]. Considering the limitations of reliance on exertional dyspnea, other tools for early detection are required.

In the past issue of *Journal of Rheumatic Diseases*, Jung et al. [12] examined the prevalence of PH in patients with CTD and evaluated the various clinical manifestations

Received : November 1, 2018, Revised : November 30, 2018, Accepted : November 30, 2018

Corresponding to : Chan Hong Jeon  <http://orcid.org/0000-0002-2430-7264>

Division of Rheumatology, Department of Internal Medicine, Soonchunhyang University Bucheon Hospital, 170 Jomaru-ro, Wonmi-gu, Bucheon 14584, Korea. E-mail : [chjeon@gmail.com](mailto:chjeon@gmail.com)

Copyright © 2019 by The Korean College of Rheumatology. All rights reserved.

This is an Open Access article, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

and biomarkers as potential tools to detect PH early. The results of the pulmonary function test, total cholesterol, red cell volume distribution width, alkaline phosphatase, and New York Heart Association functional class III or IV differed significantly according to the presence of PH. On the other hand, there was no difference in the incidence of PH according to the presence of interstitial lung disease (ILD). However, the study should be interpreted with caution because PH was diagnosed with echocardiography without confirmation of RHC. In addition, while the incidence of ILD was compared between groups with PH or without PH, the extent of ILD was not considered in the analysis.

Several clinical risk factors for the development of PAH in SSc have been suggested: advanced age, post-menopausal state, degree of skin involvement, disease duration of more than 5 years, severe peripheral vascular disease, presence of pulmonary fibrosis, microstomia, gastroesophageal reflux and dysphagia, telangiectasia, and abnormal nail fold capillaroscopy [9,13-15].

SSc-ILD is a frequent manifestation of SSc, occurring in up to 75% of patients [16]. Coexistent ILD and PAH may be identified in 2%~18% of SSc patients [17]. SSc-PH can occur in isolation or in combination with ILD as a direct consequence of severe ILD (group 3 PH) or just co-occurrence with ILD (PAH and mild ILD insufficient to cause PH). The discrimination of the two conditions has been defined variably according to the extent of chest involvement on high-resolution computed tomography together with lung function testing abnormalities [18]. On the other hand, the causality and contribution of ILD on the significance of PH are uncertain in some cases [14,15,19]. Mild ILD has been suggested as a risk factor for the subsequent development of PAH [16,20]. Furthermore, the combination of ILD and PH shows a dismal prognosis with 3-year survival rates of 35% [21].

Although echocardiography is the most widely used noninvasive modality to screen for PAH, other noninvasive markers have been studied for the detection of SSc-PAH. A low diffusing capacity for carbon monoxide (DLCO) can also predict the occurrence of PAH in SSc patients. Patients with SSc-PAH typically have moderate-to-severe reductions in DLCO at the time of diagnosis, and a predicted DLCO of <60% is associated with a markedly increased probability of PAH [13,22-25].

B-type natriuretic peptide (BNP) is released from ventricular myocytes when the wall tension increases. In patients with PAH, BNP and N-terminal pro-brain natriu-

retic peptide (NT-pro-BNP) correlate with the disease severity and is also highly predictive of SSc-PAH [13,26-28].

Several biomarkers are also considered as a tool for the early diagnosis and screening of PAH, but a few of them have been tested in prospective studies [29,30]. Subtypes of antinuclear antibody, such as anti-nucleosome antibodies, anti-centromere antibodies, anti-U3-ribonucleoprotein (anti-U3-RNP), and Anti-Th/To are associated with PAH while anti-U1-RNP is associated with a reduced risk [29,30].

Anti-angiotensin receptor type-1 antibody and anti-endothelin receptor type A, which affect the inflammation and fibrotic process by direct receptor activation, are also considered diagnostic markers for CTD-PAH, particularly in SSc [29,31,32].

Chemokines, growth differentiation factor-15, and markers of vascular injury, such as von Willebrand factor, endothelin-1, and vascular endothelial growth factor, are also suggested for use in the early detection of PAH in CTDs, especially SSc [29,31].

Longitudinal observational studies have shown that systematic screening for SSc-PAH can improve the outcomes [33]. Moreover, the accuracy of screening algorithms for SSc-PAH can be improved by combining different tests that are predictive of the presence of PAH [33]. The DETECT (early, simple and reliable detection of pulmonary arterial hypertension in SSc) protocol is a two-step scoring algorithm based on forced vital capacity, DLCO, telangiectasia, anti-centromere antibody, NT-pro BNP, serum urate, electrocardiogram, and subsequent echocardiography to determine the need for RHC in systemic sclerosis. The protocol results in fewer false negatives than an approach using echocardiography alone [34]. The new 2015 ESC/ERS guidelines also recommend a combined screening approach (using biomarkers, lung function, and echocardiography) to predict the presence of PAH in SSc [4].

In conclusion, an evaluation of the clinical risk factors, biomedical markers, and their combinations are valuable tools for the early detection of CTD-PH, thereby enabling prompt intervention to improve the survival of patients.

## CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

## REFERENCES

1. Hoepfer MM, Bogaard HJ, Condliffe R, Frantz R, Khanna D, Kurzyna M, et al. Definitions and diagnosis of pulmonary hypertension. *J Am Coll Cardiol* 2013;62(25 Suppl): D42-50.
2. Bhatnagar A, Wiesen J, Dweik R, Chaisson NF. Evaluating suspected pulmonary hypertension: a structured approach. *Cleve Clin J Med* 2018;85:468-80.
3. Lau EM, Manes A, Celermajer DS, Galiè N. Early detection of pulmonary vascular disease in pulmonary arterial hypertension: time to move forward. *Eur Heart J* 2011;32:2489-98.
4. Galiè N, Humbert M, Vachiery JL, Gibbs S, Lang I, Torbicki A, et al. 2015 ESC/ERS guidelines for the diagnosis and treatment of pulmonary hypertension: the joint task force for the diagnosis and treatment of pulmonary hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS): endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). *Eur Heart J* 2016;37:67-119.
5. Maron BA, Galiè N. Diagnosis, treatment, and clinical management of pulmonary arterial hypertension in the contemporary era: a review. *JAMA Cardiol* 2016;1:1056-65.
6. Hoepfer MM, Simon R, Gibbs J. The changing landscape of pulmonary arterial hypertension and implications for patient care. *Eur Respir Rev* 2014;23:450-7.
7. Condliffe R, Kiely DG, Peacock AJ, Corris PA, Gibbs JS, Vrapai F, et al. Connective tissue disease-associated pulmonary arterial hypertension in the modern treatment era. *Am J Respir Crit Care Med* 2009;179:151-7.
8. Thenappan T, Shah SJ, Rich S, Gomberg-Maitland M. A USA-based registry for pulmonary arterial hypertension: 1982-2006. *Eur Respir J* 2007;30:1103-10.
9. Aithala R, Alex AG, Danda D. Pulmonary hypertension in connective tissue diseases: an update. *Int J Rheum Dis* 2017;20:5-24.
10. Benza RL, Miller DP, Gomberg-Maitland M, Frantz RP, Foreman AJ, Coffey CS, et al. Predicting survival in pulmonary arterial hypertension: insights from the Registry to Evaluate Early and Long-Term Pulmonary Arterial Hypertension Disease Management (REVEAL). *Circulation* 2010;122:164-72.
11. Haddad RN, Mielniczuk LM. An evidence-based approach to screening and diagnosis of pulmonary hypertension. *Can J Cardiol* 2015;31:382-90.
12. Jung JY, Lee CH, Kim HA, Choi ST, Lee JH, Yoon BY, et al. Pulmonary hypertension in connective tissue disease is associated with the New York Heart Association functional class and forced vital capacity, but not with interstitial lung disease. *J Rheum Dis* 2018;25:179-87.
13. Steen V, Medsger TA Jr. Predictors of isolated pulmonary hypertension in patients with systemic sclerosis and limited cutaneous involvement. *Arthritis Rheum* 2003;48:516-22.
14. Avouac J, Airò P, Meune C, Beretta L, Dieude P, Caramaschi P, et al. Prevalence of pulmonary hypertension in systemic sclerosis in European Caucasians and metaanalysis of 5 studies. *J Rheumatol* 2010;37:2290-8.
15. Condliffe R, Howard LS. Connective tissue disease-associated pulmonary arterial hypertension. *F1000Prime Rep* 2015;7:06.
16. Bussone G, Mouthon L. Interstitial lung disease in systemic sclerosis. *Autoimmun Rev* 2011;10:248-55.
17. Michelfelder M, Becker M, Riedlinger A, Siegert E, Drömann D, Yu X, et al. Interstitial lung disease increases mortality in systemic sclerosis patients with pulmonary arterial hypertension without affecting hemodynamics and exercise capacity. *Clin Rheumatol* 2017;36:381-90.
18. Goh NS, Desai SR, Veeraraghavan S, Hansell DM, Copley SJ, Maher TM, et al. Interstitial lung disease in systemic sclerosis: a simple staging system. *Am J Respir Crit Care Med* 2008;177:1248-54.
19. Hoepfer MM, Andreas S, Bastian A, Claussen M, Ghofrani HA, Gorenflo M, et al. Pulmonary hypertension due to chronic lung disease: updated recommendations of the Cologne consensus conference 2011. *Int J Cardiol* 2011;154 Suppl 1:S45-53.
20. Morrisroe K, Huq M, Stevens W, Rabusa C, Proudman SM, Nikpour M; Australian Scleroderma Interest Group (ASIG). Risk factors for development of pulmonary arterial hypertension in Australian systemic sclerosis patients: results from a large multicenter cohort study. *BMC Pulm Med* 2016;16:134.
21. Lefèvre G, Dauchet L, Hachulla E, Montani D, Sobanski V, Lambert M, et al. Survival and prognostic factors in systemic sclerosis-associated pulmonary hypertension: a systematic review and meta-analysis. *Arthritis Rheum* 2013;65:2412-23.
22. Hachulla E, Gressin V, Guillemin L, Carpentier P, Diot E, Sibilia J, et al. Early detection of pulmonary arterial hypertension in systemic sclerosis: a French nationwide prospective multicenter study. *Arthritis Rheum* 2005;52:3792-800.
23. Iudici M, Codullo V, Giuggioli D, Riccieri V, Cuomo G, Breda S, et al. Pulmonary hypertension in systemic sclerosis: prevalence, incidence and predictive factors in a large multicentric Italian cohort. *Clin Exp Rheumatol* 2013;31(2 Suppl 76):31-6.
24. Hsu VM, Chung L, Hummers LK, Wigley F, Simms R, Bolster M, et al. Development of pulmonary hypertension in a high-risk population with systemic sclerosis in the Pulmonary Hypertension Assessment and Recognition of Outcomes in Scleroderma (PHAROS) cohort study. *Semin Arthritis Rheum* 2014;44:55-62.
25. Hinchcliff M, Fischer A, Schioppa E, Steen VD; PHAROS Investigators. Pulmonary Hypertension Assessment and Recognition of Outcomes in Scleroderma (PHAROS): baseline characteristics and description of study population. *J Rheumatol* 2011;38:2172-9.
26. Leuchte HH, Holzapfel M, Baumgartner RA, Ding I, Neurohr C, Vogeser M, et al. Clinical significance of brain natriuretic peptide in primary pulmonary hypertension. *J Am Coll Cardiol* 2004;43:764-70.
27. Thakkar V, Stevens WM, Prior D, Moore OA, Byron J, Liew D, et al. N-terminal pro-brain natriuretic peptide in a novel screening algorithm for pulmonary arterial hypertension in systemic sclerosis: a case-control study. *Arthritis Res Ther* 2012;14:R143.
28. Allanore Y, Borderie D, Avouac J, Zerkak D, Meune C, Hachulla E, et al. High N-terminal pro-brain natriuretic peptide levels and low diffusing capacity for carbon monoxide

- as independent predictors of the occurrence of precapillary pulmonary arterial hypertension in patients with systemic sclerosis. *Arthritis Rheum* 2008;58:284-91.
29. Odler B, Foris V, Gungl A, Müller V, Hassoun PM, Kwapiszewska G, et al. Biomarkers for pulmonary vascular remodeling in systemic sclerosis: a pathophysiological approach. *Front Physiol* 2018;9:587.
  30. Warwick G, Thomas PS, Yates DH. Biomarkers in pulmonary hypertension. *Eur Respir J* 2008;32:503-12.
  31. Affandi AJ, Radstake TR, Marut W. Update on biomarkers in systemic sclerosis: tools for diagnosis and treatment. *Semin Immunopathol* 2015;37:475-87.
  32. Becker MO, Kill A, Kutsche M, Guenther J, Rose A, Tabeling C, et al. Vascular receptor autoantibodies in pulmonary arterial hypertension associated with systemic sclerosis. *Am J Respir Crit Care Med* 2014;190:808-17.
  33. Thakkar V, Lau EM. Connective tissue disease-related pulmonary arterial hypertension. *Best Pract Res Clin Rheumatol* 2016;30:22-38.
  34. Coghlan JG, Denton CP, Grünig E, Bonderman D, Distler O, Khanna D, et al. Evidence-based detection of pulmonary arterial hypertension in systemic sclerosis: the DETECT study. *Ann Rheum Dis* 2014;73:1340-9.