

Activation domain in P67^{phox} regulates the steady state reduction of FAD in gp91^{phox}

Chang-Hoon Han and Mun-Han Lee^{1*}

Department of Biochemistry, Swiss Federal Institute of Technology in Zurich, Universitatstrasse 16, 8092 Zurich, Switzerland

¹Department of Biochemistry, College of Veterinary Medicine, Seoul National University, Suwon 441-744, Korea

An activation domain in p67^{phox} (residues 199-210) is critical for regulating NADPH oxidase activity in cell-free system [10]. To determine the steady state reduction of FAD, thioacetamide-FAD was reconstituted in gp91^{phox}, and the fluorescence of its oxidised form was monitored. Omission of p67^{phox} decreased the steady state reduction of the FAD from 28% to 4%, but omission of p47^{phox} had little effect. A series of the truncated forms of p67^{phox} were expressed in *E.coli* to determine the domain in p67^{phox} which is essential for regulating the steady state of FAD reduction. The minimal length of p67^{phox} for regulating the steady state of FAD reduction is shown to be 1-210 using a series of truncation mutants which indicates that the region 199-210 is also important for regulating electron flow within flavocytochrome b₅₅₈. The deletion of this domain not only decreased the superoxide generation but also decreased the steady state of FAD reduction. Therefore, the activation domain on p67^{phox} regulates the reductive half-reaction for FAD, consistent with a dominant effect on hydride/electron transfer from NADPH to FAD.

Key words: the activation domain on p67^{phox}; the steady state of FAD reduction.

Introduction


Neutrophils and macrophages produce superoxide and other reactive oxygen species that participate in intracellular killing of phagocytized microorganisms [2,5]. Superoxide generation is catalyzed by NADPH oxidase which consists of both cytosolic factors (p47^{phox} and p67^{phox}) and plasma membrane associated flavocytochrome b₅₅₈. In cell resting state, cytosolic factors p47^{phox}, p67^{phox} exist in the cytosol as a complex along with a third component,

p40^{phox}, which appears to stabilize a 240 kDa complex of cytosolic factors [9, 21]. Upon activation, in response to microorganisms or to a variety of soluble agonists, cytosolic factors p47^{phox}, p67^{phox}, and possibly p40^{phox} translocate to membrane where they bind directly or indirectly with flavocytochrome b₅₅₈ [7, 18]. The small GTP-binding protein, Rac, translocates to membrane independently of the other cytosolic components [8, 11], and thereby assembled complex catalyzes the reduction of oxygen to superoxide.

Flavocytochrome b₅₅₈ is a membrane-associated heterodimer (p22^{phox} and gp91^{phox}) that contains putative binding sites for NADPH, FAD, and heme [16, 19] and considered to be redox center of the NADPH oxidase. Three cytosolic components (p47^{phox}, p67^{phox}, and small GTPase Rac) are considered to be regulatory subunits of NADPH oxidase. A great deal of current research involves understanding the protein-protein interactions among the components of NADPH oxidase complex, and how these change with the activation state. Supporting the importance of these interactions, individuals with genetic deficiencies or mutations in p47^{phox}, p67^{phox}, or one of the subunits of cytochrome b₅₅₈ (gp91^{phox} and p22^{phox}) exhibit chronic granulomatous disease [5], which is characterized by the inability of phagocytic leukocytes (neutrophils, eosinophils, monocytes, and macrophages) to generate active oxygen species which are necessary for killing of phagocytized pathogens (reviewed in 11).

NADPH oxidase activity can be reconstituted *in vitro* using purified cytosolic factors p47^{phox}, p67^{phox}, GTPγS preloaded Rac, and phospholipid-reconstituted flavocytochrome b₅₅₈ along with an anionic amphiphiles such as arachidonate [1, 17].

Based on chemical precedent and structural models of the enzyme [22], the pathway for electron flow within flavocytochrome b₅₅₈ has been proposed in Scheme I.



Our recent study identified an activation domain in p67^{phox}

*Corresponding author

Phone: 82-31-290-2741; Fax: 82-31-293-0084;

E-mail: vetlee@snu.ac.kr

that is essential for NADPH oxidase activity [10]. Deletion of this region within residues 199~210 completely eliminated NADPH oxidase activity.

In the present study, we observed that the activation domain is also essential for regulating electron flux in the complex. We propose that the activation domain on p67^{phox} directly activates a particular step in the electron transfer pathway depicted above in Scheme I. We provide evidence that the activation domain on p67^{phox} regulates the reduction of FAD by NADPH, consistent with the regulation of the NADPH → FAD hydride/electron transfer reaction.

Materials and Methods

Preparation of plasma membrane, cytochrome b₅₅₈ and recombinant proteins:

Plasma membranes were isolated as described by Burnham et al. [4]. Further purification steps were done for isolating cytochrome b₅₅₈ from plasma membrane as described previously [14]. Rac cDNA cloned in pGEX-2T was expressed in DH5a cells as a GST fusion form, and purified by using glutathione-Sepharose followed by thrombin cleavage as described by Kreck et al. [12]. Recombinant proteins p47^{phox} and wild-type p67^{phox} were expressed in insect cells (sf9 cell) as described previously [10]. A series of truncated p67^{phox} and their mutants were expressed in *E. coli*, were purified with glutathione-Sepharose followed by glutathione elution as described previously [10], and were dialyzed to remove free glutathione. Protein concentrations were determined according to Bradford [3]. The purity of the proteins were confirmed by SDS-PAGE and Coomassie Blue staining.

Truncations of p67^{phox}:

A series of truncated p67^{phox} clones were obtained by PCR using p67^{phox} DNA cloned in pGEX-2T as the template. For all PCR reactions, the forward primer (CGTGGATCC ATGTCCCTGGTGGAG GCC) was designed to anneal to 5' end of p67^{phox} sequence and to introduce a BamHI site (shown in bold) and the initiation codon (underlined). For each truncation, the reverse primer (e.g. for p67^{phox}(1-210) mutant, GATGAATTCTTAATCCACCACAGATGC) was designed to anneal to the p67^{phox} sequence immediately 5' to the region to be truncated, and to introduce the stop codon (underlined) and a EcoRI site (shown in bold). These PCR products were ligated into the BamHI and EcoRI sites of pGEX-2T vector, and were transformed into DH5a for expression of the protein. The PCR products were sequenced to verify that no unexpected mutations were introduced by PCR and to confirmed the truncations.

NADPH oxidase activity assay:

Superoxide generation was measured by SOD-inhibitable

reduction of cytochrome c as described by Burnham et al. [4] using a Thermomax Kinetic Microplate reader (Molecular Devices, Menlo Park, CA). Rac was preloaded with 5-fold molar excess of GTPγS for 15 min at room temperature in the absence of MgCl₂ as described previously [12]. For the standard assay condition, the cell-free reaction mixtures include 60 nM flavocytochrome b₅₅₈ that had been reconstituted with FAD or FAD analog and phospholipids, 800 nM p47^{phox}, 900 nM p67^{phox}, 450 nM Rac, 10 mM GTPγS, and 200-240 mM arachidonate in a total of 50 ml. Three 10 ml aliquots of each reaction mixture were transferred to 96-well assay plates and preincubated for 5 min at 25°C. For each well, 240 μl of substrate cocktail containing 200 mM NADPH and 80 mM cytochrome c in buffer A (100 mM KCl, 3 mM NaCl, 4 mM MgCl₂, 1 mM EGTA, and 10 mM PIPES, pH 7.0), was added to initiate superoxide generation. NADPH oxidase activity was measured by monitoring absorbance change at 550 nm. An extinction coefficient at 550 nm of 21 mM⁻¹ cm⁻¹ was used to calculate the quantity of cytochrome c reduced [13].

Spectrophotometric and fluorometric assays:

Heme content was determined by reduced minus oxidized difference spectroscopy at 424~440 nm using an extinction coefficient of 161 mM⁻¹ cm⁻¹ [6]. The flavin content of FAD analog-reconstituted cytochrome b₅₅₈ was estimated fluorimetrically. Fluorescence spectra were recorded with a Hitachi model F-3000 spectrofluorimeter. Fluorescence changes at 525 nm induced by NADPH-FAD analog oxidoreduction during cell-free NADPH oxidase activation occurred slowly for about 5 min, and the total fluorescence change due to the complete reduction of the FAD analog was measured by adding a few crystals of sodium dithionite. To calculate the percent reduction of the FAD analog at steady state, the fluorescence change at 525 nm attributable to NADPH oxidation was subtracted from that due to oxidoreduction of NADPH and the FAD analog. The time course of heme reduction was derived from the absorbance changes at 558 minus 540 nm, using an extinction coefficient of 21.6 mM⁻¹ cm⁻¹ [6].

Results

Effect of cytosolic factors on the reduction of fad and heme:

The steady state reduction levels were calculated based on the percent fluorescence bleaching achieved at 5 min, correcting for the decrease in fluorescence contributed by NADPH oxidation. Based on this calculation the fraction reduction of flavin after steady state has been achieved is 28 ± 3% (Table 1). In contrast to flavin reduction, addition of NADPH produced < 2% steady state reduction of heme based on absorbance changes at 558 nm minus 540 nm

Table 1. Effects of cytosolic factors on NADPH oxidase activity and on the steady state reduction of FAD and heme 8-Thioacetamido-FAD was reconstituted into purified cytochrome b₅₅₈ in the presence of phospholipids as described in "Materials and Methods". NADPH-dependent superoxide generation was monitored in the presence or absence of p47^{phox} and p67^{phox}.

Components	NADPH oxidase activity	Steady state reduction level	Heme
	(nmol/min/nmol of heme)	FAD analog	
Complete	320 ± 50	28 ± 3%	<2%
-p47 ^{phox}	210 ± 30	21 ± 2%	<2%
-p67 ^{phox}	0	4 ± 1%	<2%

(Table 1). The steady state percent reduction of the FAD analog and heme was determined as above in the complete system or in the absence of either p47^{phox} or p67^{phox} (Table 1). When p47^{phox} was omitted, there was still significant reduction of FAD (21% compared with 28%). However, when p67^{phox} was omitted, FAD was almost completely oxidised (Table 1). The steady state of reduction of FAD correlated with the rate of superoxide generation under the same conditions, indicating a functional relationship between FAD reduction and superoxide generation (Table 1). In contrast, heme was completely oxidised regardless of the presence of the cytosolic factors (Table 1).

Expression of truncated p67^{phox}:

A series of truncated mutant p67^{phox} (Fig. 1) was generated to determine the region which is important for regulating the steady state FAD reduction. As shown in Fig. 2, p67^{phox} (1-246) partially (approximately 50% of V_{max}) activates flavocytochrome b₅₅₈ which is consistent with previous observation [10]. Further truncated mutants p67^{phox} (1-235), p67^{phox} (1-221), p67^{phox} (1-216), and p67^{phox} (1-210) thoroughly regain their abilities for activating NADPH oxidase almost same as wild-type p67^{phox} (Fig. 2). Further




Fig 1. Truncation of p67^{phox} and its effect on NADPH oxidase activation. Various domains of p67^{phox}, including two SH3 (src homologous region 3) domains, a Rac-binding domain (RBD), and the region from amino acid residues 198 to 246 (hatched). This region is expanded to show the amino acid sequence and residue number. The activation domain is underlined




Fig 2. NADPH oxidase activation by truncated p67^{phox}. Superoxide generation was measured as described under Materials and Methods. The reaction mixture was consisted of 60 nM flavocytochrome b₅₅₈ that had been reconstituted with FAD and phospholipids. In cubations contained 600 nM p47^{phox}, 450 nM Rac1, 900 nM of truncated p67^{phox}, and 0.2 mM arachidonate. Error bars show the standard error of the mean (n = 3).




Fig 3. Effect of truncated p67^{phox} on the steady state reduction of FAD. The reaction mixture was consisted of 60 nM of FAD analog reconstituted flavocytochrome b₅₅₈, 800 nM p47^{phox}, 450 nM Rac1, 900 nM of truncated p67^{phox}, and 0.2 mM arachidonate. The fluorescence emission spectrum of 525 nm (excitation wavelength, 475 nm) was recorded as described in Materials and Methods. Error bars show the standard error of the mean (n = 3).

truncated mutants, p67^{phox} (1-204) and p67^{phox} (1-198), dramatically reduces superoxide generation, which suggests that p67^{phox} (1-210) is the minimal-size active domain, and the region 199~210 of p67^{phox} is critical for activating flavocytochrome b₅₅₈ in cell-free oxidase reconstitution. Therefore, the activation domain is important for regulating

electron flow within flavocytochrome b_{558} , and data suggest that it does so by interacting directly with this catalytic component.



Role of the activation domain in $p67^{phox}$ in FAD reduction:

As shown in Fig. 2 and 3, $p67^{phox}$ derivatives which have an activation domain showed both higher rate of superoxide generation and higher level of steady state reduction of 8-thioacetamido-FAD. However, the truncation of the activation domain resulted in much lower rate of superoxide generation (Fig. 2) and a very low steady state reduction of 8-thioacetamido-FAD (Fig. 3). The activation domain is not involved in the interaction with Rac1 or $p47^{phox}$ [10]. Therefore, the truncation of the activation domain suppresses the reduction of FAD by NADPH in flavocytochrome b_{558} .

Discussion

Based on the sequence homologies between $p67^{phox}$ and the putative pyridine nucleotide-binding sites of NADPH-dependent enzymes, the 193~212 amino acid region of $p67^{phox}$ was proposed as the one of the candidates for NADPH-binding site [20]. NADPH-binding site on the b subunit of flavocytochrome b_{558} ($gp91^{phox}$) was also postulated on the basis of sequence homologies; alignment of the amino acid sequence of $gp91^{phox}$ with other flavoprotein revealed that five peptide segments in the 403~570 amino acid region of $gp91^{phox}$ are likely to be NADPH-binding domain [16, 19]. The docking site of $p67^{phox}$ on flavocytochrome b_{558} is still unknown. Therefore, one of the possible role of the 201~210 amino acid region is transferring NADPH from cytosol to the substrate binding site of $gp91^{phox}$ to form [E-S] complex by opening the NADPH-binding site in $gp91^{phox}$.

A model has been proposed that attempts to explain individual roles for cytosolic factors during the protein assembly associated with activation of the respiratory burst (see Introduction). According to this model, it is $p67^{phox}$ that directly regulates the rate-limiting transfer of electrons within the $gp91^{phox}$ subunit through its activation domain within the 199~210 region. In the present study, we have investigated the influence of this region on regulating the rate of specific catalytic steps involved in transferring electrons from NADPH to O_2 . The reductive half-reaction (Reaction 1) and reoxidative half-reaction (Reaction 2) with respect to FAD within $gp91^{phox}$ are summarized as follows.



We first used steady state kinetics to investigate whether the activation domain in $p67^{phox}$ stimulates the reductive

half-reaction (Reaction 1) or the reoxidative half-reaction (Reaction 2). If the former were the case then the $p67^{phox}$ should increase the steady state reduction level of FAD, and truncations should lead to a more oxidized state. The opposite should be true if $p67^{phox}$ were functioning as an activator for Reaction 1. In addition, the heme should become more reduced. Thus, monitoring the steady state reduction of flavin and heme during turnover will distinguish between these two models.

We propose that the activation domain on $p67^{phox}$ was critical for regulating FAD reduction, since the deletion of this domain not only decreased the superoxide generation but also decreased the steady state of FAD reduction. Thus, the activation domain on $p67^{phox}$ regulates the reductive half-reaction for FAD (Reaction 1), consistent with a dominant effect on hydride/electron transfer from NADPH to FAD.

References

1. Abo A, Boyhan A, West I, Thrasher AJ, and Segal AW. Reconstitution of neutrophil NADPH oxidase activity in the cell-free system by four components: $p67^{phox}$, $p47^{phox}$, $p21^{rac1}$, and cytochrome b_{245} . *J Biol Chem.* 1992, **267**(24), 16767-70.
2. Badwey JA, and Karnovsky ML. Active oxygen species and the functions of phagocytic leukocytes. *Annu Rev Biochem.* 1980, **49**, 695-726.
3. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976, **72**, 248-54.
4. Burnham DN, Uhlinger DJ, and Lambeth JD. Diradylglycerol synergizes with an anionic amphiphile to activate superoxide generation and phosphorylation of $p47^{phox}$ in a cell-free system from human neutrophils. *Biol Chem.* 1990, **265**(29), 17550-9.
5. Clark RA. The human neutrophil respiratory burst oxidase. *J Infect Dis.* 1990, **161**(6), 1140-7.
6. Cross AR, Higson FK, Jones OT, Harper AM, and Segal AW. The enzymic reduction and kinetics of oxidation of cytochrome b_{245} of neutrophils. *Biochem J.* 1982, **204**(2), 479-85.
7. De Leo FR, Ulman KV, Davis AR, Jutila KL, and Quinn MT. Assembly of the human neutrophil NADPH oxidase involves binding of $p67^{phox}$ and flavocytochrome b to a common functional domain in $p47^{phox}$. *J Biol Chem.* 1996, **271**(29), 17013-20.
8. Dorseuil O, Quinn MT, and Bokoch GM. Dissociation of Rac translocation from $p47^{phox}$ / $p67^{phox}$ movements in human neutrophils by tyrosine kinase inhibitors. *J Leukoc Biol.* 1995, **58**(1), 108-13.
9. Fuchs A, Dagher MC, and Vignais PV. Mapping the domains of interaction of $p40^{phox}$ with both $p47^{phox}$ and $p67^{phox}$ of the neutrophil oxidase complex using the two-hybrid system. *J Biol Chem.* 1995, **270**(11), 5695-7.

10. **Han CH, Freeman JL, Lee T, Motalebi SA, and Lambeth JD.** Regulation of the neutrophil respiratory burst oxidase. Identification of an activation domain in p67(phox). *J Biol Chem.* 1998, **273(27)**, 16663-8.
11. **Heyworth PG, Bohl BP, Bokoch GM, and Curnutte JT.** Rac translocates independently of the neutrophil NADPH oxidase components p47^{phox} and p67^{phox}. Evidence for its interaction with flavocytochrome b₅₅₈. *J Biol Chem.* 1994, **269(49)**, 30749-52.
12. **Kreck ML, Uhlinger DJ, Tyagi SR, Inge KL, and Lambeth JD.** Participation of the small molecular weight GTP-binding protein Rac1 in cell-free activation and assembly of the respiratory burst oxidase. Inhibition by a carboxyl-terminal Rac peptide. *J Biol Chem.* 1994, **269(6)**, 4161-8.
13. **Lambeth JD, Burnham DN, and Tyagi SR.** Sphinganine effects on chemoattractant-induced diacylglycerol generation, calcium fluxes, superoxide production, and on cell viability in the human neutrophil. Delivery of sphinganine with bovine serum albumin minimizes cytotoxicity without affecting inhibition of the respiratory burst. *J Biol Chem.* 1988, **263(8)**, 3818-22.
14. **Nisimoto Y, Otsuka-Murakami H, and Lambeth DJ.** Reconstitution of flavin-depleted neutrophil flavocytochrome b₅₅₈ with 8-mercapto-FAD and characterization of the flavin-reconstituted enzyme. *J Biol Chem.* 1995, **270(27)**, 16428-34.
15. **Roos D, de Boer M, Kuribayashi F, Meischl C, Weening RS, Segal AW, Ahlin A, Nemet K, Hossle JP, and Bernatowska-Matuszkiewicz E, Middleton-Price H.** Mutations in the X-linked and autosomal recessive forms of chronic granulomatous disease. *Blood.* 1996, **87(5)**, 1663-81.
16. **Rotrosen D, Yeung CL, Leto TL, Malech HL, and Kwong CH.** Cytochrome b₅₅₈: the flavin-binding component of the phagocyte NADPH oxidase. *Science.* 1992, **256(5062)**, 1459-62.
17. **Rotrosen D, Yeung CL, and Katkin JP.** Production of recombinant cytochrome b₅₅₈ allows reconstitution of the phagocyte NADPH oxidase solely from recombinant proteins. *J Biol Chem.* 1993, **268(19)**, 14256-60.
18. **Sathyamoorthy M, de Mendez I, Adams AG, and Leto TL.** p40(phox) down-regulates NADPH oxidase activity through interactions with its SH₃ domain. *J Biol Chem.* 1997, **272(14)**, 9141-6.
19. **Segal AW, West I, Wientjes F, Nugent JH, Chavan AJ, Haley B, Garcia RC, Rosen H, and Scrace G.** Cytochrome b-245 is a flavocytochr phagocytes. *Biochem J.* 1992, **284(Pt 3)**, 781-8.
20. **Smith RM, Connor JA, Chen LM, and Babior BM.** The cytosolic subunit p67^{phox} contains an NADPH-binding site that participates in catalysis by the leukocyte NADPH oxidase. *J Clin Invest.* 1996, **98(4)**, 977-83.
21. **Someya A, Nagaoka I, Yamashita T.** Purification of the 260 kDa cytosolic complex involved in the superoxide production of guinea pig neutrophils. *FEBS Lett.* 1993, **330(2)**, 215-8.
22. **Taylor WR, Jones DT, Segal AW.** A structural model for the nucleotide binding domains of the flavocytochrome b-245 beta-chain. *Protein Sci.* 1993, **2(10)**, 1675-85.