



# 14q12q13.3 Deletion Diagnosed Using Chromosomal Microarray Analysis in an Infant Showing Seizures, Hypoplasia of the Corpus Callosum, and Developmental Delay

Jae Hyuk Kwon, MD<sup>1</sup>, Young Hwa Song, MD<sup>1</sup>, Jung Min Yoon, MD<sup>1</sup>, Eun Jung Cheon, MD<sup>1</sup>, Kyung Ok Ko, MD<sup>1</sup>, Jae Woo Lim, MD<sup>1</sup>, and Hyon J. Kim, MD, FACMG<sup>2</sup>

Departments of <sup>1</sup>Pediatrics and <sup>2</sup>Medical Genetics, Konyang University College of Medicine, Daejeon, Korea

## ABSTRACT

14q12q13.3 Deletion is a rare microdeletion syndrome associated with neurodevelopmental delay, failure to thrive, seizures, and abnormal brain development. Symptoms vary depending on the sites of gene deletion, and establishing the diagnosis is often difficult, as the condition cannot be detected with routine chromosome analysis. In this report, we present a patient with intrauterine growth retardation, microcephaly, muscle weakness, seizures, and hypoplasia of the corpus callosum who underwent diagnostic tests, including karyotyping in the neonatal period without leading to a specific diagnosis. The patient was confirmed with a serious developmental disorder, and a chromosomal microarray analysis was performed at 8 months of age, revealing a 14q12q13.3 deletion. In this case, the condition was diagnosed in early infancy, in contrast to previously reported cases, and the patient had diverse and severe symptoms. Establishing the diagnosis of 14q12q13.3 deletion syndrome allows better management of patient care and genetic counseling for the parents.

**Key Words:** Microdeletion 14q12q13.3; Chromosomal microarray; Developmental delay; Failure to thrive; Genetic counseling

## INTRODUCTION

14q12q13 Microdeletion syndrome is a major chromosome anomaly associated with neurodevelopmental delay and learning disability, as well as microcephaly, failure to thrive, seizure, and abnormal development of part of the brain<sup>1)</sup>. The symptoms may vary depending on the gene deleted. Deletion of the Forkhead box protein G1 (*FOXP1*) gene manifests as a neurodevelopmental disability or severe developmental delay, while a deletion of NK2 homeobox 1 (*NKX2-1*) has been associated with the brain-lung-thyroid syndrome and has

Received: 18 August 2020

Revised: 25 September 2020

Accepted: 6 October 2020

**Correspondence to:** Hyon J. Kim, MD  
Department of Medical Genetics,  
Konyang University College of Medicine,  
158 Gwanjeodong-ro, Seo-gu,  
Daejeon 35365, Korea

**Tel:** +82-42-600-9230

**Fax:** +82-42-600-9090

**E-mail:** [raredisease@hanmail.net](mailto:raredisease@hanmail.net)

Copyright(c)

By Korean Society of Neonatology.

All right reserved.

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

25 weeks of gestation. Although no abnormalities were found during prenatal care, the amniotic fluid index checked on 35 weeks of gestation was 5.1 cm, based on which oligohydramnios was suspected, and the fetus showed intrauterine growth delay. The results of placental biopsy conducted postpartum were normal. The father of the baby was 29 years of age, without any notable family or medical history (Figure 1). The baby's Apgar score at birth was 7 at 1 minute and 9 at 5 minutes. The baby showed symmetrical intrauterine growth retardation, with a birth weight below the 3rd percentile and a head circumference (31 cm) and height (44 cm) below the 10th percentile. The baby had small capillary hemangiomas at the center of the forehead and posterior scalp, a simian line on the right hand, and slightly diminished muscle tone. He had stridor and tachypnea requiring oxygen via a nasal cannula, and nasogastric tube feeding was initiated due to tachypnea. The first blood test taken on the day of the birth showed a white blood cell count of  $7,600/\text{mm}^3$ , a platelet count of  $116,000/\text{mm}^3$ , and a C-reactive protein (CRP) level of 3.5 mg/dL, raising suspicion for early sepsis; thus, antibiotic therapy was initiated. On hospital day (HD) 6, the platelet count dropped to  $35,000/\text{mm}^3$ . The prothrombin time (PT), activated partial thromboplastin time (aPTT), and international normalized ratio (INR) were 16.3 seconds, 46.9 seconds, and 1.30, respectively, indicating slightly increased PT with aPTT and INR within the normal ranges. Antithrombin III was 46%, D-dimer/fibrin degradation product tests were positive, and systemic bruising was noted. Therefore, platelet concentrate and fresh frozen plasma were transfused, and antithrombin was administered. On HD 12, no bacteria were identified in the blood culture, the platelet count increased to  $104,000/\text{mm}^3$ , and the CRP level improved, leading to the discontinuation of antibiotic therapy. Tests for congenital infection, such as cytomegalovirus urine polymerase chain reaction and TORCH immunoglobulin M level, were normal. Oral feeding was attempted on HD 6; however, the baby was not feeding well, because he was not strong enough to suck the bottle. Therefore, central venous parenteral nutrition was started. On HD 15, the baby had a seizure, with stiffening and shaking of the extremities and was placed on a ventilator due to repeated episodes of apnea and reduced oxygen saturation. An electroencephalogram (EEG) showed moderate encephalopathy with frequent negative sharp waves on both central regions. Seizures were poorly controlled with phenobarbital; therefore, topiramate was administered. Brain computed tomography performed on HD 18 revealed suspected hypoplasia of the corpus callosum.


Ventilator use was discontinued on HD 22 after the baby's seizures ceased and his breathing stabilized. Hypoplasia of the corpus callosum was confirmed on brain magnetic resonance imaging (MRI), with no additional anomalies discovered (Figure 2). As the baby had symmetrical intrauterine growth retardation, simian line, hypoplasia of the corpus callosum, and demonstrated muscle weakness and seizure, routine chromosomal karyotyping was performed using peripheral blood. This confirmed the baby to be a male with normal 46 XY chromosomes. Echocardiography, renal ultrasonography, and eye examination performed to reveal any other anomalies showed normal findings, and diagnostic auditory brainstem response confirmed normal hearing in the left ear but mild hearing loss with a hearing threshold of 40 dB in the right ear.

While the initial thyroid stimulating hormone (TSH) level on the neonatal metabolic screening test (NST) was normal, a second NST taken after the seizure showed an elevated TSH level of 17.1  $\mu$ IU/mL. On HD 20, hypothyroidism was diagnosed with a TSH level of 37.80  $\mu$ IU/mL, free thyroxine level of 0.92 ng/dL, and triiodothyronine level of 1.06 ng/dL. The thyroglobulin level was elevated to 586.59 ng/mL, and anti-thyroglobulin antibody (Ab) and anti-thyroid peroxidase (anti-TPO) Ab were negative. Thyroid ultrasonographic findings were unremarkable. Under the diagnosis of hypothyroidism, the baby was administered 10  $\mu$ g/kg of levothyroxine, after which the thyroid function test results were normal.

The baby gained weight to 2.92 kg with increased oral feeding, and the seizures were controlled. The baby was diagnosed with

seizure and transient hypothyroidism and was discharged on HD 54 with continued phenobarbital, topiramate, and levothyroxine therapy. At this time, the baby continued to show muscle weakness equivalent to a level 4 grading using the Manual Muscle Testing method and was scheduled for rehabilitation and follow-up in the outpatient clinic.

Two days after discharge, the baby was brought to the emergency department for poor feeding and lethargy. He was administered intravenous fluid therapy and discharged. Subsequently, he was brought to the hospital several times due to irritability and repeated hypertonia until 4 months of age, and each time, conservative treatment was conducted. He had no seizures and a normal EEG at 4 months of age; therefore, anticonvulsants were ceased. At 5 months of age, the baby was diagnosed with failure to thrive with a weight of 4.6 kg (<1st percentile), height of 50 cm (<1st percentile), head circumference of 39 cm (<3rd percentile), hypertonia, sleep disturbance with irritability, and constipation. The baby developed some neck strength but could not make eye contact. On the Bayley Scales of infant and toddler development, 3rd edition, at 6 months of age his cognition composite score was 55 (0.1st percentile), language composite score was 59 (0.3st percentile), and motor composite score was 46 (<0.1st percentile), which revealed severe developmental delay equivalent to the development of a 1-month-old. At 8 months of age, the baby was referred to a Genetic Counseling Clinic and underwent genetic evaluation including pedigree analysis, which was essentially negative except diabetic mellitus in maternal great aunts, and chromosomal microarray testing. The test detected an



**Figure 2.** Brain magnetic resonance imaging findings. (A) Sagittal T1-weighted image showing hypoplasia of the corpus callosum (white arrow). (B) Coronal T2-weighted image showing the “bat wings” appearance of the lateral ventricles (black arrows).


11.5 Mb deletion in chromosome 14q [14q12q13.3 (25,363,718-36,872,996)×1] (Figure 3). At 19 months of age, he again began to have seizures with the arms and legs flexed. Valproate therapy was initiated followed by topiramate, when valproate alone could not control the seizures. It was difficult to increase the amount of feeding, as he was able to feed using a bottle but had difficulty swallowing. Further, he was frequently given antibiotic therapy due to repeated respiratory infections. The Denver developmental screening test conducted at 25 months of age revealed that the child was at 3 months for large motor development, 2 months for personal-social, and 3 months for language development. Thyroid function tests revealed normal findings until 36 months of age, and levothyroxine was stopped. However, after 2 weeks, his TSH level rose again to 18.34  $\mu$ IU/mL; therefore, levothyroxine therapy was resumed, resulting in a level within the normal range. At 41 months of age, the baby's weight was 7.7 kg (<1st percentile), height was 72 cm (<1st percentile), and head circumference was 44 cm (<1st percentile). The child was still only able to feed from a bottle, with weight gain stalled, so enteric nutrients were added for calorie supplementation. Pharmacological therapy for seizures and hypothyroidism was continued. His large motor and language development were equivalent to those of a 3-month-old, where he developed neck strength, was able to lift his head 45 degrees when placed on his stomach, and was able to produce only meaningless sounds.

## DISCUSSION

Approximately 2% of neonates born worldwide have a major anomaly<sup>5</sup>, 10% of which are known to result from a chromosome abnormality<sup>6</sup>. The incidence is gradually on the rise due to the increasing age of mothers at childbirth, and the increase in *in vitro* fertilization<sup>7</sup>. The banding karyotyping technique is useful for confirming abnormalities resulting from chromosome translocations, inversions, ring chromosomes, marker chromosomes, and deletions<sup>8,9</sup>. However, more than half of neonatal chromosomal abnormalities cannot be detected by physical examination or banding karyotyping<sup>8,10,11</sup>, and these are cases in which a sex chromosomal abnormality has not yet manifested or cases of microdeletion syndrome that are discovered via chromosomal microarray analysis. Microdeletion syndrome refers to cases in which the chromosomal defect is smaller than 5 Mb, which is difficult to detect even with a high-resolution banding technique. Chromosomal microarray analysis has a higher sensitivity to microdeletions, subtelomeres, and pericentric regions than routine karyotyping and is thus known to be conducive to the diagnosis of microdeletion or microduplication syndrome<sup>12,13</sup>.

The present case is of an infant with intrauterine growth retardation, microcephaly, muscle weakness, seizures, hypothyroidism, and hypoplasia of the corpus callosum who showed severe growth and neurodevelopmental abnormalities during follow-up and was diagnosed with 14q12q13.3 deletion via chromosomal microarray analysis.

The patient in this case had 14q12q13.3 deletion, which in-



**Figure 3.** Chromosomal microarray data from the patient showing the chromosome 14q12q13.3 deletion (red inverse triangle).


cludes the *FOXG1*, *COCHAP4S1*, *CFL2*, *PSMA6*, *NFKBIA*, and *NKX2-1* genes (Figure 4). Accordingly, he showed microcephaly, structural anomaly of the corpus callosum, seizures, and severe developmental delays, which are observed with *FOXG1* gene anomalies<sup>14,15</sup>, along with neonatal hypotonicity that progresses to hypertonia, developmental delay, and seizures, which are observed with *AP4S1* mutation<sup>16</sup>.

Furthermore, while the baby had normal findings on the initial NST after birth, a thyroid function test performed after the onset of seizures confirmed hypothyroidism. The *NKX2-1* gene is located on 14q13.3 and is involved in the formation of homeobox protein Nkx-2.1; therefore, this gene affects thyroid hormone production. In the present case, we speculate that the patient's hypothyroidism is caused by the abnormality of this gene<sup>17</sup>.

The *NFKBIA* gene is associated with ectodermal dysplasia; affects hair, teeth, and sweat gland development; and results in T cell immunodeficiency<sup>18</sup>. The child in the present case had no dental development and experienced repeated respiratory

infections. *COCH*, *CFL2*, and *PSMA6* are associated with hearing abnormality, nemaline myopathy, and ophthalmologic diseases, respectively, and the patient in the present case did not show these symptoms.

Existing descriptions of 14q12q13.3 microdeletion are those reported by Fonseca et al.<sup>19</sup> and Ponzi et al.<sup>1)</sup>, and we compared the clinical manifestations with these cases (Table 1). Fonseca et al.<sup>19</sup> reported various symptoms including growth and developmental delays. In this case, EEG showed abnormal focal theta semi-rhythmic activity, and there were no abnormal findings on brain MRI<sup>19</sup>. The patient in Ponzi et al.'s report<sup>1)</sup> had a high birth weight of greater than the 97th percentile and hypotonia, sleep disturbance, and attention problems. This patient had normal EEG, and brain MRI revealed hypoplasia of the corpus callosum including posterior portion abnormalities of the lateral ventricles<sup>1)</sup>. The patient in our case differed in that he was a preterm infant who had showed seizures and hypotonia since the neonatal period, which progressed to hypertonia in



**Figure 4.** Schematic representation of the 14q12q13.3 overlapping deletions in this patient. The key genes included in the deleted region are marked with a red box.

**Table 1.** Comparison of Clinical and Genetic Data of the 14q12q13.3 Microdeletion Cases Reported in Literature and Our Case

| Variable                       | Fonseca et al. <sup>19)</sup>         | Ponzi et al. <sup>1)</sup>            | Present case                          |
|--------------------------------|---------------------------------------|---------------------------------------|---------------------------------------|
| Breakpoint nucleotide position | 14q12q13.3<br>(35,520,645-37,271,596) | 14q12q13.3<br>(32,217,109-35,157,847) | 14q12q13.3<br>(25,363,718-36,872,996) |
| Deletion size                  | 5.264 Mb                              | 2.94 Mb                               | 11.5 Mb                               |
| Inheritance                    | <i>De novo</i>                        | <i>De novo</i>                        | <i>De novo</i>                        |
| Sex                            | Female                                | Male                                  | Male                                  |
| Gestation                      | 42 wk                                 | Full-term                             | 36 wk                                 |

infancy. In addition, compared to the other cases, he had severe growth and neurodevelopmental delay, and both EEG and brain MRI showed abnormal findings. All three cases were *de novo* mutations with no parental influence.

Clinical manifestations of microdeletion syndrome vary widely depending on the genes included in the deleted regions of the chromosome. The present case was also difficult to diagnose due to the presence of multiple comorbidities, and a specific diagnosis was not made despite several tests, including chromosomal karyotyping in the neonatal period. However, through continuous follow-up, we were able to confirm progressive developmental and growth delay and perform chromosomal microarray analysis at an early age of 8 months. Nowadays, chromosomal microarray analysis is an essential test, and important for confirming the diagnosis of multiple congenital anomalies and neurodevelopmental disabilities during the neonatal period. Further, the present case had a large deletion that resulted in a wide array of symptoms with severe developmental and growth delay compared to previous cases. Thus, the specific diagnosis of 14q12q13.3 microdeletion syndrome in this case enabled early intervention for better management of the patient and genetic counseling for the parents.

## ARTICLE INFORMATION

### Ethical statement

This case report was reviewed and approved by the Institutional Review Board of Konyang University Hospital (KYUH 2020-07-021). Informed consent was waived by the board.

### Conflicts of interest

No potential conflict of interest relevant to this article was reported.

### Author contributions

Conception of design: J.H.K., Y.H.S., J.W.L., H.J.K.

Acquisition, analysis, or interpretation of data: J.H.K., H.J.K.

Drafting the work or revising: Y.H.S., J.M.Y., E.J.C., K.O.K., J.W.L., H.J.K.

Final approval of the manuscript: H.J.K.

### ORCID

Jae Hyuk Kwon <https://orcid.org/0000-0002-1682-3218>

Hyon J. Kim <https://orcid.org/0000-0002-2945-8731>

### Acknowledgments

This study was funded by the Genetic Counseling Service Support Funding from the Korean Foundation for Rare Disease.

## REFERENCES

1. Ponzi E, Gentile M, Agolini E, Matera E, Palumbi R, Buonadonna AL, et al. 14q12q13.2 microdeletion syndrome: clinical characterization of a new patient, review of the literature, and further evidence of a candidate region for CNS anomalies. *Mol Genet Genomic Med* 2020;8:e1289.
2. Santen GW, Sun Y, Gijsbers AC, Carre A, Holvoet M, Haeringen Av, et al. Further delineation of the phenotype of chromosome 14q13 deletions: (positional) involvement of FOXG1 appears the main determinant of phenotype severity, with no evidence for a holoprosencephaly locus. *J Med Genet* 2012;49:366-72.
3. Resta N, Memo L. Chromosomal microarray (CMA) analysis in infants with congenital anomalies: when is it really helpful? *J Matern Fetal Neonatal Med* 2012;25 Suppl 4:124-6.
4. Miller DT, Adam MP, Aradhya S, Biesecker LG, Brothman AR, Carter NP, et al. Consensus statement: chromosomal microarray is a first-tier clinical diagnostic test for individuals with developmental disabilities or congenital anomalies. *Am J Hum Genet* 2010;86:749-64.
5. Feldkamp ML, Carey JC, Byrne JL, Krikov S, Botto LD. Etiology and clinical presentation of birth defects: population based study. *BMJ* 2017;357:j2249.
6. Brent RL. Addressing environmentally caused human birth defects. *Pediatr Rev* 2001;22:153-65.
7. Mai CT, Isenburg JL, Canfield MA, Meyer RE, Correa A, Alverson CJ, et al. National population-based estimates for major birth defects, 2010-2014. *Birth Defects Res* 2019;111:1420-35.
8. Hook EB. Contribution of chromosome abnormalities to human morbidity and mortality. *Cytogenet Cell Genet* 1982;33:101-6.
9. Schinzel A. Catalog of unbalanced chromosome aberrations in man. 2nd ed. Berlin: Walter de Gruyter, 2001.
10. Emy Dorfman L, Leite JC, Giugliani R, Riegel M. Microarray-based comparative genomic hybridization analysis in neonates with congenital anomalies: detection of chromosomal imbalances. *J Pediatr (Rio J)* 2015;91:59-67.
11. Pickering DL, Eudy JD, Olney AH, Dave BJ, Golden D, Stevens J, et al. Array-based comparative genomic hybridization analysis of 1176 consecutive clinical genetics investigations. *Genet Med* 2008;10:262-6.
12. Ellison JW, Ravnan JB, Rosenfeld JA, Morton SA, Neill NJ, Williams MS, et al. Clinical utility of chromosomal microarray

- analysis. *Pediatrics* 2012;130:e1085-95.
13. Lalani SR. Current genetic testing tools in neonatal medicine. *Pediatr Neonatol* 2017;58:111-21.
14. Florian C, Bahi-Buisson N, Bienvenu T. FOXP1-related disorders: from clinical description to molecular genetics. *Mol Syndromol* 2012;2:153-63.
15. Kortum F, Das S, Flindt M, Morris-Rosendahl DJ, Stefanova I, Goldstein A, et al. The core FOXP1 syndrome phenotype consists of postnatal microcephaly, severe mental retardation, absent language, dyskinesia, and corpus callosum hypogenesis. *J Med Genet* 2011;48:396-406.
16. Hardies K, May P, Djemie T, Tarta-Arsene O, Deconinck T, Craiu D, et al. Recessive loss-of-function mutations in AP4S1 cause mild fever-sensitive seizures, developmental delay and spastic paraplegia through loss of AP-4 complex assembly. *Hum Mol Genet* 2015;24:2218-27.
17. Gentile M, de Mattia D, Pansini A, Schettini F, Buonadonna AL, Capozza M, et al. 14q13 distal microdeletion encompassing NKX2-1 and PAX9: patient report and refinement of the associated phenotype. *Am J Med Genet A* 2016;170:1884-8.
18. McDonald DR, Mooster JL, Reddy M, Bawle E, Secord E, Geha RS. Heterozygous N-terminal deletion of IkappaBalpha results in functional nuclear factor kappaB haploinsufficiency, ectodermal dysplasia, and immune deficiency. *J Allergy Clin Immunol* 2007;120:900-7.
19. Fonseca DJ, Prada CF, Siza LM, Angel D, Gomez YM, Restrepo CM, et al. A de novo 14q12q13.3 interstitial deletion in a patient affected by a severe neurodevelopmental disorder of unknown origin. *Am J Med Genet A* 2012;158A:689-93.