



Ameliorative Effects of Ombuoside on Dopamine Biosynthesis in PC12 Cells

Uchralsaikhan Davaasambuu¹, Keun Hong Park¹, Hyun Jin Park^{1,2}, Hyun Sook Choi³, Chong Kil Lee^{1,2}, Bang Yeon Hwang¹, and Myung Koo Lee^{1,2,*}

¹College of Pharmacy and ²Research Center for Bioresource and Health, Chungbuk National University, Cheongju 28160, Korea

³Department of Food and Nutrition, Chungcheong University, Cheongju 28171, Republic of Korea

Abstract – This study investigated the effects of ombuoside, a flavonol glycoside, on dopamine biosynthesis in PC12 cells. Ombuoside at concentrations of 1, 5, and 10 µM increased intracellular dopamine levels at 1 - 24 h. Ombuoside (1, 5, and 10 µM) also significantly increased the phosphorylation of tyrosine hydroxylase (TH) (Ser40) and cyclic AMP-response element binding protein (CREB) (Ser133) at 0.5 - 6 h. In addition, ombuoside (1, 5, and 10 µM) combined with L-DOPA (20, 100, and 200 µM) further increased intracellular dopamine levels for 24 h compared to L-DOPA alone. These results suggest that ombuoside regulates dopamine biosynthesis by modulating TH and CREB activation in PC12 cells.

Keywords – Ombuoside, Dopamine biosynthesis, TH, CREB, PC12 cells

Introduction

Dopamine levels in the brain are associated with several devastating diseases such as Parkinson's disease (PD), Alzheimer's disease, schizophrenia, and affective disorders.¹ In dopamine biosynthetic pathways, tyrosine hydroxylase (EC 1.14.16.2; TH), the rate-limiting enzyme, catalyzes the conversion of L-tyrosine to L-3,4-dihydroxyphenylalanine (L-DOPA).² TH activity is mainly regulated via the cyclic AMP (cAMP)-cAMP-dependent protein kinase A (PKA)-cAMP-response element binding protein (CREB) pathway.^{3,4} The long-term regulation of TH is also involved in TH gene transcription and induction of TH mRNA, which is regulated by the cAMP-PKA-CREB system.⁴⁻⁶ TH at Ser40 is a major phosphorylation residue.^{5,6} In addition, L-DOPA is the most effective and frequently prescribed therapy for controlling symptoms of PD.^{1,8} L-DOPA also increases intracellular levels of dopamine in PC12 cells,^{9,10} which may be applied in the *in vitro* model of PD.

Gynostemma pentaphyllum (Cucurbitaceae) has been used as a herbal tea and contains many types of gynosaponins (GPS), flavonoids, polysaccharides, vitamins, and amino acids.¹¹ The ethanol extract (80%) from *G*

pentaphyllum (GP-EX) exerts ameliorating effects on chronic stress-induced anxiety in mice.¹² GP-EX and GPS have shown protective effects against neurotoxicity by reducing TH neuronal cell death and L-DOPA-induced dyskinesia in rat models of PD.^{13,14} GP-EX and GPS also show anxiolytic effects on affective disorders in an MPTP-lesioned mouse model of PD.¹⁵

In addition, ombuoside (7,4'-di-O-methylquercetin-3-O-beta-rutinoside) is one of the flavonol glycoside components of GP-EX.¹¹ Flavonoids exhibit a variety of biological activities, such as antioxidant, antimicrobial, anti-inflammation, cytotoxic, and anti-allergy effects by scavenging free radicals and reactive oxygen species.¹⁶⁻¹⁹

PC12, rat adrenal pheochromocytoma, cells have been widely used to investigate dopamine biosynthesis and L-DOPA-induced oxidative cytotoxicity.^{9,10,20} In this study, the effects of ombuoside on dopamine biosynthesis in PC12 cells were investigated to examine whether ombuoside is a beneficial bioactive component of GP-EX, similar to GPS.

Experimental

Materials – Ombuoside (purity > 97.0%) was obtained from BioBioPhar Co. Ltd. (Kunming, Yunnan). L-DOPA, isoproterenol, and 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) were purchased from Sigma-Aldrich (St. Louis, MO). RPMI1640 medium, donor-horse serum, fetal bovine serum, and antibiotics

*Author for correspondence

Myung Koo Lee, College of Pharmacy and Research Center for Bioresource and Health, Chungbuk National University, Cheongju 28160, Korea.
Tel: +82-43-261-2822; E-mail: myklee@chungbuk.ac.kr

were purchased from Gibco BRL (Grand Island, NY). Primary antibodies against TH, phospho-TH (Ser40), CREB, phospho-CREB (Ser133), and β -actin were also purchased from Cell Signaling Technology (Danvers, MA). All other chemicals were of reagent grade.

Cell culture – PC12 cells were grown in an RPMI 1640 medium supplemented with 10% heat-inactivated horse serum, 5% heat-inactivated fetal bovine serum, penicillin (100 units/ml), and streptomycin (100 μ g/ml). The cells were placed in a humidified atmosphere with 5% CO₂ and 95% air at 37 °C, as previously described.²⁰

Determination of dopamine levels and cell viability – PC12 cells were treated with ombuoside for the designated time and the cells were harvested with phosphate buffered saline. Trichloroacetic acid (1 M, 100 μ l) and isoproterenol (200 pmol, internal standard) were added to pellet extract and dopamine levels were measured by an HPLC system (Toso, Tokyo, Japan) with a fluorescence detector (F1000, Hitachi, Tokyo) (Ex/Em, 350/460 nm).¹⁰ The dopamine levels were expressed as nmol/mg protein and percentage of the control group.

In addition, cell viability was evaluated using a conventional MTT assay,²¹ with a Bauty Diagnostic Microplate Reader (Molecular Devices, Sunnyvale, CA).¹⁰

Western blot analysis – Analysis of the phosphorylation of TH at Ser40 [phospho-TH (Ser40)], CREB at Ser133 [phospho-CREB (Ser133)], and β -actin were determined by western blot analysis.^{4,10} Protein samples (20 μ g in each lane, 50 μ g for caspase-3) were electrophoresed and the blot analysis was conducted using primary antibodies (1:1,000 in TBS-T with 5% bovine serum albumin [BSA]) at 4 °C and secondary antibodies (1:5,000 in TBS-T with 5% BSA) according to standard procedures (Amersham Pharmacia Biotech, Inc., Piscataway, NJ) as previously described.^{4,10}

Statistical analysis – Protein amounts were determined using BSA as a standard.²² All data are expressed as the means \pm S.E.M. of at least four independent experiments. Statistical analyses were performed using analysis of variance (ANOVA) followed by Dunnett's test for multiple comparisons, and a P-value < 0.05 was considered to be statistically significant.

Result

Intracellular dopamine levels and cell viability – Ombuoside (1, 5, and 10 μ M) slightly increased intracellular dopamine levels for 1 h, and then intracellular dopamine levels were significantly increased for 3 – 24 h (increase of 122 – 136%, P < 0.05). The increase in dopamine levels




Fig. 1. Effects of ombuoside on dopamine content in PC12 cells. PC12 cells were treated with ombuoside (1, 5, and 10 μ M) and then incubated for 1 – 24 h. PC12 cells were harvested and then dopamine levels were determined by an HPLC method. The mean control levels of intracellular dopamine (0 h) were 3.78 ± 0.31 nmol/mg protein. The results are expressed as the means \pm S.E.M. ($n = 6 - 8$). *P < 0.05 compared to the control levels (ANOVA followed by Dunnett's test).

by ombuoside was treatment-, time-, and dose-dependent (Fig. 1). Ombuoside at concentrations up to 50 μ M did not alter cell viability for 24 h. However, ombuoside at 100 μ M reduced cell viability to 80.4% (P < 0.05) at 24 h compared to the control group (data not shown).

TH and CREB phosphorylation – Ombuoside (1, 5, and 10 μ M) showed significant increase in TH phosphorylation at Ser40 in a time-dependent manner (Fig. 2). TH phosphorylation started to rise at 30 min and the increase in phosphorylated TH was further enhanced by ombuoside (1, 5, and 10 μ M) for 6 h by 1.48-, 1.54-, and 1.71-fold (P < 0.05) (Fig. 2).

In addition, phosphorylation of CREB at Ser133 was induced after treatment with ombuoside (1, 5, and 10 μ M) for 0.5 – 6 h (increase by 1.44-, 1.61-, and 1.68-fold at 6 h, P < 0.05) (Fig. 3).

L-DOPA-induced dopamine levels – Intracellular levels of dopamine in PC12 cells were increased by treatment with L-DOPA at 25 – 200 μ M for 24 h (Fig. 4).¹⁰ Co-treatment with ombuoside (1, 5, and 10 μ M) and L-DOPA (20, 100, and 200 μ M) increased intracellular levels of dopamine to 148 – 176% (20 μ M, P < 0.05), 197 – 228% (100 μ M, P < 0.05), and 197 – 228% (200 μ M, P < 0.05) compared to the L-DOPA-treated group after 24 h (Fig. 4).

Discussion

GP-EX has shown protective effects on dopaminergic neurons by mediating anti-oxidative functions in animal




Fig. 2. Effects of ombuoside on the phosphorylation of TH (Ser40) in PC12 cells. PC12 cells were treated with ombuoside (1, 5, and 10 μM) for 0.5 – 6 h. Immunoblots of lysates of cell pellets were probed with phospho-TH and then the phosphorylation of TH (Ser40) (p-TH) and total TH were analyzed by western blotting. The values of the relative density ratios of p-TH/TH are expressed in arbitrary units. The results are expressed as the means ± S.E.M. ($n = 4$). *P < 0.05 compared to the control levels (ANOVA followed by Dunnett's test).




Fig. 3. Effects of ombuoside on the phosphorylation of CREB (Ser133) in PC12 cells. PC12 cells were treated with ombuoside (1, 5, and 10 μM) for 0.5 – 6 h. Immunoblots of lysates of cell pellets were probed with phospho-CREB and then the phosphorylation of CREB (Ser133) (p-CREB) and total CREB were analyzed by western blotting. The values of the relative density ratios of p-CREB/CREB are expressed in arbitrary units. The results are expressed as the means ± S.E.M. ($n = 4$). *P < 0.05 compared to the control levels (ANOVA followed by Dunnett's test).

models of PD.^{13–15} Ombuoside is a flavonol glycoside of GP-EX.¹¹ In this study, we investigated the effects of ombuoside on dopamine biosynthesis in PC12 cells.

Ombuoside at concentrations up to 50 μM did not reduce cell viability for 24 h (data not shown). Ombuoside (1, 5, and 10 μM) significantly enhanced intracellular dopamine levels in a dose-dependent manner at 6 – 24 h (Fig. 1). TH phosphorylation (Ser40) was also significantly increased after treatment with ombuoside (1, 5, and 10 μM) in a time-dependent manner (Fig. 2). TH phosphorylation (Ser40) was enhanced by ombuoside at 30 min and this increase in phosphorylated TH was maintained for 6 h. In addition, ombuoside (1, 5, and 10 μM) significantly increased the phosphorylated levels of CREB at Ser133 for 0.5 – 6 h (Fig. 3).

Dopamine biosynthesis is mainly regulated by TH

activation. Short-term activation of TH occurs through phosphorylation of TH at Ser40 by the cAMP-CREB system in dopaminergic neurons and PC12 cells^{4,7} and long-term regulation is achieved through *TH* gene expression by activating the PKA-CREB system.^{4,6} In addition, CREB is activated by phosphorylation at Ser133 through the cAMP-PKA system^{3,4,7} and then binds to the CRE region, a TH promoter, which is essential for the regulation of *TH* gene expression.^{4,7,24} These results suggest that ombuoside enhances dopamine biosynthesis through CREB and TH phosphorylation in PC12 cells.

L-DOPA at 20 – 200 μM significantly elevates the intracellular levels of dopamine by activation of the cAMP-PKA-CREB system in PC12 cells.^{9,10} However, L-DOPA at high concentration (200 μM) reduces the phosphorylation of TH, PKA, and CREB at 24 – 48 h,




Fig. 4. Effects of ombuoside on L-DOPA-induced dopamine levels in PC12 cells. PC12 cells were exposed to L-DOPA (20, 100, and 200 μ M) in the absence or presence of ombuoside (1, 5 and 10 μ M) for 24 h. PC12 cells were harvested and then dopamine levels were determined by an HPLC method. The mean control levels of intracellular dopamine were 3.56 ± 0.24 nmol/mg protein. The results are expressed as the means \pm S.E.M. ($n = 6 - 8$). * $P < 0.05$ compared to the control levels (untreated cells), # $P < 0.05$ compared to corresponding L-DOPA treatment (20, 100, and 200 μ M) (ANOVA followed by Dunnett's test).

which leads to decrease in dopamine levels because of L-DOPA-induced oxidative cytotoxicity.^{10,24} Treatment with L-DOPA alone (20, 100, and 200 μ M) increased intracellular dopamine levels (Fig. 4).¹⁰ Ombuoside (1, 5, and 10 μ M) co-administered with L-DOPA (20, 100, and 200 μ M) for 24 h further increased the intracellular levels of dopamine in PC12 cells compared to L-DOPA alone (Fig. 4.), indicating that ombuoside enhanced L-DOPA-induced dopamine biosynthesis in PC12 cells.

In general, flavonoids exhibit anti-oxidative functions.¹⁶ The anti-oxidative functions of ombuoside may serve as the mechanism of dopamine biosynthesis in PC12 cells. GP-EX and GPS have been shown to exhibit protective effects against neurotoxicity by reducing TH neuronal cell death and L-DOPA-induced cytotoxicity in rat models of PD.^{13,14} The functional relationships between GP-EX, GPS, and ombuoside need to be further elucidated for pharmaceutical application.

In conclusion, ombuoside enhanced dopamine biosynthesis by regulating the CREB-TH system in PC12 cells. Further *in vivo* work should be conducted to elucidate its efficacy using animal models.

Acknowledgements

This research was financially supported by the National

Research Foundation of Korea (grant no. 2013R1A1A2058230, 2015-2016), Republic of Korea.

References

- (1) Fahn, S. *Ann. NY Acad. Sci.* **2003**, *991*, 1-14.
- (2) Nagatsu, T.; Levitt, M.; Udenfriend, S. *J. Biol. Chem.* **1964**, *239*, 2910-2917.
- (3) Kilbourne, E. J.; Nankova, B. B.; Lewis, E. J.; McMahon, A.; Osaka, H.; Sabban, D. B.; Sabban, E. L. *J. Biol. Chem.* **1992**, *267*, 7563-7569.
- (4) Kim, K. S.; Lee, M. K.; Carroll, J.; Joh, T. H. *J. Biol. Chem.* **1993**, *268*, 15689-15695.
- (5) Campbell, D. G.; Hardie, D. G.; Vulliet, P. R. *J. Biol. Chem.* **1986**, *261*, 10489-10492.
- (6) Haycock, J. W. *Neurochem. Res.* **1993**, *18*, 15-26.
- (7) Kim, K. S.; Park, D. H.; Wessel, T. C.; Song, B.; Wagner, J. A.; Joh, T. H. *Proc. Natl. Acad. Sci. USA* **1993**, *90*, 3471-3475.
- (8) Marsden, C. D.; Parkes, J. D. *Lancet* **1977**, *1*, 345-349.
- (9) Migheli, R.; Godani, C.; Sciola, L.; Delogu, M. R.; Serra, P. A.; Zangani, D.; De Natale, G.; Miele, E.; Desole, M. S. *J. Neurochem.* **1999**, *73*, 1155-1163.
- (10) Jin, C. M.; Yang, Y. J.; Huang, H. S.; Lim, S. C.; Kai, M.; Lee, M. K. *Eur. J. Pharmacol.* **2008**, *591*, 88-95.
- (11) Razmovski-Naumovski, V.; Huang, T. H. W.; Tran, V. H.; Li, G. Q.; Duke, C. C.; Roufogalis, B. D. *Phytochem. Rev.* **2005**, *14*, 197-219.
- (12) Choi, H. S.; Zhao, T. T.; Shin, K. S.; Kim, S. H.; Hwang, B. Y.; Lee, C. K.; Lee, M. K. *Molecules* **2013**, *18*, 4342-4356.
- (13) Choi, H. S.; Park, M. S.; Kim, S. H.; Hwang, B. Y.; Lee, C. K.; Lee, M. K. *Molecules* **2010**, *15*, 2814-2824.
- (14) Shin, K. S.; Zhao, T. T.; Park, K. H.; Park, H. J.; Hwang, B. Y.; Lee, C. K.; Lee, M. K. *BMC Neurosci.* **2015**, *16*, 23.
- (15) Shin, K. S.; Zhao, T. T.; Choi, H. S.; Hwang, B. Y.; Lee, C. K.; Lee, M. K. *Brain Res.* **2014**, *1567*, 57-65.
- (16) Rice-Evans, C. A.; Miller, N. J.; Paganga, G. *Free Radic. Biol. Med.* **1996**, *20*, 933-956.
- (17) Amaro-Luis, J. M.; Adrián, M.; Díaz, C. *Ann. Pharm. Fr.* **1997**, *55*, 262-268.
- (18) Pollard, S. E.; Kuhnle, G. G.; Vauzour, D.; Vafeiadou, K.; Tzounis, X.; Whiteman, M.; Rice-Evans, C.; Spencer, J. P. *Biochem. Biophys. Res. Commun.* **2006**, *350*, 960-968.
- (19) Kumar, S.; Pandey, A. K. *Scientific World Journal* **2013**, *2013*, 162750.
- (20) Tischler, A. S.; Perlman, R. L.; Morse, G. M.; Sheard, B. E. *J. Neurochem.* **1983**, *40*, 364-370.
- (21) Mosmann, T. *J. Immunol. Methods* **1983**, *65*, 55-63.
- (22) Lowry, O. H.; Rosebrough, N. J.; Farr, A. L.; Randall, R. L. *J. Biol. Chem.* **1951**, *193*, 265-275.
- (23) Gonzalez, G. A.; Montminy, M. R. *Cell* **1989**, *59*, 675-680.
- (24) Basma, A. N.; Morris, E. J.; Nicklas, W. J.; Geller, H. M. *J. Neurochem.* **1995**, *64*, 825-832.

Received September 1, 2017

Revised February 8, 2018

Accepted February 11, 2018