



Integracide K: A New Tetracyclic Triterpenoid from *Desmodium uncinatum* (Jacq.) DC. (Fabaceae)

Borice Tapondjou Tsafack¹, Beaudelaire Kemvoufo Ponou¹, Rémy Bertrand Teponno¹, Raymond Ngansop Nono¹, Kristina Jenett-Siemss², Matthias F. Melzig², Hee Juhn Park^{3,*}, and Léon Azefack Tapondjou^{1,*}

¹Department of Chemistry, Faculty of Science, University of Dschang, Box 67, Dschang, Cameroon

²Institut für Pharmazie (Pharmazeutische Biologie), Freie Universität Berlin, Königin-Luise-Str. 2-4, D-14195 Berlin, Germany

³Department of Pharmaceutical Engineering, Sangji University, 26339 Wonju, Korea

Abstract – A new tetracyclic triterpenoid [4,4,24-trimethylcholesta- $\Delta^{8,9;14,15;24,28}$ -trien-3 β ,11 β ,12 α -triol-12-acetate, 3-sulfate] sodium salt (**1**), together with eight known compounds including ergosterol 5 α ,8 α -endoperoxide (**2**), 1,9-dihydroxy-3-methoxy-2-methylpterocarpan (**3**), 3-O- β -D-2-acetyl-amino-2-deoxyglucopyranosyloleanoic acid (**4**), hydnocarpin (**5**), derrone (**6**), isovitexin (**7**), erythrinin C (**8**), and 5,4'-dihydroxy-2"-hydroxyisopropylidihydrofuran [4,5:7,8]-isoflavone (**9**), were isolated from the EtOAc soluble fraction of the methanol extract of aerial part of *Desmodium uncinatum* collected in the western highland of Cameroon. The structures of these compounds were established by comprehensive interpretation of their spectral data mainly including 1D- (¹H and ¹³C), 2D-NMR (¹H-¹H COSY, HMQC, HMBC) spectroscopic and ESI-TOF-MS mass spectrometric analysis. The isolation of an integracide-like compound from plant origin is a very unusual finding.

Keywords – *Desmodium uncinatum*, Fabaceae, Tetracyclic triterpenoid, Integracide, Isoflavonoids

Introduction

The genus *Desmodium* is a large group of the Fabaceae family. It contains about 350 species which are mainly distributed in tropical and subtropical region of the world and about 28 species in China.¹ Most of its plants are herbs, shrubs or sub-shrubs but rarely trees. Besides their popularity as feeding stuffs, they are also used in traditional medicine.² Many species in this genus have a long history of medicinal uses and are recorded to have hepatoprotective effect.³ *Desmodium uncinatum* is a large perennial legume with stems that may grow several meters long and trail over surrounding vegetation. These cylindrical or angular stems are covered with short, hooked hairs that stick to hair or clothing. The roots of *D. uncinatum* have been found to stimulate germination of striga seeds and to inhibit radicle growth of the resulting seedlings; this combination represents the allelopathic mechanism associated with striga control.^{4,5} Previous

phytochemical investigations on these roots growing in Kenya have revealed the presence of isoflavanones including uncinanone A, B, C, D and E, and a pterocarpan uncinacarpan.^{4,6} The present study was undertaken in order to characterize the chemical constituents of the aerial part of this plant growing in the western region of Cameroon and this report deals with the isolation and structural elucidation of nine compounds from the ethyl acetate soluble fraction of its methanol extract, amongst which one new tetracyclic triterpenoid derivative (integracide K) (**1**) and eight known compounds (**2–9**).

Experimental

General procedure – The optical rotation was determined in MeOH with a MCP200 modular circular polarimeter. IR spectra were obtained on a Shimadzu FTIR-8400S spectrometer (Japan). ¹H NMR, ¹³C NMR, COSY, HMQC and HMBC spectra were performed in deuterated solvents on a Bruker DRX-500 Spectrometer at 500 MHz/125 MHz and on a Bruker AVANCE 700 spectrometer (Bruker, Germany) at 700 MHz/175 MHz. All chemical shifts (δ) are given in ppm units with reference to tetramethylsilane (TMS) as internal standard and the coupling constants (J) are in Hz. ESI mass spectra

*Author for correspondence

Leon Azefack Tapondjou, Department of Chemistry, Faculty of Science, University of Dschang, Box 67, Dschang, Cameroon
Tel: +237-675004826; E-mail: tapondjou2001@yahoo.fr

Hee-Juhn Park, Department of Pharmaceutical Engineering, Sangji University, 26339 Wonju, Korea
Tel: +82-33-730-0564; E-mail: hjpark@sangji.ac.kr

were carried out on an Agilent 6210 ESI-TOF mass spectrometer (Agilent Technologies, Santa Clara, CA, USA). Column chromatography was performed using 70 - 230 mesh and 230 - 400 mesh silica gel 60 (Merck), and sephadex LH-20. TLC was carried out on precoated silica gel 60 F₂₅₄ (Merck) plates and spots were visualized by a UV lamp multiband UV-254/365 nm (Model UVGL-58 Upland CA 91786, U.S.A) and by spraying with 50% H₂SO₄ and heating for 10 min at 110 °C.

Plant material – The aerial part of *D. uncinatum* was collected in the city of Dschang (Menoua Division, Western region of Cameroon) in August 2015. The plant identification was confirmed at the Cameroon National Herbarium, Yaoundé, where a voucher specimen was kept under the reference number HNC/5315.

Extraction and isolation – The powdered aerial part of *D. uncinatum* (4 kg) was extracted three times (each time for 24 h) with MeOH at room temperature followed by heating for 30 min before filtration. The combined filtrate was concentrated under reduced pressure to give the crude MeOH extract (168 g). Part of this extract (162 g) was suspended in distilled water (400 ml) and partitioned successively with ethyl acetate and *n*-butanol to yield after evaporation of solvents 78 g and 13 g of EtOAc and *n*-BuOH fractions, respectively. Part of the ethyl acetate fraction (73 g) was fractionated by a silica gel column chromatography using a gradient of ethyl acetate in *n*-hexane and thereafter a gradient of MeOH in ethyl acetate to give seven main fractions (A-G). Fraction E (3.5 g) was submitted to silica gel column chromatography eluting with hexane-EtOAc (1:1) to yield compound **5** (31 mg) and a mixture (sub-fraction E₁). Recrystallization of the sub-fraction E₁ in MeOH yielded compound **7** (26 mg). Fraction B (7 g) was purified over silica gel column chromatography eluted with an isocratic mixture of hexane-EtOAc (9:1) to afford compound **2** (15.6 mg) and a mixture (sub-fraction B₁). Sub-fraction B₁ was chromatographed on a silica gel column using the mixture hexane-EtOAc (8.5:1.5) as an eluent to give compounds **3** (7.8 mg) and **6** (6 mg). Recrystallization of Fraction C (11 g) (eluted with hexane-EtOAc 40%) in EtOAc yielded compound **9** (9.5 mg) and the resulting filtrate was chromatographed on the silica gel column using the mixture hexane-EtOAc (6:4) as an eluent to afford compound **8** (46 mg). Fraction G (12 g) (eluted with EtOAc-MeOH (8:2)) was chromatographed on a silica gel column with EtOAc-MeOH (9.5:0.5) as an eluent to give seven sub-fractions (G1-G7). The sub-fraction G5 was subjected to silica gel column chromatography eluted with EtOAc-MeOH (9.5:0.5) to give compound **4** (12 mg) and a mixture

Table 1. ¹H-NMR (400 MHz) and ¹³C-NMR (100 MHz) data of compound 1 (*J* in Hz)

Position	δ_{H}	δ_{C}
1	1.54 (2H, m)	34.2
2	2.18 (1H, m), 1.51 (1H, m)	33.3
3	3.62 (1H, d, <i>J</i> = 9.7; 4.2)	82.7
4		38.8
5	1.55 (1H, m)	50.9
6	1.69 (1H, m) 1.60 (1H, m)	18.4
7	2.32 (1H, m), 2.20 (1H, m)	26.8
8		125.5
9		140.3
10		36.7
11	4.10 (1H, brs)	67.3
12	4.96 (1H, d, <i>J</i> = 1.5)	77.5
13		46.8
14		147.3
15	5.50 (1H, brt)	120.3
16	2.31 (1H, m), 2.00 (1H, m)	36.2
17	1.85 (1H, m)	48.6
18	1.03 (3H, s)	24.0
19	0.99 (3H, s)	17.1
20	1.55 (1H, m)	33.6
21	0.84 (3H, d, <i>J</i> = 6.6)	18.3
22	1.51 (2H, m)	33.6
23	2.01 (1H, m), 1.88 (1H, m)	31.2
24		156.4
25	2.18 (1H, m)	34.9
26	0.98 (3H, d, <i>J</i> = 3.1)	22.4
27	1.00 (3H, d, <i>J</i> = 3.1)	22.3
28	4.71 (1H, brs) 4.66 (1H, brs)	107.8
29	0.74 (3H, s)	16.8
30	0.92 (3H, s)	28.7
31		170.4
32	1.96 (3H, s)	21.2
11-OH	5.31 (1H, d)	

The data were measured in DMSO with reference to TMS;

(sub-fraction B₅₋₁). The above mixture was purified by silica gel column chromatography eluted with EtOAc-MeOH (9.5:0.5) and thereafter by a sephadex LH-20 column chromatography eluted with MeOH to give compound **1** (25 mg).

Integracide K – Amorphous yellow powder. $[\alpha]^{20}_{\text{D}} = -93.84$ (c 0.065, MeOH); IR ν_{max} (NaCl) = 3400, 1643, 1226 cm⁻¹; ¹H and ¹³C NMR: see Table 1; ESI-MS (positive mode): *m/z* 623.3024 [M+Na]⁺, *m/z* 1223.6158 [2M+Na]⁺, (calcd. for C₃₂H₄₉O₇SnNa₂, 623.2994)

Ergosterol 5 α ,8 α -endoperoxide (2) – white amorphous powder; ESIMS (positive mode): m/z 451.3263 [$M+Na$]⁺ ($C_{28}H_{44}O_3Na$); ¹H-NMR (500 MHz, CD₃OD) : δ 0.87 (3H, s, H-18), 0.88 (6H, d, $J=6.5$ Hz, H-26/H-27), 0.92 (3H, s, H-19), 0.94 (3H, d, $J=6.8$ Hz, H-28), 1.00 (2H, m, H-15/H-16), 1.04 (3H, d, $J=6.6$ Hz, H-21), 1.28 (1H, m, H-17), 1.38 (2H, m, H-11), 1.44 (2H, m, H-15/H-16), 1.46 (2H, m, H-1), 1.49 (1H, m, H-25), 1.54 (1H, m, H-2), 1.79 (1H, m, H-2), 1.80 (1H, m, H-14), 1.88 (1H, m, H-23), 1.95 (2H, m, H-4), 1.98 (2H, m, H-12), 2.08 (1H, m, H-20), 3.75 (1H, m, H-3), 5.20 (1H, d, $J=7.9$ Hz, H-22), 5.25 (1H, d, $J=7.9$ Hz, H-23), 6.28 (1H, d, $J=8.5$ Hz, H-6), 6.56 (1H, d, $J=8.5$ Hz, H-7); ¹³C-NMR (125 MHz, CD₃OD): δ 12.1 (C-18), 16.6 (C-28), 18.1 (C-19), 19.5 (C-21), 19.9 (C-26), 20.2 (C-27), 21.1 (C-11), 23.6 (C-15), 29.0 (C-2), 33.3 (C-25), 34.6 (C-4), 35.1 (C-1), 37.4 (C-10), 39.0 (C-16), 39.5 (C-12), 40.1 (C-20), 43.0 (C-24), 44.7 (C-13), 51.8 (C-9), 52.0 (C-14), 56.3 (C-17), 65.5 (C-3), 79.3 (C-8), 82.0 (C-5), 131.0 (C-7), 132.3 (C-23), 136.0 (C-22), 136.1 (C-6).

1,9-Dihydroxy-3-methoxy-2-methylpterocarpan (3) – yellow amorphous powder; ESIMS (positive mode): m/z , 323.0920 [$M+Na$]⁺ ($C_{17}H_{16}O_5Na$); ¹H-NMR (500 MHz, CD₃OD): δ 2.10 (3H, s, 2-CH₃), 3.40 (1H, m, H-6a), 3.47 (1H, m, H-6), 3.78 (3H, s, 3-OCH₃), 4.22 (1H, m, H-6), 5.64 (1H, d, $J=6.3$ Hz, H-11a), 6.12 (1H, s, H-4), 6.31 (1H, d, $J=2.2$ Hz, H-10) 6.34 (1H, dd, $J=8.0$ and 2.2 Hz, H-8), 7.10 (1H, d, $J=8.0$ Hz, H-7); ¹³C-NMR (125 MHz, CD₃OD): δ 8.0 (CH₃-2), 38.2 (C-6a), 54.2 (3-OCH₃), 66.3 (C-6), 77.1 (C-11a), 91.2 (C-4), 98.6 (C-10), 101.3 (C-11b), 105.3 (C-2), 108.7 (C-8), 118.5 (C-6b), 125.4 (C-7), 155.2 (C-4a), 156.1 (C-1), 158.8 (C-9), 159.8 (C-3).

3-O- β -D-2-acetyl-amino-2-deoxyglucopyranoxyleanoic acid (4) – white amorphous powder, ESIMS (positive mode): m/z 682.4247 [$M+Na$]⁺ ($C_{38}H_{61}NO_8Na$); ¹H-NMR (500 MHz, CD₃OD) : δ 0.66 (3H, s, H-23), 0.72 (1H, m, H-5), 0.86 (3H, s, H-26), 0.88 (6H, s, H-29/H30), 0.90 (6H, s, H-26/H-24), 1.06 (1H, m, H-19), 1.10 (3H, s, H-27), 1.30 (2H, m, H-16), 1.46 (2H, m, H-1), 1.48 (1H, m, H-2), 1.49 (1H, m, H-11), 1.57 (2H, m, H-6), 1.57 (1H, m, H-19), 1.60 (2H, m, H-7/H-22), 1.75 (1H, m, H-7), 1.76 (3H, s, CH₃CO), 1.78 (1H, m, H-22), 1.80 (1H, m, H-11), 1.82 (1H, m, H-2), 1.90 (1H, m, H-9), 2.30 (2H, m, H-17), 2.76 (1H, d, $J=10.0$ and 2.8 Hz, H-18), 2.98 (1H, dd, $J=11.5$ and 4.1 Hz, H-3), 3.04 (2H, m, H-4'/H-5'), 3.27 (1H, m, H-3'), 3.40 (1H, m, H-2'), 3.41 (1H, m, H-6'), 3.67 (1H, m, H-6'), 4.27 (1H, d, $J=8.3$ Hz, H-1'), 5.18 (1H, brt, H-5), 7.70 (1H, d, $J=9.2$ Hz, H-NAc); ¹³C-NMR (125 MHz, CD₃OD): δ 15.5 (C-25), 16.7 (C-24),

17.6 (C-26), 18.3 (C-6), 22.6 (C-30), 23.0 (C-11), 23.2 (CH₃CO), 23.4 (C-16), 26.2 (C-2), 26.3 (C-27), 27.6 (C-23), 28.1 (C-15), 30.8 (C-20), 32.7 (C-29), 32.9 (C-22), 33.3 (C-7), 36.7 (C-10), 33.8 (C-21), 38.4 (C-4), 38.8 (C-1), 39.4 (C-8), 41.7 (C-18), 41.8 (C-14), 45.7 (C-19), 45.8 (C-17), 47.5 (C-9), 55.4 (C-5), 56.4 (C-2'), 61.6 (C-6'), 71.5 (C-4'), 74.3 (C-3'), 77.2 (C-5'), 88.4 (C-3), 104.0 (C-1'), 122.0 (C-12), 144.4 (C-13), 169.2 (CH₃CO), 179.3 (C-28).

Hydnocarpin (5) – yellow amorphous powder, ESIMS (positive mode): m/z 465.1244 [$M+Na$]⁺ ($C_{25}H_{20}O_9Na$); ¹H-NMR (500 MHz, DMSO): δ 3.40 (1H, m, H_a of 3-CH₂OH), 3.59 (1H, m, H_b of 3-CH₂OH), 3.79 (3H, s, 3"-OCH₃), 4.28 (1H, m, H-2), 5.03 (1H, m, H-3), 6.20 (1H, d, $J=2.0$ Hz, H-8), 6.52 (1H, d, $J=2.0$ Hz, H-6'), 6.82 (1H, d, $J=8.1$ Hz, H-8), 6.88 (1H, dd, $J=8.1$ and 1.8 Hz, H-7), 6.90 (1H, s, H-3'), 7.05 (1H, d, $J=1.8$ Hz, H-3), 7.10 (1H, d, $J=8.5$ Hz, H-5"), 7.62 (1H, dd, $J=8.5$ and 2.2 Hz, H-6"), 7.68 (1H, d, $J=2.2$ Hz, H-2"); ¹³C-NMR (125 MHz, DMSO): δ 55.1 (3"-OCH₃), 60.1 (CH₂OH), 71.1 (C-2), 77.1 (C-3), 93.5 (C-8'), 98.5 (C-6), 103.6 (C-3'), 103.7 (C-10'), 111.4 (C-5), 114.6 (C-2"), 115.3 (C-8), 117.2 (C-6"), 119.5 (C-5"), 120.6 (C-7), 123.8 (C-6), 124.6 (C-1"), 144.0 (C-10), 147.1 (C-9), 148.0 (C-4"), 148.3 (C-3"), 157.6 (C-9'), 162.1 (C-5'), 163.2 (C-7'), 163.3 (C-2'), 182.1 (C-4')

Derrone (6) – yellow amorphous powder, ESIMS (positive mode): m/z 337.1101 [$M+H$]⁺ ($C_{20}H_{18}O_6$), [$M+Na$]⁺ 359.0920; ¹H-NMR (500 MHz, CD₃OD): δ 1.30 (6H, s, H-5"/H-6"), 5.72 (1H, d, $J=10.0$ Hz, H-3"), 6.25 (1H, s, H-6), 6.75 (1H, d, $J=10.0$ Hz, H-4"), 6.80 (2H, d, $J=8.7$ Hz, H-3'/H-5'), 7.40 (2H, d, $J=8.7$ Hz, H-2'/H-6'); ¹³C-NMR (125 MHz, CD₃OD): δ 26.7 (C-5"/C-6"), 77.8 (C-2"), 99.3 (C-6), 101.4 (C-8), 113.0 (C-4"), 114.8 (C-3'/C-5'), 121.5 (C-1'), 124.6 (C-3), 127.3 (C-3"), 130.1 (C-2'/C-6'), 152.6 (C-2), 153.4 (C-8a), 156.1 (C-4'), 159.8 (C-7), 181.0 (C-4).

Isovitexin (7) – yellow amorphous powder; ¹H-NMR (500 MHz, CD₃OD): δ 3.37 (1H, m, H-4"), 3.49 (1H, m, H-3"), 3.78 (1H, m, H-5"), 3.91 (2H, m, H-6"), 4.19 (1H, m, H-2"), 4.92 (1H, d, $J=9.2$ Hz, H-1"), 6.48 (1H, s, H-8), 6.57 (1H, s, H-3), 6.92 (2H, d, $J=8.7$ Hz, H-3'/H-5'), 7.81 (2H, d, $J=8.7$ Hz, H-2'/H-6'); ¹³C-NMR (125 MHz, CD₃OD): δ 61.4 (C-6"), 70.4 (C-4"), 71.1 (C-2"), 73.9 (C-1"), 78.7 (C-3"), 81.20 (C-5"), 93.9 (C-8), 102.3 (C-3), 103.7 (C-10), 107.7 (C-6), 115.6 (C-3'/C-5'), 121.6 (C-1'), 128.0 (C-2'/C-6'), 157.2 (C-9), 160.5 (C-5), 161.3 (C-4'), 163.4 (C-7), 164.7 (C-2), 182.5 (C-4).

Erythrinin C (8) – yellow amorphous powder; ESIMS (positive mode): m/z 355.1221 [$M+H$]⁺ ($C_{20}H_{18}O_6$), 377.1041

$[M+Na]^+$; $^1\text{H-NMR}$ (500 MHz, CD_3OD): δ 1.23 (3H, s, H-5"), 1.29 (3H, s, H-4"), 3.14 (2H, dd, J = 2.5 and 8.7 Hz, H-1"), 4.78 (1H, t, J = 8.7 Hz), 6.40 (1H, s, H-8), 6.84 (2H, d, J = 9.1, H-3'/H-5'), 7.37 (2H, d, J = 9.1, H-2'/H-6'), 8.07 (1H, s); $^{13}\text{C-NMR}$ (125 MHz, CD_3OD): δ 24.0 (C-5"/C-4"), 26.0 (C-1"), 71.0 (C-3"), 88.1 (C-8), 91.7 (C-2"), 105.8 (C-4a), 109.2 (C-6), 114.9 (C-3'/C-5'), 121.7 (C-1'), 123.2 (C-3), 130.3 (C-2'/C-6'), 153.5 (C-2), 157.6 (C-5), 157.5 (C-4'), 158.7 (C-8a), 166.4 (C-7).

5,4'-Dihydroxy-2"-hydroxyisopropyldihydrofuranone [4,5:7,8]-isoflavone (9)—yellow amorphous powder, ESIMS (positive mode): m/z 355 $[M+\text{H}]^+$ ($\text{C}_{20}\text{H}_{19}\text{O}_6$), $[M+\text{Na}]^+$ 377; $^1\text{H-NMR}$ (500 MHz, DMSO): δ 1.15 (3H, s, H-5"), 1.17 (3H, s, H-4"), 3.20 (2H, dd, J = 8.5 and 5.3 Hz, H-1"), 4.80 (1H, t, J = 8.6 Hz, H-2"), 6.31 (1H, s, H-6), 6.83 (2H, d, J = 8.5 Hz, H-3'/H-5'), 7.37 (2H, d, J = 8.5 Hz, H-2'/H-6'), 8.35 (1H, s, H-2), 9.64 (1H, s, OH-9) 13.20 (1H, s, OH-5); $^{13}\text{C-NMR}$ (125 MHz, DMSO): δ 25.2 (C-5"), 26.1 (C-4"), 26.5 (C-1"), 70.6 (C-3"), 91.7 (C-2"), 94.0 (C-6), 104.0 (C-8), 105.3 (C-4a), 115.5 (C-3'/C-5'), 121.6 (C-1'), 123.0 (C-3), 130.7 (C-2'/C-6'), 152.5 (C-8a), 154.8 (C-2'), 157.9 (C-4'), 162.8 (C-5), 167.8 (C-7).

Results and Discussion

The structures of compounds **2** - **7** (Fig. 1) were deter-

mined by comparing their NMR (^1H , ^{13}C) and MS spectral data with those found of the literatures of ergosterol 5 α ,8 α -endoperoxide (**2**),⁷ 1,9-dihydroxy-3-methoxy-2-methylpterocarpan (**3**),⁵ 3-O- β -D-2-acetyl-amino-2-deoxy-glucopyranoxyloleanoic acid (**4**),⁸ hydnocarpin (**5**),⁹ derrone (**6**),¹⁰ isovitexin (**7**),¹¹ erythrinin C (**8**),¹² and 5,4'-dihydroxy-2"-hydroxyisopropyldihydrofuranone[4,5:7,8]-isoflavone (**9**).¹³

Compound **1** obtained as yellow powder was positively reacted with Liebermann-Burchard reagent, indicating that it is a triterpenoid. Its positive ESIMS displayed two important pseudo-molecular ion peaks at m/z 623.3024 $[M+\text{Na}]^+$ ($[\text{C}_{32}\text{H}_{49}\text{O}_7\text{SNa}+\text{Na}]^+$) and m/z 1223.6158 $[2\text{M}+\text{Na}]^+$ ($[\text{2C}_{32}\text{H}_{49}\text{O}_7\text{SNa}+\text{Na}]^+$), thus indicating a molecular weight of 600 a.m.u and the presence of a sulfate group (SO_4Na). The presence of this function was confirmed by the characteristic vibration band in the IR spectrum at 1226 cm^{-1} .¹⁴ When the ^1H NMR spectra of **1** were both recorded in $\text{DMSO}-d_6$ and in pyridine- d_5 , the spectra obtained in $\text{DMSO}-d_6$ was quite indicative as it revealed the signal of only one free OH group. This fact was quite determinant for the elucidation of the structure. The ^1H , ^{13}C and HMQC NMR spectra of **1** showed eight methyl groups at δ_{H} 0.98 (CH_3 -26)/ δ_{C} 22.4 (d, J = 3.1), δ_{H} 1.00 (CH_3 -27)/ δ_{C} 22.3 (d, J = 3.1); δ_{H} 0.84 (CH_3 -21)/ δ_{C} 18.3; δ_{H} 0.99 (CH_3 -19)/ δ_{C} 17.1; δ_{H} 1.03 (CH_3 -18)/ δ_{C} 24.1; δ_{H}




Fig. 1. Chemical structures of compounds isolated from *Desmodium uncinatum*.

0.74 ($\text{CH}_3\text{-}29$)/ δ_{C} 16.9; δ_{H} 0.92 ($\text{CH}_3\text{-}30$)/ δ_{C} 28.7, seven methylenes, seven methine groups (three of which are oxygenated), and eight quaternary carbons including one carbonyl and six olefinic carbons (three of them are tri-substituted olefinic double bonds conjugated at δ_{C} 140.3 (C-9); δ_{C} 125.5 (C-8) and δ_{C} 147.3 (C-14)). These NMR data were quite similar to those of integracide-like compounds;^{15,16,17} therefore the dienic system was positioned at C-14–C-15 and C-8–C-9 based on the literature. This was thereafter confirmed by HMBC correlations of $\text{CH}_3\text{-}19$ (δ_{H} 0.99) to C-14, C-12 (δ_{C} 77.5), C-17 (δ_{C} 49.1) and C-13 (δ_{C} 46.8); H-12 (δ_{H} 4.96) to C-14 and C-9; OH-11 (δ_{H} 5.31) to C-9; $\text{CH}_3\text{-}18$ (δ_{H} 1.03) to C-9, C-1 (δ_{C} 34.2) and C-10 (δ_{C} 36.7). The position of this dienic system was also justified by the ^1H - ^1H COSY cross peaks of H-15 (δ_{H} 5.50) to the methylene protons at δ_{H} 2.31 (m H_a-16) and 2.00 (m, H_b-16). Furthermore, the presence of signals at δ_{H} 4.71 (brs, H_a-28) and δ_{H} 4.66 (brs, H_b-28)/ δ_{C} 107.8 (C-28) and δ_{C} 156.4 (C-24) indicate the presence of an exomethylene group. The position of this group was confirmed by HMBC cross peaks of $\text{CH}_3\text{-}26$ (δ_{H} 0.98) and $\text{CH}_3\text{-}27$ (δ_{H} 1.00) to C-24 (Fig. 2). The HMBC spectrum also showed cross peaks from $\text{CH}_3\text{-}29$ (δ_{H} 0.92) and $\text{CH}_3\text{-}30$ (δ_{H} 0.74) to C-3 (δ_{C} 82.7), C-4 (δ_{C} 38.8) and C-5 (δ_{C} 50.9) establishing the location of the gem dimethyl group at C-4 (δ_{C} 38.8). Other important cross peaks such as $\text{CH}_3\text{-}21$ (δ_{H} 0.84) to C-20 (δ_{C} 33.6), C-17 (δ_{C} 49.1) and C-22 (δ_{C} 33.6) indicated the presence of a methyl group at C-20. The presence of an isopropyl group in **1** was characterized by the signals at δ_{H} 0.98 (d, J =3.1 Hz, $\text{CH}_3\text{-}26$)/ δ_{C} 22.4, δ_{H} 1.00 (d, J =3.1 Hz, $\text{CH}_3\text{-}27$)/ δ_{C} 22.3 and δ_{H} 2.18 (m, H-25)/ δ_{C} 34.9 and confirmed by the ^1H - ^1H COSY correlation of H-26 and H-27 with H-25. The connectivity of the isopropyl fragment at C-24 was confirmed by the HMBC correlation of $\text{CH}_3\text{-}26$, $\text{CH}_3\text{-}27$ to C-25 (δ_{C} 34.9) and C-24 (δ_{C} 156.4). The signal at δ_{H} 4.96 (d, J =1.5 Hz, H-12)/ δ_{C} 77.5 (C-12); δ_{H} 4.10 (brs H-11)/ δ_{C} 67.3 (C-11) and δ_{H} 3.62 (dd, J =9.7, 4.3 Hz, H-3)/ δ_{C} 82.7 (C-3) indicated the presence of three oxygenated methine groups, they were located at C-12, C-11 and C-3, respectively, based on the observed ^1H - ^1H COSY correlations of H-11 to H-12 and OH-11 (δ_{H} 5.31) and HMBC correlation of $\text{CH}_3\text{-}29$ and $\text{CH}_3\text{-}30$ to C-3. The ^1H NMR spectrum of **1** showed one singlet methyl signal at δ_{H} 1.96 ($\text{CH}_3\text{-}32$) correlated to the carbon signal resonating at δ_{C} 21.2 (C-32) in the HMQC spectrum. In the HMBC spectrum, this methyl group shows a cross peak with the carbonyl carbon at δ_{C} 170.4 (C-31), indicating the presence of one acetoxy group in **1**. Moreover, the presence of only one singlet signal at δ_{H} 5.31 in the ^1H NMR spectrum and




Fig. 2. Selected HMBC correlations of Integracide K (**1**).

assigned to OH-11 group indicated the substitution of other two oxymethines located at C-3 and C-12. As such, the acetoxy group was located at C-12 based on the literature comparison of the NMR data of **1** with those of integracide A-J,^{15,16,17} thus the sulphate group was located at C-3. The stereochemistry of the different chiral centers were established based on the comparison of their spectral data with those of reported compounds. More precisely, the stereochemistry at C-11, C-12 and C-17 were confirmed based on the coupling constants observed for their corresponding proton atoms (Table 1) by comparing with those of reported integracides.^{15,16,17} Thus the structure of compound **1** was elucidated as 4,4,24-trimethylcholesta- $\Delta^{8,9;14,15;24,28}$ -trien-3 β ,11 β ,12 α -triol-12-acetate,3-sulfate sodium salt.

Integracides are oxygenated tetracyclic 4,4-dimethyler-gostane terpenoids possessing a 12-acetyl- $\Delta^{8,14}$ dien-11-ol moiety. Up to date, they have been reported only from fungi of *Fusarium* species such as *Fusarium sporotrichioides* 921,¹⁸ *Fusarium compactum*¹³ and *Fusarium sp*^{14,15,16} and therefore seem to be the chemotaxonomic markers of these species. The present work is then the first report of an integracide-like compound from plants origin and so far from *Desmodium* species. Integracides have been found to exhibit important inhibitory activity against the HIV-1 integrase^{15,19} and to display potent cytotoxic and anti-leishmanial activities.^{16,17} Therefore, various bioactivities of *D. uncinatum* used in Cameroonian folk medicine might not only be attributed to its isoflavonoid but also to integracide K.

Acknowledgements

The authors are grateful to the Alexander von Humboldt Foundation (AvH), Bonn, Germany for the financial support of this work.

References

- (1) Liguo, F.; Tanqing, C.; Kaiyong, L.; Hong, T.; Lin, Qi. Higher plants of China; Qingdao Publishing House: Qingdao, **2008**, p 152.
- (2) Ma, X.; Zheng, C.; Hu, C.; Rahman, K.; Qin, L. *J. Ethnopharmacol.* **2011**, *138*, 314-332.
- (3) Vedpal, Dhanabal, S. P.; Dhamodaran, P.; Chaitnya, M. V. N. L.; Duraiswamy, B.; Jayaram, U.; Srivastava, N. *J. Chem. Pharm. Res.* **2016**, *8*, 91-97.
- (4) Tsanuo, M. K.; Hassanali, A.; Hooper, A. M.; Khan, Z.; Kaberia, F.; Pickett, J. A.; Wadhams, L. *J. Phytochemistry* **2003**, *64*, 265-273.
- (5) Khan, Z. R.; Hassanali, A.; Overholt, W.; Khamis, T. M.; Hooper, A. M.; Pickett, J. A.; Wadhams, L. J.; Woodcock, C. M. *J. Chem. Ecol.* **2002**, *28*, 1871-1885.
- (6) Guchu, S. M.; Yenesew, A.; Tsanuo, M. K.; Gikonyo, N. K.; Pickett, J. A.; Hooper, A. M.; Hassanali, A. *Phytochemistry* **2007**, *68*, 646-651.
- (7) Correa, E.; Quñones, W.; Torres, F.; Cardona, D.; Franco, A. E.; Robledo, S.; Echeverri, F. *Actual. Biol.* **2005**, *27*, 39-42.
- (8) Wang, P.; Wang, J.; Guo, T.; Li, Y. *Carbohydr. Res.* **2010**, *345*, 607-620.
- (9) Afifi, M. S. A.; Ahmed, M. M.; Pezzuto, J. M.; Kinghorn, A. D. *Phytochemistry* **1993**, *34*, 839-841.
- (10) Máximoa, P.; Lourenço, A.; Savluchinske Feio, S.; Roseiro, J. C. Z. *Naturforsch. C* **2002**, *57*, 609-613.
- (11) Peng, J.; Fan, G.; Hong, Z.; Chai, Y.; Wu, Y. *J. Chromatogr. A* **2005**, *1074*, 111-115.
- (12) Tahara, S.; Ingham, J. L.; Mizutani, J. *Agric. Biol. Chem.* **1985**, *49*, 1775-1783.
- (13) Dai, J.; Shen, D.; Yoshida, W.Y.; Parrish, S. M.; Williams, P. G *Planta Med.* **2012**, *78*, 1357-1362.
- (14) Brill, G. M.; Kati, W. M.; Montgomery, D.; Karwowski, J. P.; Humphrey, P. E.; Jackson, M.; Clement, J. J.; Kadam, S.; Chen, R. H.; McAlpine, J. B. *J. Antibiot.* **1996**, *49*, 541-546.
- (15) Singh, S. B.; Zink, D. L.; Dombrowski, A. W.; Polishook, J. D.; Ondeyka, J. G.; Hirshfield, J.; Felock, P.; Hazuda, D. *J. Bioorg. Med. Chem.* **2003**, *11*, 1577-1582.
- (16) Ibrahim, S. R. M.; Mohamed, G. A.; Ross, S. A. *Phytochem. Lett.* **2016**, *15*, 125-130.
- (17) Ibrahim, S. R. M.; Abdallah, H. M.; Mohamed, G. A.; Ross, S. A. *Fitoterapia* **2016**, *112*, 161-167.
- (18) Chatterjee, S.; Sarkar A.; Dutta, P. C. *J. Chem. Soc., Perkin Trans. 1* **1979**, *1*, 2914-2919.
- (19) Singh, S. B.; Ondeyka, J. G.; Schleif, W. A.; Felock, P.; Hazuda, D. *J. J. Nat. Prod.* **2003**, *66*, 1338-1344.

Received January 29, 2017

Revised February 28, 2017

Accepted March 7, 2017