

Original Research



OPEN ACCESS

Received: Jan 4, 2021
Revised: Jan 10, 2021
Accepted: Feb 3, 2021

*Corresponding Author:

Kyunghee Song

Department of Food and Nutrition, Myongji University, 116 Myongji-ro, Cheoin-gu, Yongin 17058, Korea.

Tel. +82-31-330-6206

Fax. +82-31-330-6200

E-mail. khsong@mju.ac.kr

©2021 The Korean Nutrition Society and the Korean Society of Community Nutrition
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Onjeong Choi

<https://orcid.org/0000-0001-5093-8476>

Jiwon Kim

<https://orcid.org/0000-0002-3307-1441>

Yujin Lee

<https://orcid.org/0000-0002-6948-2525>

Youngmi Lee

<https://orcid.org/0000-0001-9965-0748>

Kyunghee Song

<https://orcid.org/0000-0001-9549-0716>

Conflict of Interest

The authors declare no potential conflicts of interests.

Association between stress and dietary habits, emotional eating behavior and insomnia of middle-aged men and women in Seoul and Gyeonggi

Onjeong Choi , Jiwon Kim , Yujin Lee , Youngmi Lee , and Kyunghee Song *

Department of Food and Nutrition, Myongji University, Yongin 17058, Korea

ABSTRACT

BACKGROUND/OBJECTIVES: The study was performed to investigate the degree of perceived stress by sex in middle-aged people, and to provide basic data for appropriate nutrition education and interventional measures for middle-aged subjects through comparative analysis of association between stress and dietary habits, emotional eating behavior, and insomnia.

SUBJECTS/METHODS: A survey was conducted in 670 middle-aged subjects (320 men and 350 women) aged 40–64 years, resided in Seoul and Gyeonggi area. Collected data were analyzed using SPSS WIN 25.0 program.

RESULTS: Perceived stress in middle-aged subjects was not different by sex. Stress was higher in men with low dietary habits level compared to other groups ($P < 0.001$), and was higher in women with low and moderate dietary habits level compared to women with high dietary habits level ($P < 0.05$). Stress in men was higher in the group with emotional eaters ($P < 0.05$), and the same result was shown in women ($P < 0.001$). Stress was higher in both men and women with poor sleep quality ($P < 0.001$). Stress score was positively correlated with emotional dietary behavior and insomnia ($P < 0.001$) and negatively correlated with dietary habits ($P < 0.001$).

CONCLUSIONS: It is considered that proper nutrition education and interventional measures according to sex are needed for stress control and proper dietary behavior and lifestyle in middle-aged people population, along with projects and policies at the national level.

Keywords: Stress; dietary habit; emotional eating behavior; insomnia; middle-aged people

INTRODUCTION

While the life expectancy has been increased since 2005, the health-span tended to decrease compare to 2016 [1]. As of 2018 the life expectancy was 82.7 years but the healthy life year was 64.4 years with about 17 years of discrepancy. This suggests that it might be necessary to improve the quality of life in the elderly, for which mental and physical health management is needed in the middle-aged. The middle-aged is a period of being exposed to various stresses which are major factors for lowering the quality of life and health [2]. Middle-aged men experience psychological stress due to various role performance and physical changes. Stress causes physical and mental symptoms such as peripheral vascular disorders, gastrointestinal discomfort, and muscular tension in middle-aged women [3].

Author Contributions

Conceptualization: Song K; Formal analysis: Choi O, Kim J; Methodology: Choi O; Supervision: Song K; Writing - original draft: Choi O; Writing - review & editing: Lee Y¹, Lee Y².

Lee Y,¹ Lee Yujin; Lee Y,² Lee Youngmi.

Stress is associated with dietary behavior, and also affects dietary intake patterns and can induce overeating or meal-skipping. Overeating leads to obesity and meal-skipping can reduce body weight resulting in serious adverse effect on health [4]. Proper dietary habits are the basics of balanced nutritional intake and a major factor to determine the individual health status. Irregular dietary habits not only disrupt physical health but also can affect psychological condition and emotional stability [5].

Stress is known to be a major risk factor for sleep disorder [6]. Sleep disorder includes difficulties in falling asleep, waking up frequently or too early causing fatigue during daily activities [7]. Sleep disorder has been increased by 8.1% annually during the past 5 years, and the number of patients with sleep disorder has been increased by age according to 2018 Health Insurance Report [8]. Except for population over 65 years of age, sleep disorder was the highest in the middle-aged group. Also, sleep disorder has increased the incidence of cardiovascular disease, diabetes, hypertension, and metabolic syndrome in adults [9,10].

In middle-aged people, proper management of stress, dietary habits, emotional dietary behaviors and sleep can improve the quality of life and health. But it is thought that the improved quality of life and healthy lifestyle can be maintained more effectively when the relationship between stress and lifestyle is checked and managed together rather than separate management of stress and lifestyle. Thus, the study was performed to investigate and analyze dietary habits, emotional dietary behavior, and sleep according to sex and stress of middle-aged people and then to provide basic data for nutrition education and nutritional intervention of middle-aged subjects.

SUBJECTS AND METHODS

Subjects and data collection

A survey was conducted using online and self-administered methods in middle-aged subjects of 40–64 years of age in Seoul and Gyeonggi area from May 5 to May 22, 2020. Among a total of 724 questionnaire collected, 670 questionnaire including 417 online questionnaire (180 men, 237 women) and 253 written questionnaire (140 men, 113 women) were used in the study, after those with 54 missing responses were excluded. A survey response rate was 92.5%.

General characteristics

General characteristics of the subjects included age, body weight status, perceived health status, education level, occupation, and monthly household income. Anthropometric data of the subjects were obtained from self-administered questionnaire and the body mass index (BMI, kg/m²) was calculated using height and weight. BMI was divided into underweight (BMI < 18.5), normal weight (18.5 ≤ BMI < 23), overweight (23 ≤ BMI < 25), and obesity (BMI ≥ 25) according to the Asia-Pacific BMI classifications (World Health Organization 2000) [11].

Stress

The stress score used PSS (perceived stress scale) developed by Cohen *et al.* [12] and adapted by Lee *et al.* [13]. It was made of a total of 10 questions, and each question was evaluated by Likert 5-point scale. The range of total score was 0–40 points and the perceived stress level was severe as the score increased. The Cronbach's alpha was 0.79 in the study.

Dietary habits

A total of 10 questions on dietary habits included three meals a day, breakfast, meal regularity and amount, food combination, green-and-yellow vegetable intake, fruit intake, protein intake, milk and dairy product intake, seaweed intake, and water intake. Each question was evaluated by Likert 5-point scale. The higher score meant good dietary habits. In this study, the upper 30% was classified as high level, the next 40% as middle level, and the lower 30% as low level for analysis. The Cronbach's alpha was 0.83.

Emotional eating behavior

The level of emotional dietary behavior used emotional eater questionnaire (EEQ) developed by Garaulet *et al.* [14]. It was made of a total of 10 questions, and each question was evaluated by Likert 4-point scale. In the study, 0–5 points were classified as non-emotional eater group, 6–10 points as low emotional eater group, and over 11 points as emotional eater group for analysis. Before the study, the validity and reliability of the questionnaire were established. The Cronbach's alpha was 0.78 in the previous study [15] and 0.87 in this study.

Sleep quality

The insomnia level used insomnia severity index (ISI), which was developed by Bastien *et al.* [16] and adapted by Cho *et al.* [17] as Korean version of ISI. It was made of a total of 7 questions and each question was evaluated by Likert 5-point scale. In the study, 0–7 points were classified as no clinically significant insomnia group, 8–14 points as subthreshold insomnia group, and over 15 points as clinical insomnia group for analysis. In the previous study [17], the Cronbach's alpha was 0.92; in this study, it was 0.86.

Statistical analysis

Data were analyzed using SPSS Win 25.0. For general characteristics, the frequency and percentage were calculated, and continuous variables were analyzed using independent samples t-test. One-way analysis of variance test was performed to obtain stress score according to dietary habits, emotional dietary behavior, and insomnia level, and *post hoc* analysis was performed using Duncan's new multiple range test. Pearson's correlation coefficient test was performed to find the association among dietary habits, emotional dietary behavior, insomnia, and stress. All results were determined the statistical significance at $P < 0.05$.

Research ethics

For written questionnaire, the instruction was presented to the subjects and the consent form was obtained from the subjects who agreed to the instruction after sufficient explanation before the survey. For online questionnaire, the same instruction and agreement form as in the case of written questionnaire were presented and then the survey was proceeded only if the subject checked the 'I agree' box for agreement. It was clearly stated in the instruction and agreement form that the withdrawal of the agreement could be made after the participation in the study. Also, it was explained that data would not be used except for this study purpose, and related data would be stored in the Institutional Review Board (IRB) organization for 3 years after the termination of the study, and then all data would be discarded. The study was performed after the deliberation and approval of Myongji University IRB (deliberation No.: MJU-2020-04-002-01).

RESULTS

General characteristics of the subjects

General characteristics of the study subjects are shown in **Table 1**.

The highest age range was 50–54 years as 27.5% in men and 27.4% in women. For the weight status, ‘obese’ was the highest as 46.3% in men and ‘normal weight’ was the highest as 51.7% in women ($P < 0.001$). The BMI was 25.17 kg/m² in men and 22.96 kg/m² in women ($P < 0.001$). For the perceived health status, ‘average’ was the highest as 61.6% in men and 62.6% in women. The ratio of positive perceived health status was higher in men while the ratio of negative perceived health status was higher in women ($P < 0.01$). For education level, college graduate was 42.2% and over graduate school was 32.2% in men; college graduate was 50.6% and high school graduate was 31.4% in women ($P < 0.001$). For occupation, office

Table 1. General characteristics of subjects by sex

Classification	Male (n = 320)	Female (n = 350)	Total (n = 670)	t or χ^2	P-value
Age (yrs)				3.447	0.486
40–44	46 (14.4)	51 (14.6)	97 (14.5)		
45–49	87 (27.2)	79 (22.6)	166 (24.8)		
50–54	88 (27.5)	96 (27.4)	184 (27.5)		
55–59	64 (20.0)	72 (20.6)	136 (20.3)		
60–64	35 (10.9)	52 (14.9)	87 (13.0)		
Weight status ¹⁾				83.862***	< 0.001
Underweight	2 (0.6)	11 (3.1)	13 (1.9)		
Normal	66 (20.6)	181 (51.7)	247 (36.9)		
Overweight	104 (32.5)	81 (23.1)	185 (27.6)		
Obese	148 (46.3)	77 (22.0)	225 (33.6)		
BMI	25.17 ± 2.90	22.96 ± 2.89	24.01 ± 3.09	9.881***	< 0.001
Perceived health status				19.079**	0.001
Very bad	3 (0.9)	10 (2.9)	13 (1.9)		
Bad	15 (4.7)	41 (11.7)	56 (8.4)		
Average	197 (61.6)	219 (62.6)	416 (62.1)		
Good	93 (29.1)	71 (20.3)	164 (24.5)		
Very good	12 (3.8)	9 (2.6)	21 (3.1)		
Education level				26.623***	< 0.001
≤ Middle school	5 (1.6)	9 (2.6)	14 (2.1)		
High school	77 (24.1)	110 (31.4)	187 (27.9)		
College	135 (42.2)	177 (50.6)	312 (46.6)		
≥ Graduate school	103 (32.2)	54 (15.4)	157 (23.4)		
Occupation				141.708***	< 0.001
Production Worker/laborer	50 (15.6)	18 (5.2)	68 (10.1)		
Sales/service	24 (7.5)	40 (11.6)	64 (9.6)		
Professional	71 (22.1)	78 (22.2)	149 (22.2)		
Office staff	104 (32.5)	59 (16.7)	163 (24.3)		
Administration management	14 (4.4)	4 (1.2)	18 (2.7)		
Self employed	57 (17.9)	44 (12.5)	101 (15.1)		
Housekeeping	0	107 (30.7)	107 (16.0)		
Monthly household income (10,000 won)				9.937	0.127
< 200	21 (6.6)	39 (11.1)	60 (9.0)		
200–299	38 (11.9)	53 (15.1)	91 (13.6)		
300–399	51 (15.9)	52 (14.9)	103 (15.4)		
400–499	55 (17.2)	58 (16.6)	113 (16.9)		
500–599	43 (13.4)	54 (15.4)	97 (14.5)		
600–699	24 (7.5)	24 (6.9)	48 (7.2)		
≥ 700	88 (27.5)	70 (20.0)	158 (23.6)		

Data are shown as mean ± SD or number (%).

¹⁾Underweight: < 18.5, normal: 18.5–22.9, overweight: 23–24.9, obese: ≥ 25.

** $P < 0.01$, *** $P < 0.001$ by independent samples t-test or χ^2 test.

Table 2. Stress score of subjects by sex

Classification	Male (n = 320)	Female (n = 350)	Total (n = 670)	t	P-value
Been upset because of something that happened unexpectedly?	2.67 ± 0.66	2.70 ± 0.73	2.69 ± 0.70	-0.580	0.562
Felt that you were unable to control the important things in your life?	2.76 ± 0.78	2.83 ± 0.83	2.79 ± 0.81	-1.159	0.247
Felt nervous and "stressed?"	2.95 ± 0.82	3.01 ± 0.75	2.98 ± 0.78	-1.071	0.285
Felt confident about your ability to handle your personal problems?	2.70 ± 0.75	2.82 ± 0.79	2.76 ± 0.77	-1.912	0.056
Felt that things were going your way?	2.88 ± 0.73	2.91 ± 0.80	2.90 ± 0.77	-0.360	0.719
Fund that you could not cope with all the things that you had to do?	2.70 ± 0.77	2.66 ± 0.74	2.68 ± 0.76	0.688	0.492
Been able to control irritations in your life?	2.71 ± 0.77	2.65 ± 0.79	2.68 ± 0.78	0.907	0.365
Felt that you were on top of things?	2.97 ± 0.73	3.02 ± 0.82	2.99 ± 0.78	-0.807	0.420
Been angered because of things that were outside of your control?	2.80 ± 0.74	2.75 ± 0.74	2.77 ± 0.74	1.001	0.317
Felt difficulties were piling up so high that you could not overcome them?	2.61 ± 0.79	2.55 ± 0.82	2.57 ± 0.81	0.972	0.331
Total stress score	17.75 ± 4.31	17.89 ± 4.76	17.82 ± 4.55	-0.395	0.693

Data are shown as mean ± SD.

staff was the highest as 32.5% and then in the order of 22.1% of professionals and 17.9% of self-employed in men. On the other hand, in women, housekeeping was the highest as 30.7% and then in the order of 22.2% of professionals and 16.7% of office staff ($P < 0.001$). For monthly household income, the highest was over 7 million won in 27.5% of men and 20.0% of women. The lowest distribution was under 2 million won as 6.6% in men and 6–6.99 million won as 6.9% in women.

Stress

Table 2 shows the stress score of middle-aged men and women. The perceived stress score by sex was not significantly different in middle-aged subjects: The total stress score was 17.75 ± 4.31 in men and 17.89 ± 4.76 in women, and 17.82 ± 4.55 in all subjects.

Stress according to dietary habits

Table 3 shows the stress score according to dietary habits level of middle-aged men and women. Stress score according to dietary habits was significantly higher in men with 'low' dietary habits level compared to those with 'moderate' and 'high' levels ($P < 0.001$). In women, it was significantly higher in those with 'low' and 'moderate' dietary habits levels compared to those with 'high' level ($P < 0.05$). In all subjects, stress was significantly higher in groups with low dietary habits level ($P < 0.001$).

Table 3. Stress score according to dietary habits level of middle-aged men and women

Classification	Level of dietary habits	Stress score	F	P-value
Male (n = 320)	Low (n = 103) ¹⁾	19.14 ± 3.95 ^a	9.130***	< 0.001
	Moderate (n = 143)	17.36 ± 4.00 ^b		
	High (n = 74)	16.57 ± 4.90 ^b		
	Total	17.75 ± 4.31		
Female (n = 350)	Low (n = 122)	18.64 ± 4.73 ^a	4.393*	0.013
	Moderate (n = 125)	18.05 ± 4.45 ^a		
	High (n = 103)	16.80 ± 4.98 ^b		
	Total	17.89 ± 4.76		
Total (n = 670)	Low (n = 225)	18.87 ± 4.39 ^a	11.823***	< 0.001
	Moderate (n = 268)	17.68 ± 4.22 ^b		
	High (n = 177)	16.70 ± 4.93 ^c		
	Total	17.82 ± 4.55		

Data are shown as mean ± SD. Means with different superscripts are significantly different from each other at $P = 0.05$ by Duncan's multiple range test.

¹⁾Number of person.

* $P < 0.05$, *** $P < 0.001$ by one-way analysis of variance test.

Table 4. Stress score according to emotional eating behavior level of middle-aged men and women

Classification	Level of emotional eating behavior	Stress score	F	P-value
Male (n = 320)	Non-emotional eater (n = 168) ¹⁾	17.25 ± 4.51 ^b	4.219*	0.016
	Low emotional eater (n = 104)	17.85 ± 3.88 ^b		
	Emotional eater (n = 48)	19.27 ± 4.24 ^a		
	Total	17.75 ± 4.31		
Female (n = 350)	Non-emotional eater (n = 115)	16.23 ± 4.36 ^c	22.376***	< 0.001
	Low emotional eater (n = 136)	17.54 ± 4.33 ^b		
	Emotional eater (n = 99)	20.28 ± 4.83 ^a		
	Total	17.89 ± 4.76		
Total (n = 670)	Non-emotional eater (n = 283)	16.83 ± 4.47 ^b	24.554***	< 0.001
	Low emotional eater (n = 240)	17.68 ± 4.14 ^b		
	Emotional eater (n = 147)	19.95 ± 4.66 ^a		
	Total	17.82 ± 4.55		

Data are shown as mean ± SD. Means with different superscripts are significantly different from each other at $P = 0.05$ by Duncan's multiple range test.

¹⁾Number of person.

* $P < 0.05$, *** $P < 0.001$ by one-way analysis of variance test.

Stress according to emotional dietary behavior

Table 4 shows the stress score according to emotional dietary behavior level of middle-aged men and women.

Stress score was higher in emotional eaters in men ($P < 0.05$) and all subjects ($P < 0.001$) with significant differences. In women, stress was higher in emotional eaters, with a significant difference ($P < 0.001$).

Stress according to insomnia level

Table 5 shows the stress score according to insomnia level of middle-aged men and women.

Stress score was high in men, women, and all subjects in the order of group with no clinically significant insomnia < subthreshold insomnia < clinical insomnia, with a significant difference ($P < 0.001$).

Correlations among stress, dietary habits, emotional dietary behavior, and insomnia

Table 6 shows the correlations among stress and dietary habits, emotional dietary behavior, and insomnia.

Table 5. Stress score according to insomnia level of middle-aged men and women

Classification	Level of insomnia	Stress score	F	P-value
Male (n = 320)	No clinically significant insomnia (n = 178) ¹⁾	16.38 ± 4.20 ^c	27.628***	< 0.001
	Subthreshold insomnia (n = 112)	18.97 ± 3.42 ^b		
	Clinical insomnia (n = 30)	21.30 ± 4.68 ^a		
	Total	17.75 ± 4.31		
Female (n = 350)	No clinically significant insomnia (n = 163)	16.13 ± 4.13 ^c	29.397***	< 0.001
	Subthreshold insomnia (n = 128)	18.70 ± 4.58 ^b		
	Clinical insomnia (n = 59)	20.97 ± 4.77 ^a		
	Total	17.89 ± 4.76		
Total (n = 670)	No clinically significant insomnia (n = 341)	16.26 ± 4.16 ^c	56.904***	< 0.001
	Subthreshold insomnia (n = 240)	18.83 ± 4.08 ^b		
	Clinical insomnia (n = 89)	21.08 ± 4.72 ^a		
	Total	17.82 ± 4.55		

Data are shown as mean ± SD. Means with different superscripts are significantly different from each other at $P = 0.05$ by Duncan's multiple range test.

¹⁾Number of person.

*** $P < 0.001$ by one-way analysis of variance test.

Table 6. Correlations among stress, dietary habit, emotional eating behavior and ISI-K of subjects

Classification	Dietary habit <i>r</i> (<i>P</i>)	EEQ <i>r</i> (<i>P</i>)	ISI-K <i>r</i> (<i>P</i>)
PSS			
Man (<i>n</i> = 320) ¹⁾	−0.278*** (< 0.001)	0.183** (0.002)	0.419*** (< 0.001)
Woman (<i>n</i> = 350)	−0.203*** (< 0.001)	0.372*** (< 0.001)	0.399*** (< 0.001)
Total (<i>n</i> = 670)	−0.235*** (< 0.001)	0.291*** (< 0.001)	0.406*** (< 0.001)

PSS, perceived stress scale; EEQ, emotional eating questionnaire; ISI-K, Korean version of the insomnia severity index.

¹⁾Number of person.

P* < 0.01, *P* < 0.001 by Pearson's correlation coefficient.

The results of correlation analysis of stress and dietary habit, emotional dietary behavior, and sleep showed significantly negative correlations between stress and dietary habits in middle-aged men, women, and all subjects (*P* < 0.001), and significantly positive correlations between stress and emotional dietary behavior in men (*P* < 0.01), women (*P* < 0.001), and all subjects (*P* < 0.001). Significantly positive correlation was observed between stress and insomnia in men, women, and all subjects (*P* < 0.001). Therefore, as stress increased in middle-aged subjects, poorer dietary habits and the quality of sleep habit and more emotional dietary behavior was observed.

DISCUSSION

The study was performed to investigate the degree of perceived stress by sex in middle-aged people, and to provide basic data to develop appropriate nutrition education and interventional measures for middle-aged people by determining the associations between stress and dietary habits, emotional dietary behavior, and sleep.

In the study, the perceived stress of middle-aged subjects was 17.75 ± 4.31 in men and 17.89 ± 4.76 in women, and 17.82 ± 4.55 in all subjects. In previous studies using the same stress perception measures, a study on college students showed that the perceived stress score was 15.57 ± 5.27 in male students and 16.67 ± 4.75 in female students [18]. In a study of public officials aged 19–59, the score of perceived stress was 16.17 ± 4.75 [19]. From these results, perceived stress seemed higher in middle-aged subjects compared to other age groups. Perceived stress in middle-aged subjects was not significantly different according to sex.

Stress according to dietary habits was significantly higher in men with low dietary habits level compared to other groups and significantly higher in women with low and moderate dietary habits levels compared to those with higher level. In all subjects, stress was significantly higher in groups with low dietary habits level. These results were consistent with those in a study on stress and dietary habits in adult men and women in Ulsan areas, in which stress was higher in the group with low dietary habits level [20]. Also, results of previous studies supported the results of this study in which it is highly possible for adults with higher stress to have undesirable dietary habits, causing health problems [21–23].

Stress according to emotional dietary behavior level revealed that it was significantly higher in emotional eaters in men and women, compared to less emotional eaters. These results were similar to those from a study in high school girls [24] and the elderly [25], in which higher stress and depression levels were associated with poor dietary behaviors.

Stress was high in men, women, and all subjects as the quality of sleep became worse. From previous studies on sleep and depression, too short or too long sleep hours increased depression [26,27]. Also, in a study of office workers, insufficient sleep caused higher work stress [28]. These results supported the fact that sufficient sleep can be an important factor in stress.

The results of correlation analysis of stress and dietary habit, emotional dietary behavior, and sleep in middle-aged subjects showed a significantly negative correlation between stress and dietary habits and significantly positive correlations between stress, emotional dietary behavior, and insomnia. In the previous research, the results showed the correlations among life stress, sleep, anthropometric measurement and nutrient intakes of college students [29]. Such data support our results that the causes of life stress have an association based on the sleep quality, anthropometric measurement, and nutrient intake.

In conclusion, stress was higher in middle-aged subjects as dietary habits became undesirable, as they were emotional eaters, and as the quality of sleep was lower. Thus, it is considered that stress management programs at the national level are necessary for middle-aged population to relieve and control stress first since stress might be associated with poor dietary habits, emotional dietary behavior, and sleep quality. In addition, it is necessary to investigate researches on lifestyle improvement in middle-aged people and to provide nutrition education for them, and individuals also should actively try for accomplishing these goals.

For the limitations of the study, it was impossible to meet many people as possible due to coronavirus disease 2019 situations and survey targets were middle-aged subjects who might feel uncomfortable for online survey methods; thus, the survey was performed by combining both online and self-administered methods. Since there might be some discrepancies due to survey methods, it is suggested to repeat this study using one method in the future. Also, this study did not reflect objective data of the subjects such as biochemical data or hormonal changes by stress. Therefore, it is suggested to perform a study in the future reflecting health status and objective stress level of the subjects on the basis of biochemical values. Despite these limitations, this study is meaningful in analyzing the stress status according to perceived stress and dietary habits, emotional dietary behavior, and insomnia by sex in middle-aged subjects. Thus, the study results will be used as basic data to develop nutrition education programs and nutrition intervention for lifestyle improvement according to sex and stress in middle-aged people.

REFERENCES

1. Statistics Korea. Life expectancy (life expectancy at 0 years old) and life expectancy without illness (healthy life expectancy) [Internet]. Daejeon: Statistics Korea; 2021 [cited 2021 January 13]. Available from: http://www.index.go.kr/potal/main/EachDtlPageDetail.do?id_cd=2758.
2. Oh HS. Important significant factors of health-related quality of life(EQ-5D) by age group in Korea based on KNHANES (2014). *J Korean Data Inf Sci Soc* 2017;28:573-84.
3. Han KS. Stress of the mid-life stage. *Korean J Stress Res* 2007;15:263-70.
4. Torres SJ, Nowson CA. Relationship between stress, eating behavior, and obesity. *Nutrition* 2007;23:887-94. [PUBMED](#) | [CROSSREF](#)
5. Park HE, Bae Y. Eating habits in accordance with the mental health status: the 5th Korea National Health and Nutrition Examination Survey, 2010–2012. *J Korea Acad Ind Coop Soc* 2016;17:168-81.
6. Choi HJ, Kim BJ, Kim IJ. Prevalence and risk factors of sleep disturbance in community dwelling adults in Korea. *Korean J Adult Nurs* 2013;25:183-93. [CROSSREF](#)

7. Shim HY, Kwon OJ, Kim MJ, Song EJ, Moon SY, Nam YD, Nam DH, Lee JH, Koo BS, Kim HJ. A comparative study on the quality of sleep, tongue diagnosis, and oral microbiome in accordance to the Korean medicine pattern differentiation of insomnia. *J Korean Med Obes Res* 2020;20:40-51.
CROSSREF
8. National Health Insurance Service. Sleep disorder even after sleeping, an average annual increase of 8.1% for 5 years [Internet]. Wonju: National Health Insurance Service; 2019 [cited 2019 September 19]. Available from: <https://www.nhis.or.kr/nhis/together/wbhaea01600m01.do?mode=view&articleNo=131445>.
9. Doo M, Kim Y. Sleep duration and dietary macronutrient consumption can modify the cardiovascular disease for Korean women but not for men. *Lipids Health Dis* 2016;15:17.
PUBMED | CROSSREF
10. Kim CE, Shin S, Lee HW, Lim J, Lee JK, Shin A, Kang D. Association between sleep duration and metabolic syndrome: a cross-sectional study. *BMC Public Health* 2018;18:720.
PUBMED | CROSSREF
11. World Health Organization. Regional Office for the Western Pacific. The Asia-Pacific perspective: redefining obesity and its treatment. Sydney: Health Communications Australia; 2000.
12. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983;24:385-96.
PUBMED | CROSSREF
13. Lee J, Shin C, Ko YH, Lim J, Joe SH, Kim S, Jung IK, Han C. The reliability and validity studies of the Korean version of the perceived stress scale. *Korean J Psychosom Med* 2012;20:127-34.
14. Garaulet M, Canteras M, Morales E, López-Guimera G, Sánchez-Carracedo D, Corbalán-Tutau MD. Validation of a questionnaire on emotional eating for use in cases of obesity: the emotional eater questionnaire (EEQ). *Nutr Hosp* 2012;27:645-51.
PUBMED
15. Saleh-Ghadimi S, Dehghan P, Abbasalizad Farhangi M, Asghari-Jafarabadi M, Jafari-Vayghan H. Could emotional eating act as a mediator between sleep quality and food intake in female students? *Biopsychosoc Med* 2019;13:15.
PUBMED | CROSSREF
16. Bastien CH, Vallières A, Morin CM. Validation of the insomnia severity index as an outcome measure for insomnia research. *Sleep Med* 2001;2:297-307.
PUBMED | CROSSREF
17. Cho YW, Song ML, Morin CM. Validation of a Korean version of the insomnia severity index. *J Clin Neurol* 2014;10:210-5.
PUBMED | CROSSREF
18. Jo HM. The association between sleep quality and perceived stress among students from 2 universities [master's thesis]. Seoul: Ewha Womans University; 2015.
19. Park SY, Joe SH, Kim SH, Han CS, Ham BJ, Ko YH. Fatigue and its association with socio-demographic and clinical variables in a working population. *Korean J Psychosom Med* 2014;22:3-12.
20. Kim HK, Kim JH. Relationship between stress and eating habits of adults in Ulsan. *Korean J Nutr* 2009;42:536-46.
CROSSREF
21. Seo EY, Lee SL. Factors influencing dietary behaviors and stress in male and female college students. *J Korean Soc Sch Health* 2018;31:186-95.
22. Seo YJ, Kim MH, Kim MH, Choi MK. Status and relationships among lifestyle, food habits, and stress scores of adults in Chungnam. *Korean J Community Nutr* 2012;17:579-88.
CROSSREF
23. Yoon HS. An assessment on the dietary attitudes, stress level and nutrient intakes by food record of food and nutrition major female university students. *Korean J Nutr* 2006;39:145-59.
24. Park JE, Kim SJ, Choue RW. Study on stress, depression, binge eating, and food behavior of high school girls based on their BMI. *Korean J Community Nutr* 2009;14:175-81.
25. Lee SJ, Kim YR, Seo SH, Cho MS. A study on dietary habits and food intakes in adults aged 50 or older according to depression status. *J Nutr Health* 2014;47:67-76.
CROSSREF
26. Lee YJ. Gender differences in factors associated with the severity of depression in middle-aged adults: an analysis of 2014 Korean National Health and Nutrition Examination Survey. *J Korea Converg Soc* 2018;9:549-59.

27. Watson NF, Harden KP, Buchwald D, Vitiello MV, Pack AI, Strachan E, Goldberg J. Sleep duration and depressive symptoms: a gene-environment interaction. *Sleep* 2014;37:351-8.
[PUBMED](#) | [CROSSREF](#)
28. Lee MA, Lee EJ, Soh HK, Choi BS. Analysis on stress and dietary attitudes of male employees. *Korean J Community Nutr* 2011;16:337-52.
[CROSSREF](#)
29. Sung MJ, Chang KJ. Correlations among life stress, sleep, anthropometric measurement and nutrient intakes of college students. *J Korean Soc Food Sci Nutr* 2007;36:840-8.
[CROSSREF](#)