

Prevalence and Management of Functional Gastrointestinal Disorders in Infants: An Asian Perspective

Wei Cai, Lalit Bharadia*, Mohammad Juffrie†, Fook Choe Cheah‡, Seng Hock Quak§, Vitaya Titapant||, Ruurd van Elburg¶, and Thomas Ludwig**

Xin Hua Hospital, Shanghai Jia Tong University School of Medicine, Shanghai, China, *Santokba Durlabhji Memorial Hospital, Jaipur, India, †Department of Pediatrics, Faculty of Medicine, Universitas Gadjah Mada, Yogyakarta, Indonesia, ‡Department of Paediatrics, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia, §Department of Paediatrics, National University Hospital, Singapore, ||Siriraj Hospital, Mahidol University, Bangkok, Thailand, ¶Nutricia Research, Utrecht, The Netherlands, **Nutricia Research, Singapore

To the Editor:

Scarpato et al. [1] have reported in a recent publication in *J Pediatr Gastroenterol Nutr* that in a number of surveyed Mediterranean countries the Rome diagnostic criteria are not consistently applied, and that large variability remains in the management of functional gastrointestinal disorders (FGIDs). It is known that FGIDs are very common in Europe in children up to three years of age, but little data exists on the prevalence and management of FGIDs in Asia. At a recent workshop and literature review, a group of health care professionals from China, Indonesia, Malaysia, Singapore, Thailand, Hong Kong, and India has reviewed the available data for their respective countries. Based on a systematic literature search (Embase, CAB Abstracts, MEDLINE, SciSearch, Current Contents, Biosis Previews, Pascal, Emcare, and PubMed), it was concluded that data on the prevalence of FGIDs in Asia, e.g., functional constipation ranging from 0.05% to approximately 7% in

Thailand, is scarce with considerable variation depending on the applied criteria, study design, methodology, and/or study populations. Few countries have clear guidelines on the diagnosis and management of FGIDs, and dissemination and implementation is unclear. There is a huge variety of use of over the counter remedies by parents, indicative of the absence of safety, tolerance, and efficacy data. It was concluded that I) well-designed epidemiological studies enabling comparison of the prevalence of FGIDs between Asian countries are urgently needed, II) education of parents, parental reassurance, and nutritional management should be the pillars in the management of FGIDs, and III) the potential and opportunity in nutritional prevention and management of FGIDs in Asia could be further advanced.

ACKNOWLEDGEMENTS

All authors have received unconditional support by

Received : September 27, 2017, Accepted : October 13, 2017

Corresponding author: Thomas Ludwig, Nutricia Research, 30 Biopolis Street, Singapore. Tel: +65-8498-2859, E-mail: Thomas.Ludwig @Danone.com

Copyright © 2018 by The Korean Society of Pediatric Gastroenterology, Hepatology and Nutrition

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nutricia Research for their travel expenses in the participation of a round table discussion. RvE and TL are employees of Nutricia Research.

REFERENCE

1. Scarpato E, Quitadamo P, Roman E, Jojkic-Pavkov D, Kolacek S, Papadopoulou A, et al. Functional gastrointestinal disorders in children: a survey on clinical approach in the mediterranean area. *J Pediatr Gastroenterol Nutr* 2017;64:e142-6.