



Epigenetics of Male Fertility: Effects on Assisted Reproductive Techniques

Filippo Giacone^{ID}, Rossella Cannarella^{ID}, Laura M. Mongioì^{ID}, Angela Alamo^{ID}, Rosita A. Condorelli^{ID}, Aldo E. Calogero^{ID}, Sandro La Vignera^{ID}

Department of Clinical and Experimental Medicine, University of Catania, Catania, Italy

During the last decades the study of male infertility and the introduction of the assisted reproductive techniques (ARTs) has allowed to understand that normal sperm parameters do not always predict fertilization. Sperm genetic components could play an important role in the early stages of embryonic development. Based on these acquisitions, several epigenetic investigations have been developed on spermatozoa, with the aim of understanding the multifactorial etiology of male infertility and of showing whether embryonic development may be influenced by sperm epigenetic abnormalities. This article reviews the possible epigenetic modifications of spermatozoa and their effects on male fertility, embryonic development and ART outcome. It focuses mainly on sperm DNA methylation, chromatin remodeling, histone modifications and RNAs.

Keywords: Epigenomics; Infertility; Infertility, male; Reproductive techniques, assisted

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Semen analysis is a first level diagnostic investigation for male infertility (as well as for the testicular function), since it is able to give indication on sperm fertility potential. However, the implementation of several assisted reproductive techniques (ARTs) has led to the observation that normal sperm parameters are not always capable of inducing pregnancy in a healthy woman. For this reason, several studies on sperm functional capability and bio-functional sperm parameters have been developed [1,2]. Nonetheless the evaluation of these variables is often insufficient to identify causes of male infertility [3]. To overcome this pitfall, molecular biologists have recently tried to study

the epigenetic origin of male infertility and embryonic anomalies [4].

Epigenetics study meiotic and/or mitotic inherited changes affecting gene expression without a DNA sequence modification [5]. These studies allow in some cases to better evaluate cases of so-called idiopathic infertility. DNA methylation, chromatin remodeling and residual histone modifications represent the major epigenetic changes occurring at the sperm level [6,7]. Recently, sperm RNAs seem also to play an important role [8].

This article reviews the implications of epigenetic modifications on male fertility, fertilization, embryonic development and ART outcomes, highlighting the possible mechanisms involved. To accomplish this, we

Received: Aug 6, 2018 **Revised:** Sep 16, 2018 **Accepted:** Sep 28, 2018 **Published online** Nov 27, 2018

Correspondence to: Sandro La Vignera ^{ID} <https://orcid.org/0000-0002-7113-2372>

Department of Clinical and Experimental Medicine, University of Catania, Via S. Sofia 78, 95123, Catania, Italy.

Tel: +39-95-3782641, **Fax:** +39-95-3781180, **E-mail:** sandrolavignera@unict.it

performed a search on Pubmed, Science-direct, Ovid and Scopus, using the following key words: sperm DNA methylation, sperm DNA methylation and assisted reproduction technology, sperm chromatin remodeling, sperm chromatin remodeling and assisted reproduction technology, sperm histone modification, sperm histone modification sperm RNAs and assisted reproduction technology.

DNA METHYLATION

DNA methylation is a dynamic mechanism of gene expression regulation that occurs on dinucleotides formed by cytosine and guanine. These dinucleotides are grouped in the so called differently methylated regions (DMRs) that are often placed near gene regulatory regions, such as the promoter [9]. The methylation processes are regulated by DNA methyltransferases [9]. Hypermethylation suppresses gene expression, since methyl groups prevent the recruitment of transcription factors and DNA polymerases. Conversely, hypomethylation promotes gene expression [7,10].

DNA methylation includes global DNA methylation, methylation of imprinted and non-imprinted genes and of methylation of repetitive elements.

A global DNA hypomethylation seems to occur in association with ART [11]. Furthermore, an increased risk of imprinted disorders has been reported after these procedures, thus leading to the hypothesis that ART itself may cause a loss of methylation [11]. Some studies described a significantly different global DNA methylation in sperm from patients with oligoasthenoteratozoospermia (OAT) compared to controls [12], indicating that such epigenetic abnormality may reflect on human fertility. It cannot be excluded whether a change in sperm global DNA methylation may primary affect ART outcome. Indeed, sperm global DNA methylation has been related with chromatin injury in samples from OAT patients [12,13] and, interestingly, chromatin integrity seems to impact on ART (see section "CHROMATIN REMODELING"). Thus, sperm global methylation may indirectly influence ART outcome.

Methylation of repetitive elements (*e.g.*, ALU and LINE1) at the sperm level has been reported to influence ART outcome. More in detail, ALU methylation impacts on pregnancy and live-birth rates in couples with male-factor and combined infertility, being significantly lower in sperm samples that led to abortion

compared to those leading to pregnancy and live-birth [14].

Imprinted genes show parent-specific activity (they are functionally haploids), which makes them particularly vulnerable to epigenetic dysregulation. In greater detail, both paternal and maternal alleles undergo to demethylation after fertilization. Then, genetic reprogramming consisting in new specific methylations takes place in the embryo [15]. Imprinted genes escape epigenetic reprogramming after fertilization, thus allowing the possible transmission of aberrant methylation patterns into the offspring.

Imprinting has been extensively studied in humans and mice. The role of imprinted genes in processes involved in embryo and placenta development and growth is well recognized in mice (Table 1) [16]. Since humans and mice have common placental physiology [17], these mechanisms may occur also in humans. Knowledge in humans mainly derives from pediatric growth disorders. However, epigenetic studies in human sperm and placenta allow to understand the role of imprinted genes in human fertilization. Accordingly, impaired methylation of imprinted genes has been reported in infertile patients [18]. Furthermore, modifications in sperm DNA methylation of such genes play a role in the development of male infertility and negatively impact on the ART outcomes, leading to abortion [19]. Meta-analytic data provide evidence for lower *H19* and higher *MEST* and *SNRPN* sperm methylation in infertile patients compared to fertile controls [18].

A list of imprinted genes involved in sperm quality is shown in Table 2 [18-27]. A focus on the role of *H19* DMR impaired methylation rate in human fertility is reported below.

1. *H19/IGF2*

H19/IGF2 represents the first historically charac-

Table 1. Imprinted genes involved in placenta and embryo development and growth in mice [16]

Process	Imprinted gene
Placenta establishing	<i>Peg10</i>
Nutrient transport capacity and surface area for exchange	<i>Igf2, Grb10</i>
Fetal growth	<i>Igf2, Igf2r, Cdkn1c, Grb10</i>

Cdkn1c: cyclin-dependent kinase inhibitor 1c, *Grb10*: growth factor bound protein 10, *Igf2*: insulin-like growth factor 2, *Igf2r*: insulin-like growth factor 2 receptor, *Peg10*: paternally expressed 10.

Table 2. Imprinted genes involved in sperm quality

Gene	Evidence	Reference
<i>CREM</i>	Increased methylation is associated with decreased semen quality	20, 21
<i>DAZL</i>	Increased methylation in OAT patients compared to controls	22
<i>FAM50B</i>	Reduced methylation levels are associated with asthenozoospermia	23
<i>GNAS</i>	Reduced methylation levels are associated with asthenozoospermia	23
<i>GLT2</i>	Abnormal methylation levels are associated with oligozoospermia	24, 25
<i>H19</i>	Reduced methylation associates with male infertility	18
<i>KCNQ1OT1</i>	Increased methylation in patients with abnormal sperm parameters	25
<i>MEST</i>	Increased methylation associates with male infertility	18
<i>RHOX</i>	Increased methylation associates with male infertility	26
<i>SNRPN</i>	Increased methylation associates with male infertility	18
<i>ZAC</i>	Increased methylation associates with oligozoospermia	27

CREM: cAMP responsive element modulator, *DAZL*: deleted in azoospermia-like, *FAM50B*: family with sequence similarity 50, member B, *GNAS*: guanidine nucleotide-binding protein, alpha-stimulating activity polypeptide 1, *GLT2*: gene trap locus 2 (also known as MEG3, maternally expressed gene 3), *H19*: imprinted maternally expressed non-coding transcript, *KCNQ1OT1*: *KCNQ1*-overlapping transcript 1, *MEST*: mesoderm-specific transcript, mouse, homolog of, *RHOX*: reproductive homeobox X-linked, *SNRPN*: small nuclear ribonucleoprotein polypeptide N, *ZAC*: Zac tumor suppression gene (also known as PLAGL1, pleomorphic adenoma gene-like 1).

terized couple of imprinted genes [28]. Both *H19* and *IGF2* map on the 11p15.5 chromosome. *H19* encodes for a long non-coding RNA (lncRNA), which negatively modulates human placental trophoblast cell proliferation [29]. In particular, lncRNA *H19* targets miR-675, which, in turn, represses *IGF1R* transcription [30]. *IGF2* encodes for a growth factor which is able to activate the insulin-like growth factor-1 receptor (IGF1R) and to promote fetal and placental growth [31-34]. In contrast to IGF1, which is preferentially expressed after birth, IGF2 is mainly produced in the early embryonic development. On these basis, it might be hypothesized that a low sperm *IGF2* gene expression might negatively influence the outcome of pregnancy, though no study has investigated this aspect so far.

IGF2 gene expression is regulated by the *H19* DMR methylation rate. In the maternal allele, *H19* DMR is

unmethylated, thus allowing *H19* expression and preventing the accession to the *IGF2* gene enhancer. In the paternal allele, the *H19* DMR methylation leads to *IGF2* gene expression [35]. Several studies described the occurrence of low *H19* DMR methylation rates in infertile patients [18,36-39]. At the sperm level, this may lead to low *IGF2* gene expression, thus affecting the outcome of pregnancy.

A study carried on 119 human placentas (56 from *in vitro* fertilization [IVF], 41 from intracytoplasmic sperm injection [ICSI] and 22 from natural conception) found 1.9 and 1.8-fold higher *H19* messenger RNA (mRNA) levels, respectively in ICSI and IVF-derived placentas compared with those coming from natural conception. In addition, IGF2 mRNA levels resulted significantly lower in couples undergoing ART compared to that of controls. Furthermore, although both birth and placental weight were lower in the ART group, they were not significantly different compared to controls [40]. This study provides evidence for abnormal *H19* and *IGF2* expression in ART placentas. Due to abnormal *H19* DMR methylation rate observed in infertile patients [18], men undergoing to ART programs may likely show *H19* gene hypomethylation. This might explain the abnormal *H19* and *IGF2* expression in ART placentas [40].

Finally, IGF1R has been recognized to play a role in human sperm capacitation [41]. Therefore, due to the inhibitory effect that *H19* has on *IGF1R* gene transcription [30], it might be hypothesized that low *H19* DMR methylation rates may also impair *IGF1R* gene expression, thus affecting human sperm capacitation.

Taking all this into account, *H19/IGF2* gene expression may influence human fertility (Fig. 1). Other studies relating pregnancy rate with IGF2 and IGF1R mRNA levels in human spermatozoa may further clarify their role.

CHROMATIN REMODELING

Spermatozoa are highly specialized cells. During spermatogenesis, almost all chromatin histones (90%-95%) are replaced by protamines, that are small, arginine-rich, nuclear proteins [42]. This process, typical of spermatozoa, ensures a significant DNA compaction, a decreased susceptibility to external insults and it represents a gene silencing mechanism. During early stages of protamination, there is an increased acety-




Fig. 1. Possible influence of H19/IGF2 expression on male fertility. (A) In the paternal allele of healthy fertile men, H19 DMR methylation leads to *IGF2* expression. The repression of *H19* transcription increase *IGF1R* expression. Both *IGF2* and *IGF1R* transcripts are involved in sperm capacitation, embryo and placental growth. (B) In infertile men, low H19 DMR methylation rates lead to repression of both *IGF2* and *IGF1R* transcription, thus negatively impacting on sperm capacitation, embryo and placental growth. H19: Imprinted maternally expressed non-coding transcript, IGF2: insulin-like growth, DMR: differentially methylated region, IGF1R: insulin-like growth factor 1 receptor, factor 2.

lation of histones that firstly promotes the action of DNA topoisomerase enzyme, followed by the substitution of histones with transition proteins (TP1 and TP2) [43]. These DNA binding proteins facilitate histones removal and subsequent replacement of protamines 1 (P1) and 2 (P2), which are expressed in equal amounts [44]. After sperm chromatin protamination, three levels of DNA organization can be recognized: toroidal structures formed by protamines (90%–95%), nucleosomes, involved in the primary phases of embryonic development (5%–10%) and DNA segments called matrix attachment regions (MARs), in which neither toroidal structures nor nucleosomes are present [43]. MARs provide chromatin with structural support, working as promoter in paternal pronuclear formation after fertilization, as well as contributing to normal embryogenesis.

Several studies reported that an erroneous sperm protamination or an alteration in one of the three DNA organization levels leads to a lower sperm quality and decreased pregnancy rate and embryo quality following ARTs [6,42]. In greater detail, a P1/P2 ratio ranging from 0.54 to 1.43 has been reported in normozoospermic men [45]. Abnormal low or high P1/P2 ratio is associated with poor sperm [45-47] and embryo qual-

ity and with low fertilization and pregnancy rates [48]. Indeed, abnormal sperm protamination may result in insufficient sperm DNA condensation, thus making it susceptible to the oxidative damage [49]. Accordingly, impaired sperm protamination has been related to increased DNA fragmentation [48,50]. Furthermore, several factors negatively affecting male fertility and sperm quality have been found to impair sperm chromatin compactness, such as overweight and obesity [51], nicotine [52,53], male accessory gland infections [54] and varicocele [55]. Lower chromatin compactness and poor sperm quality have been described in patients with low testicular volume [56].

In conclusion, recent scientific evidence showed that anomalies in protamine content may affect epigenetic information transmitted by the paternal DNA [49] and, not surprisingly, sperm protamination has been already suggested as a target to predict ART outcomes [57].

HISTONE MODIFICATION

After protamination, the residual histones (5%–10%) are organized into nucleosomes that are octamers composed by two dimers of H3-H4 and two of H2A-H2B

histones. These residual histones play a crucial role in sperm epigenetics. In fact, they might undergo chemical modifications that might drastically alter their DNA binding abilities, favoring the action of transcription factors and therefore gene activation or silencing [58]. These chemical modifications are mainly referred to lysine or serine residues located on the histone tail, undergoing to methylation, acetylation, phosphorylation or ubiquitination [42]. Moreover, histones may have variants such as the H2B, that is a testis-specific histone variant. Methylation is one of the main histone modifications: it is regulated by the enzyme methyltransferases and is generally associated with gene silencing [7]. Acetylation is regulated by acetyltransferase and deacetylates histones: the first one activates gene expression whereas the second inhibits it. Histone phosphorylation takes place on lysine residues and generally promotes gene activation [7]. Lastly, ubiquitination is a covalent modification that occurs on lysine residues and it is capable of activating or repressing gene expression [7].

Since retained histones are present in imprinted gene clusters [49], changes in protamination and in residual histones can be co-responsible for male infertility. Accordingly, a study carried out on 291 ART cycles described the role of histone-to-protamine ratio (HPR) on embryo development and ART outcome. For HPR ranging from 6 and 26%, the blastocyst formation rate was significantly higher (87.8%) compared to that obtained with HPR <6% (71.2%) or >6% (74.6%). Therefore, HPR seems to influence embryo development [59]. On the basis of such evidence, sperm devoid of protamine and residual histones abnormalities should be utilized in ART programs [49].

SPERM RNA

Sperm carries thousands of different RNAs, including coding (mRNA) and non-coding [micro-RNA (miRNA), small interfering (siRNA), antisense RNA, piwi-interacting RNA (piRNA) and lncRNA] RNAs [60], the latter known to be involved in the modulation of gene expression, interfering with the mRNA translation through different mechanisms.

Overall, more than 4,000 different mRNAs have been detected in human spermatozoa [61]. Due to sperm chromatin compactness, it was thought that sperm RNA may unlikely be transcribed from sperm nuclear

DNA and that it may be a residual of spermatogenesis [62,63]. However, there is some evidence of transcriptional activity in spermatozoa [64,65], thus suggesting that it may derive from imprinted genes mapping in unmethylated histones, where the transcriptional apparatus can access.

Sperm RNAs have been discovered in mammals, insects [66] and plants (pollen RNA [67]), suggesting that it might represent a conserved characteristic of the male gamete [68]. As far for sperm RNA functions, they are mostly completely unknown [66,69,70]. Gathering together the available evidence from both animal and human studies, three main roles may be hypothesized: transmission of parentally acquired phenotype, spermatozoa maturation in the epididymis and embryo development, as discussed below.

Mounting evidence has shown that sperm RNAs are involved in the transmission of paternally acquired phenotypes [8]. Recently, the comprehensive profiling of sperm epigenome from obese and lean men resulted in a different sperm non-coding RNA expression, suggesting that sperm RNAs may act as a sensitive marker of environmental exposure [71]. Sperm RNAs are capable to modulate early embryonic gene expression [8]. Therefore, due to the higher risk of developing obesity which children of obese fathers are exposed to, the differential sperm RNA content of obese patients has been hypothesized as the possible mechanism through which obesity is inherited by the offspring [71].

Mammalian sperm RNAs are involved in sperm maturation [72,73]. This role might be displayed by piRNAs, that are 30 nucleotide-long non-coding RNAs interacting with the so called Piwi proteins. The Piwi/piRNA machinery acts in gene silencing that occurs in the male gamete [74]. During spermiogenesis, Piwi proteins seem to mediate the histone-to-protamine exchange in a piRNA-independent manner in mice. *Piwi* germline mutations preventing Piwi ubiquitination and degradation have been reported in azoospermic patients [75]. Despite further research is needed, the Piwi/piRNA machinery may be involved in human spermiogenesis, during the phase of late spermatids [75], probably acting on sperm DNA protamination.

Finally, sperm RNAs seem to be play a role in embryo development. Sperm RNAs are carried into the oocyte and, hence, translated [68]. A recent study performed in mice reported a significant decrease in blastocyst formation rate after ICSI when an RNase-treat-

ed sperm was used [76]. The treatment resulted in a 90% reduction of the sperm RNA, thus supporting that the latter is required for mice embryo development [76]. No study has explored the relationship between sperm specific transcripts and ART outcomes so far.

In conclusion, sperm RNAs may be involved in several important functions. The possible role that they may play in human fertilization and embryo growth needs to be elucidated.

CONCLUSIONS

The study of the epigenetic pattern of spermatozoa can help to better understand causes of male infertility. Indeed, specific epigenetic aberrations can affect primary phases of embryonic development and ART outcomes. In particular, P1/P2 ratio, HPR and H19 DMR methylation rate may represent additional targets to investigate sperm quality before ARTs. Furthermore, sperm IGF2 and IGF1R expression should be further investigated.

In conclusion, imprinted unmethylated paternally-expressed genes in residual histones may be accessed from the transcriptional apparatus. It may be hypothesized that once transcribed, these genes contribute to the pool of sperm RNA. Abnormal DMR methylation rate of imprinted genes, alterations in chromatin compactness or in HPR represent possible mechanisms through which sperm RNA content, fertilization, embryo growth and, therefore, ART outcomes may be affected. Research characterizing sperm RNAs and their role in human fertility is needed.

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contribution

Concept and Design: La Vignera S, Giacone F. Acquisition of data or analysis: Calogero AE, Cannarella R, Mongioi LM, Alamo A, Condorelli RA. Final approval: Calogero AE, Giacone F, La Vignera S. All authors read and approved the final manuscript, and they have the accountability for all aspects of work.

REFERENCES

1. Giacone F, Condorelli RA, Mongioi LM, Bullara V, La Vignera

- S, Calogero AE. In vitro effects of zinc, D-aspartic acid, and coenzyme-Q10 on sperm function. *Endocrine* 2017;56:408-15.
2. Condorelli RA, La Vignera S, Mongioi LM, Vitale SG, Laganà AS, Cimino L, et al. Myo-inositol as a male fertility molecule: speed them up! *Eur Rev Med Pharmacol Sci* 2017;21:S30-5.
3. Cissen M, Wely MV, Scholten I, Mansell S, Bruin JP, Mol BW, et al. Measuring sperm DNA fragmentation and clinical outcomes of medically assisted reproduction: a systematic review and meta-analysis. *PLoS One* 2016;11:e0165125.
4. Karaca MZ, Konac E, Yurteri B, Bozdog G, Sogutdelen E, Bilen CY. Association between methylenetetrahydrofolate reductase (MTHFR) gene promoter hypermethylation and the risk of idiopathic male infertility. *Andrologia* 2017;49. doi: 10.1111/and.12698.
5. Jenkins TG, Aston KI, James ER, Carrell DT. Sperm epigenetics in the study of male fertility, offspring health, and potential clinical applications. *Syst Biol Reprod Med* 2017;63:69-76.
6. Gannon JR, Emery BR, Jenkins TG, Carrell DT. The sperm epigenome: implications for the embryo. *Adv Exp Med Biol* 2014;791:53-66.
7. Rajender S, Avery K, Agarwal A. Epigenetics, spermatogenesis and male infertility. *Mutat Res* 2011;727:62-71.
8. Chen Q, Yan W, Duan E. Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications. *Nat Rev Genet* 2016;17:733-43.
9. Uysal F, Akkoyunlu G, Ozturk S. DNA methyltransferases exhibit dynamic expression during spermatogenesis. *Reprod Biomed Online* 2016;33:690-702.
10. Gunes S, Agarwal A, Henkel R, Mahmutoglu AM, Sharma R, Esteves SC, et al. Association between promoter methylation of MLH1 and MSH2 and reactive oxygen species in oligozoospermic men: a pilot study. *Andrologia* 2018;50. doi: 10.1111/and.12903.
11. Owen CM, Segars JH Jr. Imprinting disorders and assisted reproductive technology. *Semin Reprod Med* 2009;27:417-28.
12. Rahiminia T, Yazd EF, Fesahat F, Moein MR, Mirjalili AM, Talebi AR. Sperm chromatin and DNA integrity, methyltransferase mRNA levels, and global DNA methylation in oligoasthenoteratozoospermia. *Clin Exp Reprod Med* 2018;45:17-24.
13. Olszewska M, Barciszewska MZ, Fraczek M, Huleyuk N, Chernykh VB, Zastavna D, et al. Global methylation status of sperm DNA in carriers of chromosome structural aberrations. *Asian J Androl* 2017;19:117-24.
14. El Hajj N, Zechner U, Schneider E, Tresch A, Gromoll J, Hahn T, et al. Methylation status of imprinted genes and repeti-

- tive elements in sperm DNA from infertile males. *Sex Dev* 2011;5:60-9.
15. Haaf T. Methylation dynamics in the early mammalian embryo: implications of genome reprogramming defects for development. *Curr Top Microbiol Immunol* 2006;310:13-22.
 16. Cassidy FC, Charalambous M. Genomic imprinting, growth and maternal-fetal interactions. *J Exp Biol* 2018;221. pii: jeb164517.
 17. Rai A, Cross JC. Development of the hemochorial maternal vascular spaces in the placenta through endothelial and vasculogenic mimicry. *Dev Biol* 2014;387:131-41.
 18. Santi D, De Vincentis S, Magnani E, Spaggiari G. Impairment of sperm DNA methylation in male infertility: a meta-analytic study. *Andrology* 2017;5:695-703.
 19. Camprubí C, Pladevall M, Grossmann M, Garrido N, Pons MC, Blanco J. Semen samples showing an increased rate of spermatozoa with imprinting errors have a negligible effect in the outcome of assisted reproduction techniques. *Epigenetics* 2012;7:1115-24.
 20. Nanassy L, Carrell DT. Analysis of the methylation pattern of six gene promoters in sperm of men with abnormal protamination. *Asian J Androl* 2011;13:342-6.
 21. Nanassy L, Carrell DT. Abnormal methylation of the promoter of CREM is broadly associated with male factor infertility and poor sperm quality but is improved in sperm selected by density gradient centrifugation. *Fertil Steril* 2011;95:2310-4.
 22. Navarro-Costa P, Nogueira P, Carvalho M, Leal F, Cordeiro I, Calhaz-Jorge C, et al. Incorrect DNA methylation of the DAZL promoter CpG island associates with defective human sperm. *Hum Reprod* 2010;25:2647-54.
 23. Xu J, Zhang A, Zhang Z, Wang P, Qian Y, He L, et al. DNA methylation levels of imprinted and nonimprinted genes DMRs associated with defective human spermatozoa. *Andrologia* 2016;48:939-47.
 24. Sato A, Hiura H, Okae H, Miyauchi N, Abe Y, Utsunomiya T, et al. Assessing loss of imprint methylation in sperm from subfertile men using novel methylation polymerase chain reaction Luminex analysis. *Fertil Steril* 2011;95:129-34, 134.e1-4.
 25. Laurentino S, Beygo J, Nordhoff V, Kliesch S, Wistuba J, Borgmann J, et al. Epigenetic germline mosaicism in infertile men. *Hum Mol Genet* 2015;24:1295-304.
 26. Richardson ME, Bleiziffer A, Tüttelmann F, Gromoll J, Wilkinson MF. Epigenetic regulation of the RHOX homeobox gene cluster and its association with human male infertility. *Hum Mol Genet* 2014;23:12-23.
 27. Hammoud SS, Purwar J, Pflueger C, Cairns BR, Carrell DT. Alterations in sperm DNA methylation patterns at imprinted loci in two classes of infertility. *Fertil Steril* 2010;94:1728-33.
 28. DeChiara TM, Robertson EJ, Efstratiadis A. Parental imprinting of the mouse insulin-like growth factor II gene. *Cell* 1991;64:849-59.
 29. Gao WL, Liu M, Yang Y, Yang H, Liao Q, Bai Y, et al. The imprinted H19 gene regulates human placental trophoblast cell proliferation via encoding miR-675 that targets Nodal modulator 1 (NOMO1). *RNA Biol* 2012;9:1002-10.
 30. Keniry A, Oxley D, Monnier P, Kyba M, Dandolo L, Smits G, et al. The H19 lincRNA is a developmental reservoir of miR-675 that suppresses growth and Igf1r. *Nat Cell Biol* 2012;14:659-65.
 31. Baker J, Liu JP, Robertson EJ, Efstratiadis A. Role of insulin-like growth factors in embryonic and postnatal growth. *Cell* 1993;75:73-82.
 32. Constância M, Hemberger M, Hughes J, Dean W, Ferguson-Smith A, Fundele R, et al. Placental-specific IGF-II is a major modulator of placental and fetal growth. *Nature* 2002;417:945-8.
 33. Fowden AL. The insulin-like growth factors and feto-placental growth. *Placenta*. 2003;24:803-12.
 34. Qiu Q, Basak A, Mbikay M, Tsang BK, Gruslin A. Role of pro-IGF-II processing by proprotein convertase 4 in human placental development. *Proc Natl Acad Sci U S A* 2005;102:11047-52.
 35. Arney KL. H19 and Igf2: enhancing the confusion? *Trends Genet* 2003;19:17-23.
 36. Marques CJ, Carvalho F, Sousa M, Barros A. Genomic imprinting in disruptive spermatogenesis. *Lancet* 2004;363:1700-2.
 37. Marques CJ, Costa P, Vaz B, Carvalho F, Fernandes S, Barros A, et al. Abnormal methylation of imprinted genes in human sperm is associated with oligozoospermia. *Mol Hum Reprod* 2008;14:67-74.
 38. Boissonnas CC, Abdalaoui HE, Haelewyn V, Fauque P, Dupont JM, Gut I, et al. Specific epigenetic alterations of IGF2-H19 locus in spermatozoa from infertile men. *Eur J Hum Genet* 2010;18:73-80.
 39. Kobayashi H, Sato A, Otsu E, Hiura H, Tomatsu C, Utsunomiya T, et al. Aberrant DNA methylation of imprinted loci in sperm from oligospermic patients. *Hum Mol Genet* 2007;16:2542-51.
 40. Sakian S, Louie K, Wong EC, Havelock J, Kashyap S, Rowe T, et al. Altered gene expression of H19 and IGF2 in placentas from ART pregnancies. *Placenta* 2015;36:1100-5.
 41. Wang J, Qi L, Huang S, Zhou T, Guo Y, Wang G, et al. Quantitative phosphoproteomics analysis reveals a key role of insulin growth factor 1 receptor (IGF1R) tyrosine kinase in human

- sperm capacitation. *Mol Cell Proteomics* 2015;14:1104-12.
42. Carrell DT. Epigenetics of the male gamete. *Fertil Steril* 2012;97:267-74.
 43. Johnson GD, Lalancette C, Linnemann AK, Leduc F, Boissonneault G, Krawetz SA. The sperm nucleus: chromatin, RNA, and the nuclear matrix. *Reproduction* 2011;141:21-36.
 44. Grassetti D, Paoli D, Gallo M, D'Ambrosio A, Lombardo F, Lenzi A, et al. Protamine-1 and -2 polymorphisms and gene expression in male infertility: an Italian study. *J Endocrinol Invest* 2012;35:882-8.
 45. Nanassy L, Liu L, Griffin J, Carrell DT. The clinical utility of the protamine 1/protamine 2 ratio in sperm. *Protein Pept Lett* 2011;18:772-7.
 46. Lakpour N, Mahfouz RZ, Akhondi MM, Agarwal A, Kharrazi H, Zeraati H, et al. Relationship of seminal plasma antioxidants and serum male hormones with sperm chromatin status in male factor infertility. *Syst Biol Reprod Med* 2012;58:236-44.
 47. Utsuno H, Miyamoto T, Oka K, Shiozawa T. Morphological alterations in protamine-deficient spermatozoa. *Hum Reprod* 2014;29:2374-81.
 48. Simon L, Castillo J, Oliva R, Lewis SE. Relationships between human sperm protamines, DNA damage and assisted reproduction outcomes. *Reprod Biomed Online* 2011;23:724-34.
 49. Francis S, Yelumalai S, Jones C, Coward K. Aberrant protamine content in sperm and consequential implications for infertility treatment. *Hum Fertil (Camb)* 2014;17:80-9.
 50. Castillo J, Simon L, de Mateo S, Lewis S, Oliva R. Protamine/DNA ratios and DNA damage in native and density gradient centrifuged sperm from infertile patients. *J Androl* 2011;32:324-32.
 51. La Vignera S, Condorelli RA, Vicari E, Calogero AE. Negative effect of increased body weight on sperm conventional and nonconventional flow cytometric sperm parameters. *J Androl* 2012;33:53-8.
 52. Condorelli RA, La Vignera S, Giaccone F, Iacoviello L, Vicari E, Mongioi' L, et al. In vitro effects of nicotine on sperm motility and bio-functional flow cytometry sperm parameters. *Int J Immunopathol Pharmacol* 2013;26:739-46.
 53. Condorelli RA, La Vignera S, Giaccone F, Iacoviello L, Mongioi LM, Li Volti G, et al. Nicotine effects and receptor expression on human spermatozoa: possible neuroendocrine mechanism. *Front Physiol* 2017;8:177.
 54. La Vignera S, Condorelli R, D'Agata R, Vicari E, Calogero AE. Semen alterations and flow-citometry evaluation in patients with male accessory gland infections. *J Endocrinol Invest* 2012;35:219-23.
 55. La Vignera S, Condorelli R, Vicari E, D'Agata R, Calogero AE. Effects of varicocele on sperm DNA fragmentation, mitochondrial function, chromatin condensation, and apoptosis. *J Androl* 2012;33:389-96.
 56. Condorelli R, Calogero AE, La Vignera S. Relationship between testicular volume and conventional or nonconventional sperm parameters. *Int J Endocrinol* 2013;2013:145792.
 57. Marchiani S, Tamburrino L, Benini F, Fanfani L, Dolce R, Rastrelli G, et al. Chromatin protamination and catsper expression in spermatozoa predict clinical outcomes after assisted reproduction programs. *Sci Rep* 2017;7:15122.
 58. Usachenko SI, Bradbury EM. Histone-DNA contacts in structure/function relationships of nucleosomes as revealed by crosslinking. *Genetica* 1999;106:103-15.
 59. Fournier C, Labrune E, Lornage J, Soignon G, Giscard d'Estaing S, Guérin JF, et al. The impact of histones linked to sperm chromatin on embryo development and ART outcome. *Andrology* 2018;6:436-45.
 60. Dadoune JP. Spermatozoal RNAs: what about their functions? *Microsc Res Tech* 2009;72:536-51.
 61. Zhao Y, Li Q, Yao C, Wang Z, Zhou Y, Wang Y, et al. Characterization and quantification of mRNA transcripts in ejaculated spermatozoa of fertile men by serial analysis of gene expression. *Hum Reprod* 2006;21:1583-90.
 62. Curry E, Safranski TJ, Pratt SL. Differential expression of porcine sperm microRNAs and their association with sperm morphology and motility. *Theriogenology* 2011;76:1532-9.
 63. Yan W, Morozumi K, Zhang J, Ro S, Park C, Yanagimachi R. Birth of mice after intracytoplasmic injection of single purified sperm nuclei and detection of messenger RNAs and microRNAs in the sperm nuclei. *Biol Reprod* 2008;78:896-902.
 64. Erickson RP. Post-meiotic gene expression. *Trends Genet* 1990;6:264-9.
 65. Vibranovski MD, Chalopin DS, Lopes HF, Long M, Karr TL. Direct evidence for postmeiotic transcription during *Drosophila melanogaster* spermatogenesis. *Genetics* 2010;186:431-3.
 66. Fischer BE, Wasbrough E, Meadows LA, Randlet O, Dorus S, Karr TL, et al. Conserved properties of *Drosophila* and human spermatozoal mRNA repertoires. *Proc Biol Sci* 2012;279:2636-44.
 67. Bourc'his D, Voinnet O. A small-RNA perspective on gametogenesis, fertilization, and early zygotic development. *Science* 2010;330:617-22.
 68. Hosken DJ, Hodgson DJ. Why do sperm carry RNA? Relatedness, conflict, and control. *Trends Ecol Evol* 2014;29:451-5.
 69. Miller D, Ostermeier GC, Krawetz SA. The controversy, potential and roles of spermatozoal RNA. *Trends Mol Med* 2005;11:156-63.

70. Boerke A, Dieleman SJ, Gadella BM. A possible role for sperm RNA in early embryo development. *Theriogenology* 2007;68 Suppl 1:S147-55.
71. Donkin I, Versteyhe S, Ingerslev LR, Qian K, Mechta M, Nordkap L, et al. Obesity and bariatric surgery drive epigenetic variation of spermatozoa in humans. *Cell Metab* 2016;23:369-78.
72. Xiao J, Wang X, Luo Y, Li XK, Li XW. Research progress in sRNAs and functional proteins in epididymosomes. *Yi Chuan* 2018;40:197-206.
73. Conine CC, Sun F, Song L, Rivera-Pérez JA, Rando OJ. Small RNAs gained during epididymal transit of sperm are essential for embryonic development in mice. *Dev Cell* 2018;46:470-80.e3.
74. Siomi MC, Sato K, Pezic D, Aravin AA. PIWI-interacting small RNAs: the vanguard of genome defence. *Nat Rev Mol Cell Biol* 2011;12:246-58.
75. Gou LT, Kang JY, Dai P, Wang X, Li F, Zhao S, et al. Ubiquitination-deficient mutations in human piwi cause male infertility by impairing histone-to-protamine exchange during spermiogenesis. *Cell* 2017;169:1090-104.e13.
76. Guo L, Chao SB, Xiao L, Wang ZB, Meng TG, Li YY, et al. Sperm-carried RNAs play critical roles in mouse embryonic development. *Oncotarget* 2017;8:67394-405.