

Detection and Typing of HSV-1, HSV-2, CMV and EBV by Quadruplex PCR

Chang Ho Shin, Gun Suk Park, Kyeong Man Hong, and Moon Kee Paik

Department of Biochemistry, Wonkwang University College of Medicine, Iksan, Korea.

The development of a multiplex polymerase chain reaction (PCR) method for rapid and accurate detection and typing of herpes simplex virus type 1 (HSV-1), and type-2 (HSV-2), cytomegalovirus (CMV) and Epstein-Barr virus (EBV) is very important for clinical diagnosis to allow the deliver of therapy as early as possible. Large scale amplifications by multiplex PCR of viral DNA can lower the cost and time for viral diagnosis. In this study, therefore sensitive quadruplex PCR was achieved by optimizing parameters such as primers, and 1.5 mM magnesium and 200 μ M dNTPs concentrations. The concentrations of HSV-1, HSV-2, CMV and EBV primers were 0.5, 0.3, 0.25 and 0.25 pmoles, respectively. Optimal annealing temperature was 54°C. Employing these conditions, we could detect 10 copies of reconstructed template plasmid DNA, which were cloned to vectors containing target sequences of viral DNA. PCR products of 271 bp for HSV-1, 231 bp for HSV-2, 368 bp for CMV, and 326 bp for EBV were separated on 5.0% polyacrylamide gel electrophoresis and confirmed by direct sequencing.

The present study showed that the quadruplex PCR assay described herein has potential application in clinical diagnosis, when rapid, accurate detection and typing of viruses HSV-1, HSV-2, CMV or EBV are necessary.

Key Words: Diagnosis, multiplex PCR, virus

INTRODUCTION

Several human virus infections play an important role in human diseases. Herpes simplex virus

type-1 (HSV-1), and type-2 (HSV-2), cytomegalovirus (CMV) and Epstein-Barr virus (EBV) are widespread human viruses frequently detected in specific tumors and immunosuppressed individuals. HSV-1 and HSV-2 cause meningitis, fatal encephalitis and other ocular diseases such as uveitis, keratitis, retinitis and acute retinal necrosis syndrome, and result in a significant mortality in neonates. Especially, CMV and HSV-2 have been demonstrated to display an oncogenic potential *in vitro*.¹⁻⁴ There have been numerous reports linking CMV and EBV to tumors, lymphoepithelial carcinoma, smooth muscle tumors and esophageal carcinoma, and suggesting that these viruses transactivate cellular oncogenes such as *fos*, *jun* and *myb*.⁵⁻⁷ As they are also associated with many diseases, primary diagnosis of the latency period is important for therapy.

In the past, antibodies were developed for their diagnosis in many clinical pathology laboratories. Diagnosis of these viruses is clinically difficult and usually depends on the isolation of the virus by tissue culture methods, by which some viruses grow slowly for one to several weeks.^{8,9} Many authors have reported diagnostic methods to distinguish between HSV types which have a very similar DNA structure. HSV-1 and HSV-2 have been distinguished by various techniques including their different sensitivity to (ϵ)-5-(2-bromovinyl)-2'-deoxy-thymidine (BVDU), differential replication in chick embryo cells versus guinea pig embryo cells,¹⁰ restriction endonuclease,¹⁰⁻¹² recognition by polyclonal or monoclonal antibodies^{10,12,13} including immunofluorescence assay methods,⁸ detection of viral nucleic acid by DNA hybridization,^{14,15} and finally polymerase chain reaction (PCR).

Serologic data, frequently used for diagnosis of

Received June 23, 2003

Accepted October 10, 2003

This work was supported by a grant from Wonkwang University in 2001.

Reprint address: requests to Dr. Moon Kee Paik, Department of Biochemistry, Wonkwang University College of Medicine, 344-2 Shinyong-dong, Iksan, Chonbuk 570-711, Korea. Tel: 82-63-850-6795, Fax: 82-63-854-1459, E-mail: forensic@hammail.net

viral infections, represent only indirect markers of infection and are often unreliable in immunocompromised patients, whereas PCR can detect only the presence of viral genomes.^{2,16-18} Because antiviral therapy is effective in the early stage of disease processing, the diagnosis of EBV, CMV, HSV-1 and HSV-2 (ECH12) has been improved by the (PCR) technology,^{19,20} which allows rapid and sensitive detection of herpes viruses. The PCR method has replaced serological assays which are based on the detection of virus-specific IgM antibodies. Furthermore, multiplex PCR can save time and cost by simultaneously amplifying multiple sequences in a single reaction.²¹ An efficient multiplex PCR usually requires strategic planning and multiple attempts to optimize reaction conditions. In many cases, in order to obtain optimal multiplex PCR condition, an optimal single PCR condition has to be first established. In such a situation, the PCR primer pairs will retain target specificity. In the present study, we described the development of a quadruplex PCR method for the detection of HSV-1, HSV-2, EBV and CMV (ECH12) in a single step reaction.

MATERIALS AND METHODS

Samples

Infected samples of HSV-1, HSV-2 and CMV (ECH12) were obtained from Seoul Clinical Laboratory (SCL), and EBV was from the Department of Clinical Pathology, Wonkwang University Medical College. These samples were used as positive controls. Genomic DNA, used as negative control, was extracted from peripheral blood by Tris buffered phenol/chloroform and precipitated with ethanol.

PCR amplification

The primers used for HSV-1 and HSV-2 PCR corresponded to the glycoprotein D coding sequence, which was the major target of vaccination studies for highly immunogenic and inducing neutralizing antibodies. The PCR primer sequences for detection of CMV and EBV were designed with the coding sequences of the structural proteins of virus, UL56 and late membrane

protein 1 (LMP-1), respectively. The HSV-1 primer sequences were 5'-CGTACCTGCGGCTCGTGAA GT-3' as the forward and 5'-AGCAGGGTGCTC GTGTATGGGC-3' as the reverse; 5'-TGGTATCG CATGGGAGACAAT-3' for the forward and 5'-CTCCGTCCAGTCGTTTATCTTG-3' for the reverse primer of HSV-2; 5'-ACGTGTTACTGGCG GAGTCG-3' for the forward and 5'-TTGAGTG TGGCCAGACTGAG-3' for the reverse primer of CMV; 5'-AGCACTGGCCAGCTCATATC-3' for the forward and 5'-TTGACGTCATGCCAAGGC AA-3' for the reverse of EBV. The expected sizes of the amplified sequences in the HSV-1, HSV-2, CMV and EBV genomes were 271, 231, 368 and 326 base pairs, respectively. Single PCR was undertaken in a final 20 μ l volume containing 1.0 U *Taq* polymerase, 1X polymerase buffer [containing 50 mM KCl, 10 mM Tris-HCl (pH 8.3)], 1.5 mM MgCl₂, 200 μ M dNTPs and 0.5 pmole primers using a GeneAmp PCR System 2400 (Applied Biosystems, Foster, California, USA). PCR amplification conditions were, predenaturation step at 95°C for 3 min., followed by denaturation at 95°C for 30 sec., annealing at 54 °C for 30 sec. and primer extension at 72°C for 30 sec. The final extension was performed for another 7 min, repeating 45 cycles of amplification.

Detection of amplified products

The amplified PCR products were electrophoresed on 2.0% agarose gel and visualized with UV light after staining with ethidium bromide. Amplification products were also separated by electrophoresis through a 0.4 mm to 1.2 mm discontinuous gel and 5% T-5% C denaturing polyacrylamide gel containing 1 X TBE and 7M urea.

Five microliters of each amplified DNA sample were mixed with 3 μ l of loading solution (2 mM EDTA, 95% formamide, 0.05% bromophenol blue, 0.05% xylene cyanol), denatured at 95°C for 3 min, and chilled on ice prior to loading. Electrophoresis was performed at 55 W for 140 min with the SA32 electrophoresis apparatus (Gibco BRL, Gaithersburg, Maryland, USA).

Silver staining

After electrophoresis, the gel was washed with

triple distilled water for 1 min, and immersed in 1% nitric acid for 5 min. The oxidized gel was washed with triple distilled water for 1 min prior to treatment with 0.12 M silver nitrate solution for 20 min. The washed gel in distilled water was developed with 0.28 M anhydrous sodium carbonate and 0.19% formalin solution. The stained gel on 3 MM Whatman paper was dried at 80°C for 1 hr with a vacuum gel dryer (Amersham Biosciences, Uppsala, Sweden).

Plasmid construction of ECH12 DNA

In order to calculate the amount of genomic DNA from PCR amplifications, PCR amplified products of 271, 231, 368 and 326 bps from the control strain were inserted into multicloning sites of 3015 bps pGEM-T Easy vector (Promega, Madison, Wisconsin, USA) plasmid DNA. Plasmid DNA from cultured bacteria was extracted and purified as described by Sambrook et al.²² The original number of each ECH12 copys was determined by optical density (OD) against the molecular weights of different dilutions of these plasmid DNAs, and the copy numbers of ECH12 DNA were calculated to be 2.724×10^{11} molecules/ $1.0 \mu\text{g}/\text{uL}$, 2.712×10^{11} molecules/ $1.0 \mu\text{g}/\text{uL}$, 2.772×10^{11} molecules/ $1.0 \mu\text{g}/\text{uL}$ and 2.802×10^{11} molecules/ $1.0 \mu\text{g}/\text{uL}$, respectively.

Sequencing of positive control DNA

For confirmation of sequences, fragments of each virus amplified through single PCR were resolved on agarose gel. After electrophoresis, the DNA band of interest was excised from the gel and eluted with a QIAquick Gel Extraction Kit (Qiagen, Hilden, Germany). About 50 ng of eluted amplified DNA were used as a template for direct sequencing. Sequencing was performed with ABI PRISM BigDyeTM Terminator Cycle Sequencing Ready Reaction Kits (Applied Biosystems) using an ABI PRISMTM 310 Genetic Analyzer (Applied Biosystems, Foster, California, USA).

Multiplex PCR

Multiplex PCR was also performed with Gene Amp PCR System 2400 (Applied Biosystems,

Foster, California, USA). Preheating for 3 min. at 95°C, followed by 45 amplification cycles of 30 sec at 94°C, 10 sec at 54°C, and 30 sec at 72°C were carried out in a final 20 uL volume containing 2 uL of GeneAmp 10X PCR Buffer (PE Biosystems, USA), 0.2 mM each of deoxynucleotide triphosphates (dNTP), 1 unit of *Taq* polymerase (Solgent, Korea) and 1.5 mM MgCl₂. The control contained 5 uL of DNA sample (positive control of ECH12 DNA or uninfected genomic DNA) which was added to the reaction mixture. After the last cycle, the samples were incubated for 7 min at 72°C to complete the extension of the primers. To optimize the buffer concentration of multiplex PCR, we performed a series of reactions to determine the optimal primer and Mg²⁺ concentrations and dNTP levels. Five microliters of each amplified products were analyzed on 2.0% agarose gel or 5% T-5% C denaturing acrylamide gel containing 7M urea.

Specificity and sensitivity

The specificity of the PCR reaction was examined by varying the cycle numbers with 1 ng of mixed positive control viral DNA which had been inserted into the plasmid vector. Also, variable annealing temperatures and times for each single PCR and quadruplex PCR were examined. To evaluate the PCR sensitivity, a decreasing number of plasmid DNA copies, ranging from 10⁸ copies to 1, was amplified. The specificity and sensitivity were examined for each primer pair and also for the combined four primer pairs in the multiplex PCR.

RESULTS

Optimization of a single PCR

PCR was carried out under various conditions which affect amplification, such as cycle numbers, MgCl₂ and dNTPs concentrations, annealing temperatures, and time. The PCR reactions were carried out for various cycle numbers, ranging from 15 to 50 cycles, at increments of 5 cycles, and for various temperatures, ranging from 54°C to 60°C, at increments of 2°C. EBV, CMV, HSV-1 and




Fig. 1. Electrophoresis-analysis on agarose gel of single PCR products obtained with HSV-1 (A), HSV-2 (B), CMV (C) and EBV (D) primers. The upper panel shows amplified products which were 271 bp and 231 bp of HSV-1 and HSV-2, respectively. The lower panel shows 368 bp and 326 bp of CMV and EBV, respectively. Lane 1 was carried with 15 cycles, at increments of 5 cycles. M; 100 bp ladder.




Fig. 2. Amplification as a function of annealing temperatures and time. PCR reactions of lanes 1 were performed at 54°C for 30 sec. Annealing temperature was increased two degrees at each reaction and annealing times were 30 sec and 60 sec at a same degree. Upper panel; HSV-1 (A) and HSV-2 (B), Lower panel; CMV (C) and EBV (D), M; 100 bp ladder.

HSV-2 were amplified to the sizes of 326 bp, 368 bp, 271 bp and 231 bp, respectively (Fig. 1 and 2).

Optimization of multiplex PCR

In order to obtain the optimal condition for multiplex PCR reactions, different dNTPs and MgCl₂ concentrations were examined. Since MgCl₂ and dNTPs concentrations are dependent on each other and also on the primer concentration, multiplex PCR reactions were performed with template plasmids of 10⁸, 10³, 10² and 10 copies, and in 1.5 mM, 2.0 mM and 2.5 mM MgCl₂ (Fig. 3). All of the amplified PCR products of ECH12 also included




Fig. 3. Effect of MgCl₂ and template concentrations on multiplex PCR. The PCR products obtained with HSV-1, HSV-2, CMV and EBV primers were electrophoresed on 2.0% agarose gel (panel A) and polyacrylamide gel (panel B) under the various concentrations of MgCl₂ and template. N; negative control (10 ng of uninfected genomic DNA), lanes 1-4; 1.5 mM MgCl₂, lanes 5-8; 2.0 mM MgCl₂, lanes 9-12; 2.5 mM MgCl₂, lanes 1, 5, 9; 10⁸ copies of template DNA, lanes 2, 6, 10; 10³ copies of template DNA, lanes 3, 7, 11; 10² copies of template DNA, lanes 4, 8, 12; 10 copies of template DNA. Arrows indicate amplified fragments, 231 bp, 271 bp, 326 bp and 368 bp. M; 100 bp ladder.

unexpected DNA under the condition of each MgCl₂ concentration, containing 10⁸ and 10⁵ copies of template DNA with silver stained gel (Fig. 3B, lanes 1, 2, 5, 6, 9, 10). With 10³ copies of template DNA at 1.5 mM and 2.5 mM MgCl₂, four kinds of amplified fragments of 231bp, 271 bp, 326 bp and 368 bp were detected (Fig. 3, lanes 2, 10), whereas 326 bp and 368 bp fragments were not detected with 2.0 mM MgCl₂ (Fig. 3, lanes 2, 6, 10). However, at all the concentrations of MgCl₂ and template plasmid tested, the fragments of HSV-1 and HSV-2 were amplified and detected (Fig. 3). When these PCR products were also electrophoresed on 5% polyacrylamide gel followed by silver staining, the results clearly indicated artifacts and a higher sensitivity than that performed on agarose gel (Fig. 3B). The best results were obtained with 1.5 mM MgCl₂ for the amplification of ECH12, therefore, this level was used to determine the optimal concentration of dNTPs. Thus, at the fixed concentration of 1.5 mM MgCl₂, multiplex PCR was performed with different concentrations of dNTPs (200 μM, 300 μM and 400 μM). As shown in Fig. 4, DNA patterns of amplified fragments were detected with 10² copies of template plasmid. How-




Fig. 4. Effect of dNTPs and template concentrations. Agarose gel (panel A) and acrylamide gel (panel B) analysis of multiplex PCR products obtained with HSV-1, HSV-2, CMV and EBV primers. N; negative control (10 ng of uninfected genomic DNA), lanes 1-4; 200 μ M dNTPs, lanes 5-8; 300 μ M dNTPs, lanes 9-12; 400 μ M dNTPs, lanes 1, 5, 9; 10^8 copies of template DNA, lanes 2, 6, 10; 10^3 copies of template DNA, lanes 3, 7, 11; 102 copies of template DNA, lanes 4, 8, 12; 10 copies of template DNA. Arrows indicate amplified fragments, 231 bp, 271 bp, 326 bp and 368 bp. M; 100 bp ladder.

ever, different concentrations of dNTPs, including 200 μ M, 300 μ M and 400 μ M, showed no difference. On the other hand, these fragments of 10-copy concentration were easily detectable on silver stained gel (Fig. 4B). These results, therefore, indicated that a 200 μ M concentration of dNTPs was sufficient to amplify 10^8 copies of template plasmid.

Specificity and sensitivity

In order to establish the lowest possible detection limit for all four primer pairs of ECH12 in the multiplex amplification, amplification was performed with serial dilutions, ranging from 10^8 to 1 copy, of template under the optimal $MgCl_2$ and dNTPs concentrations (Fig. 5). Also, 10 ng of the genomic DNA obtained from a human who was not infected with these viruses were used as a negative DNA control (Fig. 5, lane 1). Amplified PCR products of ECH12 were detectable with 10^2 copies of template plasmid, and EBV could not be detected as a clear amplification band at a concentration lower than 10^2 . On the other hand, HSV-1 and HSV-2 were detected with this copy concentration, and artifact DNA bands were




Fig. 5. Effect of copy numbers of template DNA on multiplex PCR. Multiplex PCR was performed with 1.5 mM $MgCl_2$ and 2.0 μ M dNTPs. Agarose gel (panel A) and acrylamide gel (panel B) analysis of multiplex PCR products obtained with HSV-1, HSV-2, CMV and EBV primers. N; negative control (100 ng of uninfected genomic DNA), lanes 1-9; 10^8 -1 copies of template DNA. Arrows indicate amplified fragments, 231 bp, 271 bp, 326 bp and 368 bp. M; 100 bp ladder.

observed with 10^5 copy template plasmids (Fig. 5B). Indeed, PCR products of HSV-1 and HSV-2 were detected with 1 copy template plasmid on agarose gel. Four kinds of predicted fragments of ECH12 with 10 copies were detected on silver stained polyacrylamide gel (Fig. 5B, lane 8), but not on agarose gel (Fig. 5A, lane 8).

DISCUSSION

Many molecular biological data suggest that certain types of some members of the *herpesviridae* family such as Epstein-Barr virus, human herpes simplex virus type 2 and cytomegalovirus may play an oncogenic role¹ and also induce neurological syndromes such as aseptic meningitis, polyradiculitis and acute encephalitis. Furthermore, these viruses are ubiquitously distributed, causing a wide range of acute central nerve system (CNS) diseases either during primary infection or in the course of virus reactivation or reinfection. Detection of viral DNA by PCR has been shown to be a sensitive and specific method for rapid diagnosis.¹⁶

We herein described a rapid and accurate multiplex PCR method for the detection and typing of HSV-1, HSV-2, CMV and EBV. In a multiplex

PCR, several parameters, which may affect the structure of primers and affinity to templates DNA, should be considered.²¹ Multiplex PCR includes the problems of interference between the different primers, which can lead to nonspecific amplimers, and also coordinate amplification of the bands.²³ A sensitive assay condition was established by optimizing the concentrations of dNTPs and MgCl₂ that are necessary for elongation. In each single PCR, one-half unit of *Taq* polymerase was used and the amount of newly synthesized DNA fragments was found to be propositional to different cycling numbers. However, 50 PCR cycles were carried out in the present study (Fig. 1). The primer pairs of this study were found to be independent of annealing temperature and time. With different ranges of annealing temperature and time, PCR amplified fragments were visualized with ethidium bromide staining under UV light on 2.0% agarose gel. Nonspecific bands of about 550 bp were detected in HSV-1 reactions (Fig. 2). Nonspecific bands of the same size were also detected among the multiplex PCR fragments which were performed at higher template concentrations (Fig. 2 and 3). It was highly possible that these fragments might have been produced by primers of HSV-1, because these fragments were also detectable among the products of single PCR. And with multiplex PCR using a higher concentration of template DNA (10⁵ copies), artifact bands at sizes of about 260 bp, 264 bp and 310 bp, which were not separated on agarose gel, were also detected by silver stained gel (Fig. 3B, 4B and 5B). These results strongly indicate that diagnosis using multiplex PCR methods should be carried out by polyacrylamide gel and silver staining, because rapid and accurate diagnosis is the most important factor for patients' therapy, and that their diagnosis by single PCR may be delayed. Our results indicated that multiplex PCR was successful with 10 copies of template and did not produce any amplified DNA fragments in the negative control (10 ng of genomic DNA), when agarose gel and silver stained polyacrylamide gel were used.

In the present study, optimal conditions of multiplex PCR were established, and amplified fragments were detected with high sensitivity by denaturing polyacrylamide gel. Quadruplex PCR

was capable of clinically diagnosing HSV-1, HSV-2, CMV and EBV in one PCR reaction.

ACKNOWLEDGEMENTS

We thank Dr. Woon Ki Paik for proof reading of the manuscript. The virus samples were generously provided by Dr. Young Suk Park, Seoul Clinical Laboratory.

REFERENCES

1. Koffa M, Koumantakis E, Ergazaki M, Tsatsanis C, Spandidos DA. Association of herpesvirus infection with the development of genital cancer. *Int J Cancer* 1995;63:58-62.
2. Nogueira ML, Siqueira RC, Freitas N, Amorim JB, Bonjardim CA, Ferreira PC, et al. Detection of herpesvirus DNA by the polymerase chain reaction (PCR) in vitreous samples from patients with necrotising retinitis. *J Clin Pathol* 2001;54:103-6.
3. Read SJ, Kurtz JB. Laboratory diagnosis of common viral infections of the central nervous system by using a single multiplex PCR screening assay. *J Clin Microbiol* 1999;37:1352-5.
4. Yamamoto S, Pavan-Langston D, Kinoshita S, Nishida K, Shimomura Y, Tano Y. Detecting herpesvirus DNA in uveitis using the polymerase chain reaction. *Br J Ophthalmol* 1996;80:465-8.
5. Boldogh I, AbuBakar S, Deng CZ, Albrecht T. Transcriptional activation of cellular oncogenes fos, jun, and myc by human cytomegalovirus. *J Virol* 1991;65:1568-71.
6. Chang F, Syrjanen S, Shen Q, Cintorino M, Santopietro R, Tosi P, et al. Evaluation of HPV, CMV, HSV and EBV in esophageal squamous cell carcinomas from a high-incidence area of China. *Anticancer Res* 2000;20:3935-40.
7. Weiss LM, Movahed LA, Warnke RA, Sklar J. Detection of Epstein-Barr viral genomes in Reed-Sternberg cells of Hodgkin's disease. *N Engl J Med* 1989;23:502-6.
8. Gleaves CA, Wilson DJ, Wold AD, Smith TF. Detection and serotyping of herpes simplex virus in MRC-5 cells by use of centrifugation and monoclonal antibodies 16 h postinoculation. *J Clin Microbiol* 1985;21:29-32.
9. Hashido M, Lee FK, Inouye S, Kawana T. Detection of herpes simplex virus type-specific antibodies by an enzyme-linked immunosorbent assay based on glycoprotein G. *J Med Virol* 1997;53:319-23.
10. Mayo D, Hilliard J, Landry M, Egbertson S, Mirkovic R. Comparison of four methods for typing low-passage herpes simplex virus isolates. *Intervirology* 1985;23:112-5.
11. Lonsdale DM. A rapid technique for distinguishing herpes-simplex virus type 1 from type 2 by restriction-

- enzyme technology. *Lancet* 1979;21:849-52.
12. Peterson E, Schmidt OW, Goldstein LC, Nowinski RC, Corey L. Typing of clinical herpes simplex virus isolates with mouse monoclonal antibodies to herpes simplex virus types 1 and 2: comparison with type-specific rabbit antisera and restriction endonuclease analysis of viral DNA. *J Clin Microbiol* 1983;17:92-6.
 13. Lipson SM, Schutzbank TE, Szabo K. Evaluation of three immunofluorescence assays for culture confirmation and typing of herpes simplex virus. *J Clin Microbiol* 1987;25:391-4.
 14. Kimpton CP, Corbitt G, Morris DJ. Detection of cytomegalovirus by dot-blot DNA hybridization using probes labelled with 32P by nick translation or random hexanucleotide priming. *Mol Cell Probes* 1988;2:181-8.
 15. Swierkosz EM, Scholl DR, Brown JL, Jollick JD, Gleaves CA. Improved DNA hybridization method for detection of acyclovir-resistant herpes simplex virus. *Antimicrob Agents Chemother* 1987;31:1465-9.
 16. Cinque P, Scarpellini P, Vago L, Linde A, Lazzarin A. Diagnosis of central nervous system complications in HIV-infected patients: cerebrospinal fluid analysis by the polymerase chain reaction. *AIDS* 1997;11:1-17.
 17. Temcharoen S, Hirsch P, Theamboonlers A, Likitnukul S, Seksarn P, Poovorawan Y. Cytomegalovirus infection in immunocompromised children in Thailand. *Asian Pac J Allergy Immunol* 1999;17:295-9.
 18. Webber SA, Wilson NJ, Junker AK, Byrne SK, Perry A, Thomas EE, et al. Postpericardiotomy syndrome: no evidence for a viral etiology. *Cardiol Young* 2001;11: 67-74.
 19. Linde A, Klapper PE, Monteyne P, Echevarria JM, Cinque P, Rozenberg F, et al. Specific diagnostic methods for herpesvirus infections of the central nervous system: a consensus review by the European Union Concerted Action on Virus Meningitis and Encephalitis. *Clin Diagn Virol* 1997;8:83-104. Review.
 20. Puchhammer-Stockl E. Resistance--clinical significance and diagnostics. *Wien Med Wochenschr* 1998;148:556-9.
 21. Repp R, Rhiel S, Heermann KH, Schaefer S, Keller C, Ndumbe P, et al. Genotyping by multiplex polymerase chain reaction for detection of endemic hepatitis B virus transmission. *J Clin Microbiol* 1993;31:1095-102.
 22. Sambrook J, Fritsch EF, Maniatis T, eds. *Molecular cloning. A laboratory manual*, 2nd ed. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory; 1989. p.9.16-9.19.
 23. Markoulatos P, Samara V, Siafakas N, Plakokefalos E, Spyrou N, Moncany ML. Development of a quadruplex polymerase chain reaction for human cytomegalovirus detection. *J Clin Lab Anal* 1999;13:99-105.