

## Erratum to: Effect of High Dose Rosuvastatin Loading before Primary Percutaneous Coronary Intervention on Infarct Size in Patients with ST-Segment Elevation Myocardial Infarction

Ji Won Kim, MD, Kyeong Ho Yun, MD, Eun Kyung Kim, MD, Yong Cheol Kim, MD, Dai-Yeol Joe, MD, Jum Suk Ko, MD, Sang Jae Rhee, MD, Eun Mi Lee, MD, Nam Jin Yoo, MD, Nam-Ho Kim, MD, Seok Kyu Oh, MD, and Jin-Won Jeong, MD

*Department of Cardiovascular Medicine, Regional Cardiocerebrovascular Center, Wonkwang University Hospital, Iksan, Korea*

Korean Circ J 2014;44(2):76-81

<http://dx.doi.org/10.4070/kcj.2014.44.2.76>

On page 76, reference no. 21 was inadvertently omitted.

21. Pellikka PA, Behrenbeck T, Verani MS, Mahmarian JJ, Wackers FJ, Gibbons RJ. Serial changes in myocardial perfusion using tomographic technetium-99m-hexamethylpropylene-2-methoxy-isobutylisonitrile imaging following reperfusion therapy of myocardial infarction. *J Nucl Med* 1990;31:1269-75.

The Publishing Company apologizes for these errors and any inconvenience they may have caused.