

Editorial



Elucidation of the Diagnosis and Treatment of Peripheral Arterial Disease

Hyung Oh Kim , MD, and Weon Kim , MD, PhD

Division of Cardiovascular, Department of Internal Medicine, Kyung Hee University Hospital, Kyung Hee University, Seoul, Korea

OPEN ACCESS

Received: May 4, 2018

Accepted: May 31, 2018

Correspondence to

Weon Kim, MD, PhD

Division of Cardiovascular, Department of Internal Medicine, Kyung Hee University Hospital, Kyung Hee University, 23, Kyungheedaero-ro, Dongdaemun-gu, Seoul 02447, Korea.

E-mail: mylovekw@hanmail.net

Copyright © 2018. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Hyung Oh Kim

<https://orcid.org/0000-0003-2157-441X>

Weon Kim

<https://orcid.org/0000-0003-1264-9870>

Conflict of Interest

The authors have no financial conflicts of interest.

Author Contributions

Investigation: Kim HO; Resources: Kim HO, Kim W; Supervision: Kim W; Visualization: Kim W; Writing - original draft: Kim HO; Writing - review & editing: Kim W.

► See the article “Medical Resource Consumption and Quality of Life in Peripheral Arterial Disease in Korea: PAD Outcomes (PADO) Research” in volume 48 page on 813.

Medical treatment is an important management axis of peripheral arterial disease (PAD), as the disease manifests as advanced systemic atherosclerosis. PAD is considered a coronary artery disease (CAD) risk equivalent.¹⁾ Moreover, all-cause mortality is closely linked to the occurrence and severity of PAD, reaching 20% annually in patients with limb-threatening manifestations. Management for PAD patients must include the optimal medical treatment for reducing the risk of cardiovascular complications, including cardiovascular death, myocardial infarction (MI), and stroke.²⁾ Although the disease is a serious burden on patients' health, registry-based studies about the current situations of diagnosis and treatment of PAD from the Korean Medical Society are limited,^{3,4)} unlike international studies.^{5,6)} In this issue of *Korean Circulation Journal*, Rha et al.⁷⁾ reported the analysis of multicenter, large-scale data, which may have considerable implications, which include the following.

First, Rha et al.⁷⁾ reported important facts about the current treatment situation for PAD. The authors reported that 94.1% of patients were receiving pharmacotherapy, mostly with antiplatelets, but reported a smoking cessation education rate of 12.5% and exercise education rate of 23.8%. These results could imply that the non-pharmacological approach is being neglected in many patients, even though the current guideline recommends it as a class I or IIa recommendation for smoking cessation and exercise.⁸⁾ Second, most patients had visited non-internal medical hospitals. In some respects, visiting orthopedics and general clinics in advance of visiting tertiary medical centers is unavoidable among patients. Thus, screening in primary clinics becomes important. The simple ankle-brachial index (ABI) is known to be a tool for diagnosing many patients with an unrecognized PAD.⁶⁾ The authors reported that the mean number of previous visits before diagnosis and treatment is more than 1. Education and promotion of appropriate screening tests such as the ABI to non-cardiologists may shorten the time and reduce the cost of diagnosis of unrecognized PAD. Especially considering cost-effectiveness, ABI screening in primary clinics may be greatly useful for identifying those patients. Third, the quality of life (QoL) scores statistically improved after 6 months' follow-up. Considering that exercise, diet, and smoking cessation only accounted for only a small portion of the treatment modality, the major treatment effect might be from pharmacotherapy and revascularization. As the authors mentioned, QoL is reported to be a prognostic factor of patients with PAD.⁹⁾ If QoL could be improved by medication and percutaneous intervention, it must be conducted actively by practitioners.

The contents of the report are the author's own views and do not necessarily reflect the views of the *Korean Circulation Journal*.

Although Rha et al.⁷⁾ pointed several important results from large-scale data, the article has some limitations. First, the data used were based on a multicenter-registry database, and hospital-derived databases embed an exclusion of patients outside tertiary hospitals. Second, the report suggests symptom characteristics at baseline but still lacks symptom profiles in the treatment investigation. A treatment pattern analysis may provide more information with classification of the presence of symptoms, considering that symptoms may affect the treatment modality subtly.

The study by Rha et al.⁷⁾ informs practitioners about the current practice among patients before and after visiting a tertiary hospital. The importance of a guideline-based treatment and an appropriate screening test must be emphasized, in addition to non-pharmacological treatments. In the academic aspect, this study may serve as a basis and trigger for future prospective randomized studies by the Korean Medical Society.

REFERENCES

- Berger JS, Krantz MJ, Kittelson JM, Hiatt WR. Aspirin for the prevention of cardiovascular events in patients with peripheral artery disease: a meta-analysis of randomized trials. *JAMA* 2009;301:1909-19.
[PUBMED](#) | [CROSSREF](#)
- Rooke TW, Hirsch AT, Misra S, et al. Management of patients with peripheral artery disease (compilation of 2005 and 2011 ACCF/AHA Guideline Recommendations): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013;61:1555-70.
[PUBMED](#) | [CROSSREF](#)
- Ko YG, Ahn CM, Min PK, et al. Baseline characteristics of a retrospective patient cohort in the Korean Vascular Intervention Society Endovascular Therapy in Lower Limb Artery Diseases (K-VIS ELLA) registry. *Korean Circ J* 2017;47:469-76.
[PUBMED](#) | [CROSSREF](#)
- Kim HO, Kim JM, Woo JS, et al. Effects of chronic kidney disease on clinical outcomes in patients with peripheral artery disease undergoing endovascular treatment: analysis from the K-VIS ELLA registry. *Int J Cardiol* 2018;262:32-7.
[PUBMED](#) | [CROSSREF](#)
- Bhatt DL, Steg PG, Ohman EM, et al. International prevalence, recognition, and treatment of cardiovascular risk factors in outpatients with atherothrombosis. *JAMA* 2006;295:180-9.
[PUBMED](#) | [CROSSREF](#)
- Hirsch AT, Criqui MH, Treat-Jacobson D, et al. Peripheral arterial disease detection, awareness, and treatment in primary care. *JAMA* 2001;286:1317-24.
[PUBMED](#) | [CROSSREF](#)
- Rha SW, Choi SH, Kim DI, et al. Medical resource consumption and quality of life in peripheral arterial disease in Korea: PAD Outcomes (PADO) research. *Korean Circ J* 2018;48:813-25.
[CROSSREF](#)
- Writing Committee Members Gerhard-Herman MD, Gornik HL, et al. 2016 AHA/ACC guideline on the management of patients with lower extremity peripheral artery disease: executive summary. *Vasc Med* 2017;22:NP1-43.
[PUBMED](#) | [CROSSREF](#)
- Issa SM, Hoeks SE, Scholte op Reimer WJ, et al. Health-related quality of life predicts long-term survival in patients with peripheral artery disease. *Vasc Med* 2010;15:163-9.
[PUBMED](#) | [CROSSREF](#)