

Review Article



Percutaneous Coronary Intervention and Coronary Artery Bypass Grafting for the Treatment of Left Main Coronary Artery Disease

Sang-Cheol Cho , MD, Duk-Woo Park , MD, PhD, and Seung-Jung Park , MD, PhD

Department of Cardiology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

OPEN ACCESS

Received: Apr 8, 2019

Accepted: Apr 10, 2019

Correspondence to

Seung-Jung Park, MD, PhD

Department of Cardiology, Asan Medical Center, University of Ulsan College of Medicine, 88, Olymplic-ro 43-gil, Songpa-gu, Seoul 05505, Korea.
E-mail: sjpark@amc.seoul.kr

Copyright © 2019. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Sang-Cheol Cho

<https://orcid.org/0000-0001-7587-5037>

Duk-Woo Park

<https://orcid.org/0000-0001-6643-0239>

Seung-Jung Park

<https://orcid.org/0000-0002-9187-5405>

Conflict of Interest

The authors have no financial conflicts of interest.

Author Contributions

Writing - original draft: Cho SC; Writing - review & editing: Cho SC, Park DW, Park SJ.

ABSTRACT

Severe stenosis of the left main coronary artery (LMCA) generally occurs as a result of atherosclerosis and compromises the blood supply to a wide area of myocardium, thereby increasing the risk of serious adverse cardiac events. Current revascularization strategies for patients with significant LMCA disease include coronary artery bypass grafting (CABG) and percutaneous coronary intervention (PCI), both of which have a range of advantages and disadvantages. In general, PCI is associated with a lower rate of periprocedural adverse events and provides more rapid recovery, while CABG provides more durable revascularization. Most clinical trials comparing PCI and CABG for the treatment of LMCA disease have shown PCI to be non-inferior to CABG with respect to mortality and the serious composite outcome of death, myocardial infarction, or stroke in patients with low-to-intermediate anatomical complexities. Remarkable advancements in PCI standards, including safer and more effective stents, adjunctive intravascular imaging or physiologic evaluation, and antithrombotic treatment, may have contributed to these favorable results. This review provides an update on the current management of LMCA disease with an emphasis on clinical data and academic and clinical knowledge that supports the use of PCI in an increasing proportion of patients with LMCA disease.

Keywords: Coronary artery disease; Percutaneous coronary intervention; Coronary artery bypass; Angioplasty, balloon, coronary; Drug-eluting stents; Treatment outcome

INTRODUCTION

Obstructive left main coronary artery (LMCA) disease is associated with a high rate of morbidity and mortality as a result of compromised myocardial blood supply; therefore, revascularization by coronary artery bypass grafting (CABG) surgery has been regarded as standard treatment. Over the past 20 years, there have been considerable therapeutic developments in the technique of percutaneous coronary intervention (PCI) for the treatment of obstructive coronary artery disease (CAD), involving improvements in stent technology, procedural techniques and refinement, periprocedural anticoagulation, concomitant antiplatelet agents, and cardiovascular medication.¹⁾²⁾

Several randomized clinical trials (RCTs) have been conducted to evaluate the potential therapeutic role of PCI as an alternative to standard CABG. With the introduction of first-generation drug-eluting stents (DESs), RCTs demonstrated that stenting achieved similar rates of mortality and hard clinical endpoints and a lower rate of stroke, although the rate of repeat revascularization was seen to be higher.³⁻⁶⁾ The development of second-generation DES was associated with improved efficacy and safety profiles compared with first-generation DES.^{7,8)} Subsequent RCTs were conducted and PCI has achieved greater clinical recognition as a reasonable therapeutic modality.^{9,10)} These data may impact on future clinical guidelines for myocardial revascularization and will ultimately will lead to greater use of PCI worldwide. Importantly, when undertaking PCI of the LMCA, there is increasing awareness of the need to achieve optimal procedural outcomes through the use of available technologies, including safer and more effective stents, intravascular imaging, and physiological assessment.

This review provides an update on the current management of LMCA disease with an emphasis on clinical data and procedural knowledge to support the use of PCI in a growing proportion of patients.

EVALUATION OF LMCA DISEASE

Contemporary evaluation of LMCA disease

Most patients with significant LMCA disease are symptomatic as a result of compromised blood supply to a large area of the myocardium. However, significant LMCA disease can be identified incidentally in stable patients undergoing coronary angiography. In the absence of significant LMCA stenosis or relevant clinical symptoms, the hemodynamic significance of incidental or intermediate LMCA lesions warrants further evaluation. Current clinical practice guidelines rely on angiographic lesion severity as the sole determinant of risk and for the de facto threshold for revascularization decision-making. However, this approach may be outdated in the current era of clinically proven noninvasive and invasive modalities and revascularization based solely on the angiographic appearance of LMCA stenosis of intermediate severity (50–70%) is not appropriate.¹¹⁾ In addition to the unwarranted surgical risk, premature CABG for potentially noncritical lesions may ultimately prove harmful to patients due to low graft patency rates and an accelerated rate of obstruction of bypassed native coronary vessels that may be up to 6-fold higher and which makes subsequent PCI technically challenging if required for symptom relief.¹²⁾ In the contemporary clinical setting, more detailed evaluation of the anatomic severity and hemodynamic significance of clinically ambiguous LMCA lesions can be obtained by intracoronary imaging with intravascular ultrasound (IVUS), or physiologically using pressure wire assessment of the fractional flow reserve (FFR).

IVUS characterizes the vessel size and the distribution of the plaques within the LMCA and its daughter branches, enabling accurate minimal lumen area (MLA) measurements at a cross-sectional level. Evaluation of IVUS in a prospective study showed that a MLA ≥ 6 mm² assessed by this technique is a safe value for deferring revascularization of the LMCA.¹³⁾ Another study proposed a smaller MLA of ≤ 4.5 mm² as a useful surrogate of functional significance in patients with isolated ostial and shaft intermediate LMCA stenosis.¹⁴⁾ Ideally, a dual pullback from the left anterior descending artery (LAD) and left circumflex artery (LCX) should be used to fully characterize the LMCA bifurcation and avoid overestimation of, for example, the MLA at the LCX ostium from a LAD pullback.

Physiological guidance by means of FFR or instantaneous wave-free ratio (iFR) may be helpful for the evaluation of intermediate or ambiguous LMCA lesions.¹⁵⁾ In this setting, the visual-functional mismatch between coronary angiography and FFR can be as high as 30–40%,¹¹⁾¹⁶⁾ and deferring LMCA PCI based on FFR values >0.75 or 0.80 has been shown to be safe.¹⁶⁾¹⁷⁾ Recently, deferring revascularization of LMCA disease with an iFR >0.89 has also shown favorable outcomes.¹⁸⁾¹⁹⁾ However, specific outcome studies evaluating iFR in LMCA disease are required to determine whether iFR could be widely adopted as the sole determinant of revascularization in patients with intermediate LMCA lesions.

MEDICAL TREATMENT FOR LMCA DISEASE

In the COURAGE trial,²⁰⁾ an initial strategy of optimal medical therapy vs. initial revascularization showed similar long-term outcomes in patients with stable CAD, excluding LMCA disease. The use of guideline-directed medical treatment (GDMT) and lifestyle interventions should be encouraged in patients with LMCA disease, as they are for patients with non-LMCA disease. However, the safety of deferred revascularization in patients with stable LMCA disease is not fully understood. Current clinical practice guidelines strongly recommend revascularization in all patients with $\geq 50\%$ stenosis of the LMCA.²¹⁾ This class IA recommendation is based on post hoc analysis of a few historical RCTs involving patients with chronic stable angina, demonstrating the superiority of surgical revascularization over medical treatment in terms of 5–10-year survival.²²⁾²³⁾ However, in the context of updated GDMT, the medical management of patients in these early RCTs was not adequate as only 66% of patients were treated with beta blockers and only 19% received aspirin. In addition, the current widespread use of disease-modifying pharmacological interventions (i.e., statins, inhibitors of the renin-angiotensin-aldosterone system, and more effective antiplatelet agents, such as P2Y12 inhibitors) might alleviate adverse cardiovascular events as secondary prevention.

Patients with LMCA disease may have a diverse risk spectrum. Low-risk patients, such as those with 50–70% stenosis or preserved left ventricular (LV) function, have shown more favorable survival while receiving medical therapy alone than have patients with high-risk features (such as >70% stenosis, poor LV function, elevated LV end-diastolic pressure, or prior myocardial infarction [MI]), with 3-year survival rates of 66% vs. 41%, respectively.²⁴⁾²⁵⁾ The study design of most contemporary large RCTs comparing clinical outcomes after medical vs. revascularization therapy has excluded patients with LMCA disease. Therefore, it has not yet been established whether optimal medical therapy represents a safe and appropriate alternative to revascularization in some selected LMCA groups of low-risk patients with stable LMCA disease.

PCI FOR THE TREATMENT OF LMCA DISEASE

In the 40 years history of PCI, the use of coronary stenting overcame the inherent limitations of balloon angioplasty and expanded the therapeutic role of PCI for LMCA disease. However, although PCI with bare-metal stents has demonstrated technical feasibility and acceptable clinical outcomes in highly selected, elective, low-risk patients,²⁶⁻²⁸⁾ a substantial risk of angiographic and clinical restenosis has hampered the wider use of PCI for patients with LMCA disease. In the mid-2000s, the introduction of DES was associated with superior efficacy with respect to restenosis and repeat revascularization and therefore the use of

PCI for the treatment of LMCA has increased significantly.²⁹⁻³¹⁾ In addition to significant improvements in stent technology, improved interventional techniques and adjunctive pharmacotherapy have progressively enhanced PCI outcomes in patients with LMCA disease.²⁾ Contemporary second- and third-generation DES employ improved technology and engineering, including thinner strut platforms, easier delivery profiles, biocompatible or bioresorbable polymers, and more effective antiproliferative drugs.³²⁾ In the contemporary PCI setting of LMCA disease, one issue of clinical interest is that there are substantial differences between contemporary DES with regard to efficacy and safety outcomes. A recent merged analysis involving 4,470 patients with LMCA disease who underwent PCI with second-generation DES showed no significant differences in the 3-year rate of target-vessel failure among different DES.³³⁾ In addition, the incidence of definite stent thrombosis was extremely low (<1.0%) for all types of DES.

PCI strategy and technique

LMCA PCI conducted by experienced operators (i.e., those who have performed at least 15 LMCA PCIs per year for at least 3 consecutive years) is associated with better short- and long-term outcomes than LMCA PCI performed by less experienced operators.³⁴⁾ PCI of the LMCA ostium or shaft is a straightforward procedure that is associated with a lower requirement for repeat revascularization than PCI of the distal LMCA bifurcation.³⁵⁾ The ostium of the LMCA lacks the tunica adventitia and is richer in smooth muscle cells and elastic tissue than any other portion of the LMCA and its branches, which requires attention to ensure that stent expansion is adequate.³⁶⁾

For LMCA bifurcation disease, many challenging technical issues remain. Generally, provisional stenting has been advocated as the preferred approach in bifurcation lesions, as it is technically simpler with at least similar outcomes to a systematic 2-stent strategy.³⁷⁻³⁹⁾ In practice, the single-stent crossover technique has been used more frequently, in as many as 60% of all LMCA bifurcation treatments.⁴⁰⁾ However, this was recently challenged by two RCTs that compared the double-kissing (DK) crush technique with the culotte and provisional stenting strategies for the treatment of true distal LMCA bifurcations.⁴¹⁻⁴³⁾ In both studies, the DK crush technique significantly reduced the primary composite ischemic endpoint.

Selection of a single- or 2-stent technique should be based on disease involvement of the LCX ostium, as side-branch compromise after crossover stenting is frequent. Therefore, to determine the choice of a single- or 2-stent strategy, IVUS provides accurate information for both main- and side-branch disease status in LMCA bifurcation lesions. After main-stent crossover from the proximal LAD to the LM, geometric changes in the LCX ostium were related mainly to carina shift, reduction of MLA, and increased eccentricity of the external elastic membrane and carina angle between the LAD and the LCX.⁴⁴⁾ In cases in which the LCX ostium is significantly compromised (>50%) after provisional stenting, FFR measurement should be considered first before further treatment of the LCX.

IVUS is also useful for procedural optimization of PCI of the distal LMCA bifurcation. After stent implantation, IVUS guidance ensures adequate expansion at the level of the ostial LAD, the ostial LCX, the polygon of confluence (e.g., the convergence zone of the LMCA, LAD, and LCX), and the distal LMCA.⁴⁵⁾ The best IVUS-MSA (minimal stent area) criteria that predicted angiographic in-stent restenosis were 5.0 mm² for the LCX ostium, 6.3 mm² for the LAD ostium, 7.2 mm² for the POC, and 8.2 mm² for the distal LMCA. The 2-year major adverse coronary event-free survival rate was significantly lower in patients with underexpansion of at

least one segment vs. lesions with no underexpansion ($90\pm 3\%$ vs. $98\pm 1\%$, $p<0.001$); post-stenting underexpansion was an independent predictor for major adverse cardiac events.

CLINICAL STUDIES COMPARING PCI AND CABG FOR LMCA DISEASE

Randomized clinical trials

Key clinical trials comparing PCI and CABG from the first-generation DES era to the second-generation DES era are summarized in **Table 1**. The SYNTAX study was a key pivotal trial; in the LMCA subgroup, no significant differences were seen in the 5-year rate of major adverse cardiac and cerebrovascular events (MACCEs), mortality, or MI between PCI and CABG.⁴⁾ However, the 5-year rate of repeat revascularization was higher after PCI and the rate of stroke was higher after CABG. The first LMCA-specific RCT (RECOMBAT) showed that the 5-year rate of MACCEs, death, MI, or stroke was similar between PCI and CABG, but the rate of target-vessel revascularization was significantly higher after PCI.^{6,46)} These trials prompted the initiation of two additional large-scale RCTs, EXCEL and NOBLE, which involved the use of contemporary DES.^{9,10)} In the EXCEL study, the primary composite endpoint of death, stroke, or MI at 3 years was similar between PCI and CABG (p value for non-inferiority=0.02; p value for superiority=0.98).⁹⁾ PCI was associated with a lower incidence of major periprocedural adverse events (i.e., major arrhythmias, infections, reoperations, bleeding, or transfusions). PCI was also associated with a more rapid recovery and greater improvement in quality of life at 30 days than was CABG, although both procedures resulted in similar quality of life and angina relief at 3 years.⁴⁷⁾ In the NOBLE trial, the primary composite endpoint of all-cause mortality, nonprocedural MI, stroke, or repeat revascularization at 5 years was significantly higher after PCI than after CABG (29% vs. 19% exceeding the limit for non-inferiority, respectively). The difference in favor of CABG was statistically significant (p value for superiority=0.007) and was driven by significantly higher rates of nonprocedural MI, repeat revascularization, and stroke in the PCI arm.¹⁰⁾

There may be several explanations for the inconsistent results seen in the EXCEL and NOBLE studies.⁴⁸⁾ First, different types of DES were used. In EXCEL a thin-strut, fluoropolymer-based CoCr-EES was employed, which was associated with the lowest risk of stent thrombosis of all available DES.⁴⁹⁾ The NOBLE study used first-generation, thicker-strut, stainless-steel, sirolimus-eluting Cypher stents (11%) or the biolimus-eluting Biomatrix Flex stent (89%). A substantial difference in the rate of definite stent thrombosis (0.7% in EXCEL vs. 3% in NOBLE) suggests the differential performance of stenting for LMCA disease. Secondly, the soft clinical endpoint of repeat revascularization was adopted as the key component of the primary endpoint in the NOBLE study. The majority of previous studies have consistently shown that the rate of repeat revascularization is significantly higher after PCI than after CABG. Therefore, the selection of this primary composite outcome may unfairly penalize the PCI stratum. The SYNTAX trial showed that the increase in repeat revascularization in the PCI group did not directly translate into an increase in the incidence of death or MI.⁵⁰⁾ Thirdly, the definitions used for components of the primary composite outcomes differed between the studies, particularly the definition of MI. The Society for Cardiovascular Angiography and Interventions -defined clinically relevant MI definition was used in EXCEL,⁵¹⁾ while periprocedural MI was disregarded in NOBLE. Finally, in the NOBLE study, the rate of stroke was more than two times higher after PCI than after CABG, which is not in agreement with the findings of previous clinical trials comparing PCI and CABG. This

Table 1. Randomized clinical trials of percutaneous coronary intervention vs. coronary artery bypass grafting for left main coronary artery disease

Trial	Recruitment period	PCI/CABG	F/U (years)	SS (mean)	ACS (%)	Distal (%)	MVD (%)	Stent	IMA (%)	Primary endpoint (PCI vs. CABG)	Key secondary endpoints at the longest F/U (PCI vs. CABG)
LEMANS ³⁾⁽⁶⁴⁾	2001–2004	52/53	10	NR	NR	58	91	BMS, DES	81	Change in LVEF at 1 year: 3.3±6.7% vs. 0.5±0.8%, p=0.047	<ul style="list-style-type: none"> Death, CVA, MI, or RR at 10 years: 52.2% vs. 62.5%, p=0.42 Death at 10 years: 21.6% vs. 30.2%, p=0.41 CVA at 10 years: 4.3% vs. 6.3%, p=0.58 MI at 10 years: 8.7% vs. 10.4%, p=0.68 RR at 10 years: 26.1% vs. 31.3%, p=0.39
SYNTAX-Left MAIN ⁴⁾⁽⁶⁵⁾	2005–2007	357/348	5	30	30	61	68	DP-PES	97	Death, CVA, MI, or RR at 1 year: 15.8% vs. 13.6%, p=0.4	<ul style="list-style-type: none"> Death, CVA, MI, RR at 5 years: 36.9% vs. 31.0%, p=0.12 Death/CVA/MI at 5 years: 19.0% vs. 20.8%, p=0.57 Death at 5 years: 12.8% vs. 14.6%, p=0.53 CVA at 5 years: 1.5% vs. 4.3%, p=0.03 MI at 5 years: 8.2% vs. 4.8%, p=0.10 RR at 5 years: 26.7% vs. 15.5%, p<0.001
Boudriot et al. ⁵⁾	2003–2009	100/101	1	23	NR	72	41	DP-SES	99	Death, MI, or RR at 1 year: 19.0% vs. 13.9%, p for non-inferiority=0.19	<ul style="list-style-type: none"> Death or MI at 1 year: 5.0% vs. 7.9%, p for non-inferiority<0.001 Death at 1 year: 2.0% vs. 5.0%, p for non-inferiority<0.001 MI at 1 year: 3.0% vs. 3.0%, p for non-inferiority=0.002 RR at 1 year: 14.0% vs. 5.9%, p for non-inferiority=0
PRECOMBAT ⁶⁾⁽⁴⁶⁾	2004–2009	300/300	5	25	45	64	73	DP-SES	94	Death, stroke, MI, ID-TLR at 1 year: 8.7% vs. 6.7%, p for non-inferiority=0.01	<ul style="list-style-type: none"> Death, stroke, MI, or ID-TLR at 5 years: 17.5% vs. 14.3%, p=0.26 Death, stroke, or MI at 5 years: 8.4% vs. 9.6%, p=0.66 Death at 5 years: 5.7% vs. 7.9%, p=0.32 Stroke at 5 years: 0.7% vs. 0.7%, p=0.99 MI at 5 years: 2.0% vs. 1.7%, p=0.76 RR at 5 years: 13% vs. 7.3%, p=0.02
EXCEL ⁹⁾	2010–2014	948/957	3	21	24	81	51	DP-EES	99	Death, stroke, or MI at 3 years: 15.4% vs. 14.7%, p for non-inferiority=0.02, p=0.98 for superiority	<ul style="list-style-type: none"> Death, stroke, MI, or IDR at 3 years: 23.1% vs. 19.1%, p for non-inferiority=0.01 Death at 3 years: 8.2% vs. 5.9%, p=0.11 Stroke at 3 years: 2.3% vs. 2.9%, p=0.37 MI at 3 years: 8.0% vs. 8.3%, p=0.64 IDR at 3 years: 12.6% vs. 7.5%, p<0.001
NOBLE ¹⁰⁾	2008–2015	592/592	5	22	17	81	NR	BP-BES, DP-SES	93	Death, stroke, nonprocedural MI, RR at 5 years: 29% vs. 19%, p=0.0066	<ul style="list-style-type: none"> Death at 5 years: 12% vs. 9%, p=0.77 Stroke at 5 years: 5% vs. 2%, p=0.073 Nonprocedural MI at 5 years: 7% vs. 2%, p=0.004 RR at 5 years: 16% vs. 10%, p=0.032

ACS = acute coronary syndrome; BMS = bare-metal stent; BP-BES = biodegradable-polymer biolimus-eluting stent; CABG = coronary artery bypass grafting; CVA = cerebrovascular accident; DES = drug-eluting stent; DP-EES = durable-polymer everolimus-eluting stent; DP-SES = durable-polymer sirolimus-eluting stent; DP-PES = durable-polymer paclitaxel-eluting stent; F/U = follow-up; IDR = ischemia-driven revascularization; ID-TLR = ischemia-driven target lesion revascularization; IMA = internal mammary artery; LVEF = left ventricular ejection fraction; MI = myocardial infarction; MVD = multivessel disease; NR = not reported; PCI = percutaneous coronary intervention; RR = repeat revascularization; SS = SYNTAX score.

observation lacks a clear explanation and biologic plausibility and is, therefore, likely to be due to a chance effect.⁵²⁾

Meta-analyses

In a meta-analysis of the four largest studies of LMCA revascularization with follow-up available at 3–5 years, incorporating data from the EXCEL and NOBLE trials, the hazard ratio (HR) for death, stroke, or MI with PCI compared with CABG was neutral (1.06) in a random-effects model (p=0.60).⁵³⁾ Based on individual patient data reconstruction, the Kaplan-Meier estimates of death, stroke, or MI at 5 years were 18.3% for PCI and 16.8% for CABG (p=0.52). No statistically significant subgroup interaction for this combined outcome was noted across

studies based on the generation of DES used for PCI (p value for interaction=0.25). There were no significant differences in the pooled effects for death (HR, 1.04; p=0.77) and cardiac death (HR, 1.00; p=0.99). The endpoints of MI and stroke also did not differ between the PCI and CABG groups (HR, 1.48; p=0.17 and 0.87; p=0.72, respectively), but these outcomes were confounded by high heterogeneity across the trials. Repeat revascularization was consistently higher following PCI in all trials, leading to a pooled HR of 1.70 (p<0.001). In another meta-analysis, including all the six trials available to date, missing data were collected by the principal investigators, enabling further subgroup analyses.⁵⁴ PCI was found to significantly reduce death, MI, or stroke by 36% within 30 days. PCI reduced periprocedural MI by 33%, but this effect was offset by 93% more spontaneous MIs beyond 30 days after the procedure. Cardiac death differed in relation to angiographic complexity in that it tended to be lower with PCI among patients with low SYNTAX scores and higher in patients with high SYNTAX scores.

A recent large-scale, pooled analysis of individual patient data reported a comparable treatment effect for PCI and CABG with regard to all-cause mortality up to 5 years in selected patients participating in RCTs.⁵⁵ This analysis included 11 RCTs involving 11,518 patients who were assigned to undergo PCI (n=5,753) or CABG (n=5,765). The 5-year rate of all-cause mortality was 11.2% after PCI and 9.2% after CABG (HR, 1.20; 95% CI, 1.06–1.37; p=0.004). Interestingly, the 5-year all-cause mortality differed significantly between the two interventions in patients with multivessel disease (11.5% after PCI vs. 8.9% after CABG; HR, 1.28; 95% CI, 1.09–1.49; p=0.002), including in those with diabetes (15.5% vs. 10.0%, respectively; HR, 1.48; 95% CI, 1.19–1.84; p=0.0004), but not in those without diabetes (8.7% vs. 8.0%, respectively; HR, 1.08; 95% CI, 0.86–1.36; p=0.49). By contrast, the 5-year rate of all-cause mortality was similar between the two groups in patients with LMCA disease (10.7% after PCI vs. 10.5% after CABG; HR, 1.07; 95% CI, 0.87–1.33; p=0.52), regardless of diabetes status and SYNTAX score.

Registries

Large registries of patients undergoing revascularization for LMCA disease in different geographical regions are useful resources to generalize the findings from RCTs into the daily clinical scenario. The MAIN-COMPARE, IRIS-MAIN, PRECOMBAT-2, and DELTA-2 registries cover a treatment period of 2000–2015,²⁾⁵⁶⁻⁵⁹⁾ and are summarized in **Table 2**. Compared with PCI patients enrolled in the EXCEL trial, the mean SYNTAX score in these registries tended to be higher. In the DELTA-2 study of LMCA PCI with second-generation DES (n=3,986), Kaplan-Meier estimates of events at 2 years were 9.5% and 16.7% for death or target-vessel revascularization, respectively.⁵⁶⁾ HRs for PCI vs. CABG with respect to the composite of death, stroke, or MI were similar between the PCI and CABG groups.

Recently, an observation study of the MAIN-COMPARE registry reported 10-year comparative outcomes of PCI and CABG for LMCA disease.⁶⁰⁾ Overall, there was no significant difference in the adjusted risk of death and the composite outcome between the PCI group and the CABG group up to 10 years. The risk of target-vessel revascularization was significantly higher in the PCI group. However, in the cohort comparing DES and concurrent CABG, DES was associated with a higher risk of death (HR, 1.35; 95% CI, 1.00–1.81) and the composite outcome of death, Q-wave MI, or stroke (HR, 1.46; 95% CI, 1.10–1.94) than was CABG after 5 years. In the DES era, the application of PCI for LMCA disease was substantially expanded and widely performed in patients with a broader range of clinical and anatomical complexities.⁶¹⁾ Therefore, the results of this study can be interpreted as follows: CABG is associated with superior long-term outcomes compared with multivessel PCI in patients

Table 2. Contemporary large observation registries of percutaneous coronary intervention vs. coronary artery bypass grafting for left main coronary artery disease

Study	Enrolment period	Number	SS (mean)	ACS (%)	Distal (%)	MVD (%)	Stent	IMA (%)	Key outcome (PCI vs. CABG)	Adjusted outcome (PCI vs. CABG)
MAIN-COMPARE (Wave 2) ^{57,58)}	2000–2006	PCI, 784 CABG, 690	PCI, NR CABG, NR	PCI, 63 CABG, 76	PCI, 57 CABG, 53	PCI, 58 CABG, 88	G2-DES	NR	Death, Q-wave MI, or stroke at 5 years: 12.7% vs. 16.3%, p=0.02	HR, 0.99; 95% CI, 0.73–1.36; p=0.99
PRECOMBAT-2 ⁵⁹⁾	2009–2010	PCI, 334 CABG, 272	PCI, 21 CABG, 27	PCI, 45 CABG, 54	PCI, 72 CABG, 60	PCI, 57 CABG, 75	G2-DES	NR	Death, MI, stroke, or ischemia-driven TVR at 540 days: 8.9% vs. 6.7%, p=0.23	HR, 0.84; 95% CI, 0.51–1.40; p=0.51
IRIS-MAIN (Wave 3) ²⁾	2007–2013	PCI, 1,658 CABG, 704	PCI, NR CABG, NR	PCI, 55 CABG, 57	PCI, 65 CABG, 72	PCI, 64 CABG, 91	G2-DES	95	Death, stroke, or MI at 3 years: NR	HR, 0.91; 95% CI, 0.68–1.21; p=0.50
DELTA-2 ⁵⁶⁾	2006–2015	PCI, 3,986 CABG, 901	PCI, 27 CABG, 38	PCI, 36 CABG, 65	PCI, 85 CABG, 58	PCI, 74 CABG, 94	G2-DES	NR	Death, CVA, or MI at 501 days: 10.3% vs. 11.6%, p=NR	HR, 0.73; 95% CI, 0.55–0.98; p=0.03

ACS = acute coronary syndrome; CABG = coronary artery bypass grafting; CI = confidence interval; CVA = cerebrovascular accident; DES = drug-eluting stent; G2 = second-generation; HR = hazard ratio; IMA = internal mammary artery; MI = myocardial infarction; MVD = multivessel disease; NR = not reported; PCI = percutaneous coronary intervention; SS = SYNTAX score; TVR = target vessel revascularization.

with high clinical and anatomical complexity. It is interesting to note that the suggestion of superiority of CABG over PCI during long-term follow-up was also seen in the EXCEL and NOBEL trials, which reported a trend towards late catch-up or crossover in the rate of death or the composite endpoint of death, stroke, or MI favoring CABG over PCI during the late period of follow-up. Therefore, longer-term follow-up is necessary to examine additional differences between PCI and CABG over time.

Revascularization guidelines

Existing clinical practice guidelines continue to advocate CABG surgery as the singular class I indication for myocardial revascularization of LMCA disease. However, more recent RCTs and registry studies support PCI as a reasonable alternative in selected patients with less complex LMCA anatomy.

As new evidence has become available, guideline recommendations for LMCA revascularization have slowly evolved over time in both Europe and the US (**Table 3**). Recently, the 2018 European Society of Cardiology guidelines incorporated compelling data from the EXCEL and NOBLE trials, as well as the results of the pooled analysis.⁶²⁾ The 2018 European guideline indicates the same class of recommendation, but all evidence levels have been upgraded to level A. For PCI in LMCA with intermediate anatomical complexity, the previous class IIa recommendation was maintained in view of the incomplete 5-year follow-up in the two largest RCTs in this setting. In the future, the guideline will propose less restrictive indications for PCI, thereby expanding the patient pool that might be eligible for PCI. In addition, given that SYNTAX score was not an important factor for decision-making regarding optimal revascularization and to differentiate the comparative outcomes between CABG and PCI in the EXCEL and NOBLE studies, it may be debated whether the SYNTAX score can play a pivotal role in decision-making regarding LMCA revascularization.

The heart team approach

Regardless of which method of revascularization is used, current guidelines highlight the importance of a ‘heart team’ approach to the management of LMCA disease. The heart team evaluates the risks and benefits of PCI, surgery, or medical treatment alone, taking into account the patient’s informed preference (**Figure 1**). In general, PCI offers more rapid recovery and a lower early adverse event rate, whereas CABG offers more durable revascularization. However, the relative outcomes of PCI vs. CABG can be attributed to a complex interplay of patient comorbidities, coronary anatomic complexity, and ventricular

Table 3. Secular change of myocardial revascularization guidelines for left main coronary artery disease

Guideline	Class of recommendation	Level of evidence
2005 ACC/AHA/SCAI ⁽⁶⁶⁾	III—PCI is not recommended in patients with unprotected LMCA disease and eligibility for CABG	C
2005 ESC/EACTS ⁽⁶⁷⁾	IIb—Stenting for unprotected LMCA disease should only be considered in the absence of other revascularization options	C
2009 ACC/AHA/SCAI ⁽⁶⁸⁾	IIb—PCI of the LMCA with stents as an alternative to CABG may be considered in patients with anatomic conditions that are associated with a low risk of PCI procedural complications and clinical conditions that predict an increased risk of adverse surgical outcomes	B
2010 ESC/EACTS ⁽⁶⁹⁾	IIa—LMCA isolated or β 1VD, ostium/shaft IIb—LMCA isolated or β 1VD, distal bifurcation IIb—LMCA β 2VD or 3VD, SYNTAX score \leq 32 III—LMCA β 2VD or 3VD, SYNTAX score \geq 33	B
2011 ACCF/AHA/SCAI ⁽²¹⁾	IIa—For SIHD patients when both of the following are present: • Anatomic conditions associated with a low risk of PCI procedural complications and a high likelihood of good long-term outcomes (e.g., a low SYNTAX score [\neq 22], ostial or trunk left main stenosis) • Clinical characteristics that predict a significantly increased risk of adverse surgical outcomes (e.g., STS-predicted risk of operative mortality $>$ 5%)	B
	IIb—For SIHD patients when both of the following are present: • Anatomic conditions associated with a low-to-intermediate risk of PCI procedural complications and an intermediate-to-high likelihood of good long-term outcomes (e.g., low-intermediate SYNTAX score of $<$ 33, bifurcation left main stenosis) • Clinical characteristics that predict an increased risk of adverse surgical outcomes (e.g., moderate-severe chronic obstructive pulmonary disease, disability from previous stroke, or previous cardiac surgery; STS-predicted risk of operative mortality $>$ 2%)	B
	III: HARM—For SIHD patients (vs. performing CABG) with unfavorable anatomy for PCI who are good candidates for CABG	B
2014 ESC/EACTS ⁽⁷⁰⁾	I—LMCA with a SYNTAX score \leq 22	B
	IIa—LMCA with a SYNTAX score 23–32	
	III—LMCA with a SYNTAX score \geq 33	
2014 ACC/AHA/AATS/PCNA/SCAI/STS ⁽⁷¹⁾	IIa—For SIHD patients when both of the following are present: • Anatomic conditions associated with a low risk of PCI procedural complications and a high likelihood of good long-term outcomes (e.g., a low SYNTAX score [\leq 22], ostial or trunk left main stenosis) • Clinical characteristics that predict a significantly increased risk of adverse surgical outcomes (e.g., STS-predicted risk of operative mortality $>$ 5%)	B
	IIb—For SIHD patients when both of the following are present: • Anatomic conditions associated with a low-to-intermediate risk of PCI procedural complications and an intermediate-to-high likelihood of good long-term outcome (e.g., low-intermediate SYNTAX score of $<$ 33, bifurcation left main stenosis) • Clinical characteristics that predict an increased risk of adverse surgical outcomes (e.g., moderate-severe chronic obstructive pulmonary disease, disability from previous stroke, or previous cardiac surgery; STS-predicted risk of operative mortality $>$ 2%)	B
	III: HARM—For SIHD patients (vs. performing CABG) with unfavorable anatomy for PCI and who are good candidates for CABG	B
2018 ESC/EACTS ⁽⁶²⁾	I—LMCA with a SYNTAX score \leq 22	A
	IIa—LMCA with a SYNTAX score 23–32	
	III—LMCA with a SYNTAX score \geq 33	

AATS = American Association for Thoracic Surgery; ACC = American College of Cardiology; ACCF = American College of Cardiology Foundation; AHA = American Heart Association; CABG = coronary artery bypass grafting; EACTS = European Association for Cardio-Thoracic Surgery; ESC = European Society of Cardiology; LMCA = left main coronary artery; PCI = percutaneous coronary intervention; PCNA = Preventive Cardiovascular Nurses Association; SCAI = Society for Cardiovascular Angiography and Interventions; SIHD = stable ischemic heart disease; STS = Society of Thoracic Surgeons; VD = vessel disease.

function, in addition to other less tangible factors such as operator expertise and compliance with medication. The complexity and extent of coexisting CAD with the intention of achieving complete revascularization should also be considered by the heart team. Previous evaluation has shown that major adverse cardiovascular events are higher in patients with incomplete revascularization than in those with complete revascularization regardless of the revascularization strategy.⁽⁶³⁾ The heart team approach is critical when evaluating the risks and benefits of surgery in high- and extreme-risk populations. Additional clinical factors that are not included in most risk models also need to be considered by the heart team when making




Figure 1. Heart team approach for LMCA revascularization.

Figure adapted with permission from Park et al.⁴⁸⁾

CABG = coronary artery bypass grafting; CTO = chronic total occlusion; EF = ejection fraction; DAPT = dual antiplatelet therapy; LM = left main; MI = myocardial infarction; MVD = multivessel disease; PCI = percutaneous coronary intervention.

management recommendations, including frailty, cognitive status, surgical recovery and social support, quality of life, life expectancy, patient preference, and any potential concerns regarding tolerance or adherence with long-term dual antiplatelet therapy.

CONCLUSIONS

Over the past 20 years, significant advancements in stent technology, technical refinement, image and physiological guidance, and adjunctive drug therapy have led to progressive improvements in outcomes following PCI in patients with LMCA disease. In the contemporary clinical setting, LMCA PCI has become a viable option in daily practice not only for patients with less complex clinical and anatomic characteristics (i.e., isolated left main disease, ostial or shaft left main disease, or additional less complex CAD), but also for patients with complex clinical and anatomic characteristics (i.e., distal LMCA bifurcation or those with acute MI or unsuitability for CABG).

Which approach will be of most benefit to individual patients with LMCA disease should be decided by the local heart team, which comprises a general cardiologist, interventional cardiologist, and cardiac surgeon. The heart team will consider the clinical circumstances,

any technical issues, and the likelihood of safely achieving complete revascularization with each procedure. It will also be important to consider the patient's own preference once the procedures have been explained in full.

REFERENCES

1. Park SJ, Park DW. Percutaneous coronary intervention with stent implantation versus coronary artery bypass surgery for treatment of left main coronary artery disease: is it time to change guidelines? *Circ Cardiovasc Interv* 2009;2:59-68.
[PUBMED](#) | [CROSSREF](#)
2. Lee PH, Ahn JM, Chang M, et al. Left main coronary artery disease: secular trends in patient characteristics, treatments, and outcomes. *J Am Coll Cardiol* 2016;68:1233-46.
[PUBMED](#) | [CROSSREF](#)
3. Buszman PE, Buszman PP, Banasiewicz-Szkróbka I, et al. Left main stenting in comparison with surgical revascularization: 10-year outcomes of the (left main coronary artery stenting) LE MANS trial. *JACC Cardiovasc Interv* 2016;9:318-27.
[PUBMED](#) | [CROSSREF](#)
4. Morice MC, Serruys PW, Kappetein AP, et al. Outcomes in patients with de novo left main disease treated with either percutaneous coronary intervention using paclitaxel-eluting stents or coronary artery bypass graft treatment in the Synergy Between Percutaneous Coronary Intervention with TAXUS and Cardiac Surgery (SYNTAX) trial. *Circulation* 2010;121:2645-53.
[PUBMED](#) | [CROSSREF](#)
5. Boudriot E, Thiele H, Walther T, et al. Randomized comparison of percutaneous coronary intervention with sirolimus-eluting stents versus coronary artery bypass grafting in unprotected left main stem stenosis. *J Am Coll Cardiol* 2011;57:538-45.
[PUBMED](#) | [CROSSREF](#)
6. Park SJ, Kim YH, Park DW, et al. Randomized trial of stents versus bypass surgery for left main coronary artery disease. *N Engl J Med* 2011;364:1718-27.
[PUBMED](#) | [CROSSREF](#)
7. Bangalore S, Kumar S, Fusaro M, et al. Short- and long-term outcomes with drug-eluting and bare-metal coronary stents: a mixed-treatment comparison analysis of 117 762 patient-years of follow-up from randomized trials. *Circulation* 2012;125:2873-91.
[PUBMED](#) | [CROSSREF](#)
8. Palmerini T, Benedetto U, Biondi-Zoccai G, et al. Long-term safety of drug-eluting and bare-metal stents: evidence from a comprehensive network meta-analysis. *J Am Coll Cardiol* 2015;65:2496-507.
[PUBMED](#) | [CROSSREF](#)
9. Stone GW, Sabik JF, Serruys PW, et al. Everolimus-eluting stents or bypass surgery for left main coronary artery disease. *N Engl J Med* 2016;375:2223-35.
[PUBMED](#) | [CROSSREF](#)
10. Mäkikallio T, Holm NR, Lindsay M, et al. Percutaneous coronary angioplasty versus coronary artery bypass grafting in treatment of unprotected left main stenosis (NOBLE): a prospective, randomised, open-label, non-inferiority trial. *Lancet* 2016;388:2743-52.
[PUBMED](#) | [CROSSREF](#)
11. Park SJ, Kang SJ, Ahn JM, et al. Visual-functional mismatch between coronary angiography and fractional flow reserve. *JACC Cardiovasc Interv* 2012;5:1029-36.
[PUBMED](#) | [CROSSREF](#)
12. Hwang MH, Meadows WR, Palac RT, et al. Progression of native coronary artery disease at 10 years: insights from a randomized study of medical versus surgical therapy for angina. *J Am Coll Cardiol* 1990;16:1066-70.
[PUBMED](#) | [CROSSREF](#)
13. de la Torre Hernandez JM, Hernández Hernandez F, Alfonso F, et al. Prospective application of pre-defined intravascular ultrasound criteria for assessment of intermediate left main coronary artery lesions results from the multicenter LITRO study. *J Am Coll Cardiol* 2011;58:351-8.
[PUBMED](#) | [CROSSREF](#)
14. Park SJ, Ahn JM, Kang SJ, et al. Intravascular ultrasound-derived minimal lumen area criteria for functionally significant left main coronary artery stenosis. *JACC Cardiovasc Interv* 2014;7:868-74.
[PUBMED](#) | [CROSSREF](#)

15. Modi BN, van de Hoef TP, Piek JJ, Perera D. Physiological assessment of left main coronary artery disease. *EuroIntervention* 2017;13:820-7.
[PUBMED](#) | [CROSSREF](#)
16. Hamilos M, Muller O, Cuisset T, et al. Long-term clinical outcome after fractional flow reserve-guided treatment in patients with angiographically equivocal left main coronary artery stenosis. *Circulation* 2009;120:1505-12.
[PUBMED](#) | [CROSSREF](#)
17. Puri R, Kapadia SR, Nicholls SJ, Harvey JE, Kataoka Y, Tuzcu EM. Optimizing outcomes during left main percutaneous coronary intervention with intravascular ultrasound and fractional flow reserve: the current state of evidence. *JACC Cardiovasc Interv* 2012;5:697-707.
[PUBMED](#) | [CROSSREF](#)
18. Davies JE, Sen S, Dehbi HM, et al. Use of the instantaneous wave-free ratio or fractional flow reserve in PCI. *N Engl J Med* 2017;376:1824-34.
[PUBMED](#) | [CROSSREF](#)
19. Götzberg M, Christiansen EH, Gudmundsdottir JJ, et al. Instantaneous wave-free ratio versus fractional flow reserve to guide PCI. *N Engl J Med* 2017;376:1813-23.
[PUBMED](#) | [CROSSREF](#)
20. Boden WE, O'Rourke RA, Teo KK, et al. Optimal medical therapy with or without PCI for stable coronary disease. *N Engl J Med* 2007;356:1503-16.
[PUBMED](#) | [CROSSREF](#)
21. Levine GN, Bates ER, Blankenship JC, et al. 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. *Circulation* 2011;124:2574-609.
[PUBMED](#) | [CROSSREF](#)
22. The Veterans Administration Coronary Artery Bypass Surgery Cooperative Study Group. Eleven-year survival in the Veterans Administration randomized trial of coronary bypass surgery for stable angina. *N Engl J Med* 1984;311:1333-9.
[PUBMED](#) | [CROSSREF](#)
23. Varnauskas E. Twelve-year follow-up of survival in the randomized European Coronary Surgery Study. *N Engl J Med* 1988;319:332-7.
[PUBMED](#) | [CROSSREF](#)
24. Conley MJ, Ely RL, Kisslo J, Lee KL, McNeer JF, Rosati RA. The prognostic spectrum of left main stenosis. *Circulation* 1978;57:947-52.
[PUBMED](#) | [CROSSREF](#)
25. Takaro T, Peduzzi P, Detre KM, et al. Survival in subgroups of patients with left main coronary artery disease. *Circulation* 1982;66:14-22.
[PUBMED](#) | [CROSSREF](#)
26. Park SJ, Park SW, Hong MK, et al. Stenting of unprotected left main coronary artery stenoses: immediate and late outcomes. *J Am Coll Cardiol* 1998;31:37-42.
[PUBMED](#) | [CROSSREF](#)
27. Park SJ, Hong MK, Lee CW, et al. Elective stenting of unprotected left main coronary artery stenosis: effect of debulking before stenting and intravascular ultrasound guidance. *J Am Coll Cardiol* 2001;38:1054-60.
[PUBMED](#) | [CROSSREF](#)
28. Takagi T, Stankovic G, Finci L, et al. Results and long-term predictors of adverse clinical events after elective percutaneous interventions on unprotected left main coronary artery. *Circulation* 2002;106:698-702.
[PUBMED](#) | [CROSSREF](#)
29. Park SJ, Kim YH, Lee BK, et al. Sirolimus-eluting stent implantation for unprotected left main coronary artery stenosis: comparison with bare metal stent implantation. *J Am Coll Cardiol* 2005;45:351-6.
[PUBMED](#) | [CROSSREF](#)
30. Chieffo A, Park SJ, Valgimigli M, et al. Favorable long-term outcome after drug-eluting stent implantation in nonbifurcation lesions that involve unprotected left main coronary artery: a multicenter registry. *Circulation* 2007;116:158-62.
[PUBMED](#) | [CROSSREF](#)
31. Chieffo A, Stankovic G, Bonizzoni E, et al. Early and mid-term results of drug-eluting stent implantation in unprotected left main. *Circulation* 2005;111:791-5.
[PUBMED](#) | [CROSSREF](#)
32. Stefanini GG, Holmes DR Jr. Drug-eluting coronary-artery stents. *N Engl J Med* 2013;368:254-65.
[PUBMED](#) | [CROSSREF](#)

33. Lee PH, Kwon O, Ahn JM, et al. Safety and effectiveness of second-generation drug-eluting stents in patients with left main coronary artery disease. *J Am Coll Cardiol* 2018;71:832-41.
[PUBMED](#) | [CROSSREF](#)
34. Xu B, Redfors B, Yang Y, et al. Impact of operator experience and volume on outcomes after left main coronary artery percutaneous coronary intervention. *JACC Cardiovasc Interv* 2016;9:2086-93.
[PUBMED](#) | [CROSSREF](#)
35. Naganuma T, Chieffo A, Meliga E, et al. Long-term clinical outcomes after percutaneous coronary intervention for ostial/mid-shaft lesions versus distal bifurcation lesions in unprotected left main coronary artery: the DELTA Registry (drug-eluting stent for left main coronary artery disease): a multicenter registry evaluating percutaneous coronary intervention versus coronary artery bypass grafting for left main treatment. *JACC Cardiovasc Interv* 2013;6:1242-9.
[PUBMED](#) | [CROSSREF](#)
36. Park SJ, Park DW. Left main stenting: is it a different animal? *EuroIntervention* 2010;6 Suppl J:J112-7.
[PUBMED](#) | [CROSSREF](#)
37. Maeng M, Holm NR, Erglis A, et al. Long-term results after simple versus complex stenting of coronary artery bifurcation lesions: Nordic Bifurcation Study 5-year follow-up results. *J Am Coll Cardiol* 2013;62:30-4.
[PUBMED](#) | [CROSSREF](#)
38. Hildick-Smith D, de Belder AJ, Cooter N, et al. Randomized trial of simple versus complex drug-eluting stenting for bifurcation lesions: the British Bifurcation Coronary Study: old, new, and evolving strategies. *Circulation* 2010;121:1235-43.
[PUBMED](#) | [CROSSREF](#)
39. Colombo A, Bramucci E, Saccà S, et al. Randomized study of the crush technique versus provisional side-branch stenting in true coronary bifurcations: the CACTUS (Coronary Bifurcations: Application of the Crushing Technique Using Sirolimus-Eluting Stents) Study. *Circulation* 2009;119:71-8.
[PUBMED](#) | [CROSSREF](#)
40. Kim WJ, Kim YH, Park DW, et al. Comparison of single- versus two-stent techniques in treatment of unprotected left main coronary bifurcation disease. *Catheter Cardiovasc Interv* 2011;77:775-82.
[PUBMED](#) | [CROSSREF](#)
41. Chen SL, Xu B, Han YL, et al. Comparison of double kissing crush versus Culotte stenting for unprotected distal left main bifurcation lesions: results from a multicenter, randomized, prospective DKCRUSH-III study. *J Am Coll Cardiol* 2013;61:1482-8.
[PUBMED](#) | [CROSSREF](#)
42. Chen SL, Zhang JJ, Han Y, et al. Double kissing crush versus provisional stenting for left main distal bifurcation lesions: DKCRUSH-V randomized trial. *J Am Coll Cardiol* 2017;70:2605-17.
[PUBMED](#) | [CROSSREF](#)
43. Chen SL, Xu B, Han YL, et al. Clinical outcome after DK crush versus culotte stenting of distal left main bifurcation lesions: the 3-year follow-up results of the DKCRUSH-III study. *JACC Cardiovasc Interv* 2015;8:1335-42.
[PUBMED](#) | [CROSSREF](#)
44. Kang SJ, Mintz GS, Kim WJ, et al. Changes in left main bifurcation geometry after a single-stent crossover technique: an intravascular ultrasound study using direct imaging of both the left anterior descending and the left circumflex coronary arteries before and after intervention. *Circ Cardiovasc Interv* 2011;4:355-61.
[PUBMED](#) | [CROSSREF](#)
45. Kang SJ, Ahn JM, Song H, et al. Comprehensive intravascular ultrasound assessment of stent area and its impact on restenosis and adverse cardiac events in 403 patients with unprotected left main disease. *Circ Cardiovasc Interv* 2011;4:562-9.
[PUBMED](#) | [CROSSREF](#)
46. Ahn JM, Roh JH, Kim YH, et al. Randomized trial of stents versus bypass surgery for left main coronary artery disease: 5-year outcomes of the PRECOMBAT study. *J Am Coll Cardiol* 2015;65:2198-206.
[PUBMED](#) | [CROSSREF](#)
47. Baron SJ, Chinnakondapalli K, Magnuson EA, et al. Quality-of-life after everolimus-eluting stents or bypass surgery for left-main disease: results from the EXCEL trial. *J Am Coll Cardiol* 2017;70:3113-22.
[PUBMED](#) | [CROSSREF](#)
48. Park DW, Park SJ. Percutaneous coronary intervention of left main disease: Pre- and Post-EXCEL (Evaluation of XIENCE Everolimus Eluting Stent Versus Coronary Artery Bypass Surgery for Effectiveness of Left Main Revascularization) and NOBLE (Nordic-Baltic-British Left Main Revascularization Study) Era. *Circ Cardiovasc Interv* 2017;10:e004792.
[PUBMED](#) | [CROSSREF](#)

49. Palmerini T, Biondi-Zoccai G, Della Riva D, et al. Stent thrombosis with drug-eluting and bare-metal stents: evidence from a comprehensive network meta-analysis. *Lancet* 2012;379:1393-402.
[PUBMED](#) | [CROSSREF](#)
50. Serruys PW, Morice MC, Kappetein AP, et al. Percutaneous coronary intervention versus coronary-artery bypass grafting for severe coronary artery disease. *N Engl J Med* 2009;360:961-72.
[PUBMED](#) | [CROSSREF](#)
51. Moussa ID, Klein LW, Shah B, et al. Consideration of a new definition of clinically relevant myocardial infarction after coronary revascularization: an expert consensus document from the Society for Cardiovascular Angiography and Interventions (SCAI). *J Am Coll Cardiol* 2013;62:1563-70.
[PUBMED](#) | [CROSSREF](#)
52. Mack M, Holmes DR. Randomised trials in left main disease: a NOBLE effort. *Lancet* 2016;388:2715-6.
[PUBMED](#) | [CROSSREF](#)
53. Giacoppo D, Collieran R, Cassese S, et al. Percutaneous coronary intervention vs coronary artery bypass grafting in patients with left main coronary artery stenosis: a systematic review and meta-analysis. *JAMA Cardiol* 2017;2:1079-88.
[PUBMED](#) | [CROSSREF](#)
54. Palmerini T, Serruys P, Kappetein AP, et al. Clinical outcomes with percutaneous coronary revascularization vs coronary artery bypass grafting surgery in patients with unprotected left main coronary artery disease: a meta-analysis of 6 randomized trials and 4,686 patients. *Am Heart J* 2017;190:54-63.
[PUBMED](#) | [CROSSREF](#)
55. Head SJ, Milojevic M, Daemen J, et al. Mortality after coronary artery bypass grafting versus percutaneous coronary intervention with stenting for coronary artery disease: a pooled analysis of individual patient data. *Lancet* 2018;391:939-48.
[PUBMED](#) | [CROSSREF](#)
56. Chieffo A, Tanaka A, Giustino G, et al. The DELTA 2 registry: a multicenter registry evaluating percutaneous coronary intervention with new-generation drug-eluting stents in patients with obstructive left main coronary artery disease. *JACC Cardiovasc Interv* 2017;10:2401-10.
[PUBMED](#) | [CROSSREF](#)
57. Seung KB, Park DW, Kim YH, et al. Stents versus coronary-artery bypass grafting for left main coronary artery disease. *N Engl J Med* 2008;358:1781-92.
[PUBMED](#) | [CROSSREF](#)
58. Park DW, Seung KB, Kim YH, et al. Long-term safety and efficacy of stenting versus coronary artery bypass grafting for unprotected left main coronary artery disease: 5-year results from the MAIN-COMPARE (Revascularization for Unprotected Left Main Coronary Artery Stenosis: Comparison of Percutaneous Coronary Angioplasty Versus Surgical Revascularization) registry. *J Am Coll Cardiol* 2010;56:117-24.
[PUBMED](#) | [CROSSREF](#)
59. Kim YH, Park DW, Ahn JM, et al. Everolimus-eluting stent implantation for unprotected left main coronary artery stenosis. The PRECOMBAT-2 (Premier of Randomized Comparison of Bypass Surgery versus Angioplasty Using Sirolimus-Eluting Stent in Patients with Left Main Coronary Artery Disease) study. *JACC Cardiovasc Interv* 2012;5:708-17.
[PUBMED](#) | [CROSSREF](#)
60. Park DW, Ahn JM, Yun SC, et al. 10-year outcomes of stents versus coronary artery bypass grafting for left main coronary artery disease. *J Am Coll Cardiol* 2018;72:2813-22.
[PUBMED](#) | [CROSSREF](#)
61. Park DW, Kim YH, Yun SC, et al. Complexity of atherosclerotic coronary artery disease and long-term outcomes in patients with unprotected left main disease treated with drug-eluting stents or coronary artery bypass grafting. *J Am Coll Cardiol* 2011;57:2152-9.
[PUBMED](#) | [CROSSREF](#)
62. Neumann FJ, Sousa-Uva M, Ahlsson A, et al. 2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur Heart J* 2019;40:87-165.
[PUBMED](#) | [CROSSREF](#)
63. Farooq V, Serruys PW, Garcia-Garcia HM, et al. The negative impact of incomplete angiographic revascularization on clinical outcomes and its association with total occlusions: the SYNTAX (Synergy Between Percutaneous Coronary Intervention with Taxus and Cardiac Surgery) trial. *J Am Coll Cardiol* 2013;61:282-94.
[PUBMED](#) | [CROSSREF](#)
64. Buszman PE, Buszman PP, Kiesz RS, et al. Early and long-term results of unprotected left main coronary artery stenting: the LE MANS (Left Main Coronary Artery Stenting) registry. *J Am Coll Cardiol* 2009;54:1500-11.
[PUBMED](#) | [CROSSREF](#)

65. Morice MC, Serruys PW, Kappetein AP, et al. Five-year outcomes in patients with left main disease treated with either percutaneous coronary intervention or coronary artery bypass grafting in the synergy between percutaneous coronary intervention with taxus and cardiac surgery trial. *Circulation* 2014;129:2388-94.
[PUBMED](#) | [CROSSREF](#)
66. Smith SC Jr, Feldman TE, Hirshfeld JW Jr, et al. ACC/AHA/SCAI 2005 guideline update for percutaneous coronary intervention: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/SCAI Writing Committee to Update the 2001 Guidelines for Percutaneous Coronary Intervention). *J Am Coll Cardiol* 2006;47:e1-121.
[PUBMED](#) | [CROSSREF](#)
67. Silber S, Albertsson P, Avilés FF, et al. Guidelines for percutaneous coronary interventions. *Eur Heart J* 2005;26:804-47.
[PUBMED](#) | [CROSSREF](#)
68. Kushner FG, Hand M, Smith SC Jr, et al. 2009 focused updates: ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction (updating the 2004 guideline and 2007 focused update) and ACC/AHA/SCAI guidelines on percutaneous coronary intervention (updating the 2005 guideline and 2007 focused update) a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2009;54:2205-41.
[PUBMED](#) | [CROSSREF](#)
69. Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS), European Association for Percutaneous Cardiovascular Interventions (EAPCI), Wijns W, et al. Guidelines on myocardial revascularization. *Eur Heart J* 2010;31:2501-55.
[PUBMED](#) | [CROSSREF](#)
70. Authors/Task Force members, Windecker S, Kolh P, et al. 2014 ESC/EACTS Guidelines on myocardial revascularization: The Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS) Developed with the special contribution of the European Association of Percutaneous Cardiovascular Interventions (EAPCI). *Eur Heart J* 2014;35:2541-619.
[PUBMED](#) | [CROSSREF](#)
71. Fihn SD, Blankenship JC, Alexander KP, et al. 2014 ACC/AHA/AATS/PCNA/SCAI/STS focused update of the guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines, and the American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *Circulation* 2014;130:1749-67.
[PUBMED](#) | [CROSSREF](#)