

Original Article
Preventive &
Social Medicine



Years of Life Lost due to Premature Death in People with Disabilities in Korea: the Korean National Burden of Disease Study Framework

Young-Eun Kim ,¹ Ye-Rin Lee ,² Seok-Jun Yoon ,¹ Young-Ae Kim ,³ and In-Hwan Oh ²

¹Department of Preventive Medicine, Korea University College of Medicine, Seoul, Korea

²Department of Preventive Medicine, Kyung Hee University School of Medicine, Seoul, Korea

³Cancer Policy Branch, National Cancer Control Institute, National Cancer Center, Goyang, Korea



Received: Jun 26, 2018

Accepted: Oct 10, 2018

Address for Correspondence:

In-Hwan Oh, MD

Department of Preventive Medicine, Kyung Hee University School of Medicine, 26 Kyunghaedae-ro, Dongdaemun-gu, Seoul 02447, Korea.

E-mail: parenchyme@gmail.com

© 2019 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Young-Eun Kim

<https://orcid.org/0000-0003-0694-6844>

Ye-Rin Lee

<https://orcid.org/0000-0002-4507-4877>

Seok-Jun Yoon

<https://orcid.org/0000-0003-3297-0071>

Young-Ae Kim

<https://orcid.org/0000-0002-3819-0028>

In-Hwan Oh

<https://orcid.org/0000-0002-5450-9887>

Funding

This research was supported by the R&D grant (No. 2016008) on rehabilitation by Korea National Rehabilitation Center Research Institute, Ministry of Health & Welfare, Republic of Korea.

ABSTRACT

Background: Several studies have been conducted regarding people with disabilities, however no studies have estimated the disease burden due to premature death. As such, we aimed to compare the years of life lost (YLLs) due to premature death between a population with disabilities and the total population of Korea.

Methods: To calculate the YLLs in people with disabilities, we combined disability registration data from the 2002–2011 Ministry of Health and Welfare of Korea and the causes of mortality data by Statistics Korea for 3,158,231 people.

Results: YLLs of people with disabilities were 517,337, which accounted for 23.4% of YLLs of the total population. YLLs per 100,000 people with disabilities were approximately 3.8 times higher than those in the total Korean population. Ischemic stroke was associated with the highest YLLs per 100,000 people, followed by ischemic heart disease, hemorrhagic, and other non-ischemic stroke, diabetes mellitus, and self-harm. Among individuals with physical disabilities, ischemic heart disease was associated with the highest YLLs. For intellectual disability, epilepsy contributed to the most YLLs. For individuals with mental disability, self-harm was the largest contributor to YLLs.

Conclusion: The burden of disease was higher in the population with disabilities than that in the general population. Cardiovascular diseases and cancer had a higher burden of disease than did other diseases in the population with disabilities; thus, overall, non-communicable diseases have a higher burden of disease than communicable diseases or injuries in the population with disabilities than in the general population.

Keywords: Mortality; Disability-Adjusted Life Years; Burden of Disease; Years of Life Lost; People with Disabilities

INTRODUCTION

According to the world report on disability (2011) by the World Health Organization (WHO), approximately 15% of the world's population, i.e., approximately one billion people, are estimated to have disabilities.¹ In Korea, as of 2016, registered individuals with disabilities

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Kim YE, Oh IH. Data curation: Kim YE. Formal analysis: Kim YE, Lee YR. Methodology: Kim YE, Lee YR, Kim YA. Validation: Kim YE, Yoon SJ. Writing - original draft: Kim YE. Writing - editing & editing: Kim YE, Yoon SJ.

account for 4.9% of the total population. The population with disabilities increased in number until 2011 and stabilized since 2012.²

Korea implemented a registration system for individuals with disabilities to better generate disability statistics.³ According to the registration system, disability is registered as one of the following fifteen categories: physical, brain lesions, vision, hearing, language, facial, kidney, heart, liver, respiratory, intestinal and urinary fistula, epilepsy, intellectual, mental, and autism. Each disability is classified into six grades according to severity.

Individuals with disabilities are vulnerable to preventable secondary diseases, comorbidities, untreated mental health statuses, poor oral health, high obesity rates, and early mortality.⁴ Furthermore, approximately 87% of the population with disabilities has one or more health problems, while 75.9% has two or more health problems; a report indicated that the average number of health problems was 4.1.⁴ These statistics indicate that individuals with disabilities are subject to poorer health management than individuals without disabilities, which increases their risk of developing more health problems and early death.

High morbidity and premature death are major problems affecting populations with disabilities. However, most studies on mortality and causes of death in the population with disabilities have focused on specific types of disabilities, such as intellectual disabilities, instead of examining the overall population with disabilities. For example, a British study on individuals with intellectual disabilities reported that all-cause mortality was approximately 3.6 times higher in individuals with intellectual disabilities than in those without.⁵ Furthermore, a US study on individuals with intellectual and developmental disabilities reported that all-cause mortality was approximately 1.8 times higher in individuals with disabilities than in those without.⁶

Similarly, an Irish study on individuals with intellectual disabilities reported that mortality was approximately 4 times higher in individuals with disabilities than in those without.⁷ A systematic review and meta-analysis of 203 studies also reported a 2.22 times higher all-cause mortality in the individuals with mental disabilities than in those without, and potential life lost was approximately 10 years.⁸ However, few studies have examined the burden of premature death in the overall population with disabilities—as opposed to premature death due to specific types of disabilities—in comparison with the population without disabilities and analyzed disability-specific features. In Korea, studies investigating mortality in overall population with disabilities are rare, even for specific types of disabilities.

Therefore, the present study aims to compare the disease burden of premature death between the overall population with disabilities and total population, and examine the effects with specific causes of death amongst age groups and gender. This study also identifies disability-specific features of years of life lost (YLLs) due to premature death.⁹

METHODS

We measured the disease burden of premature death in people with disabilities using YLLs. YLLs are an indicator of mortality due to premature deaths and used to measure disability-adjusted life years (DALYs), a metric introduced by the global burden of disease (GBD) study designed to measure the disease burden of premature death. This study compared the burden of disease between the population with disabilities and total population using

the methodology suggested by the 2012 burden of disease study on Koreans, and the results of that study were compared with ours. The comparisons were made in units of YLLs. The meaning and calculations of YLLs are detailed in the previous literature, and the methodology for studying burden of disease in Koreans has also been described in another study.¹⁰ Here we only outlined the method of calculating YLLs in this study.

In this study, YLLs were calculated using the equation below, with a 3% age-weighting rate and 4% time-discount rate.¹¹ To calculate YLLs due to death, we multiplied the number of deaths by standard life expectancy for each gender and age group, and summed YLLs of all age groups. The standard life expectancy suggested by the 2012 Statistics Korea survey was applied for the calculations.¹²

r = discount rate, β = age-specific weight parameter, K = use of age-specific weight (using age-weight, applied 1; not using age-weight, applied 0), C = constant, a = age at death, L = life expectancy at death.

$$YLL = \frac{KCe^{ra}}{(r + \beta)^2} \{e^{-(r+\beta)(L+a)}[-(r + \beta)(L + a) - 1] - e^{-(r+\beta)a}[-(r + \beta)a - 1]\} + \frac{1 - K}{r} (1 - e^{-rL})$$

To calculate the disease burden of premature death in people with disabilities, we combined disability registration data from the Ministry of Health and Welfare (2002–2011) and the causes of mortality data by Statistics Korea for 3,158,231 people by personal identification number. After excluding people with incomplete data, such as omissions in gender, age, type of disability, or grade of disability, a total of 3,156,038 people were analyzed. The causes of mortality of 68,323 deaths that occurred in 2012 were identified and classified using the Korean National Burden of Disease (KNBD) study classification, a Korean-adapted version of the disease classification system used in the GBD study by the WHO in 2010.¹⁰ The KNBD study classification system comprises four levels. Level 1 is divided into three categories: communicable, maternal, neonatal, and nutritional disorders; non-communicable diseases; and injuries. Diseases are subdivided into 21 categories in level 2, which are further subdivided in level 3, and further still in level 4.⁹

To improve the accuracy of causes of death, in cases where garbage codes were input, the causes of death were redistributed according to the garbage code algorithm originally developed by the GBD 2010 study and was adapted for use in Korea in the KNBD 2012 study.¹⁰ YLLs were calculated for each cause of death, based on deaths that occurred in 2012, and groups with high YLLs were identified among different groups of age (10-year age groups) and types of disabilities.

All analyses were conducted using SAS 9.3 (SAS Institute, Cary, NC, USA).

Ethics statement

The study was approved by the National Cancer Center Institutional Review Board (NCC2016-0094). This study was conducted using secondary data. So, informed consent was not required.

RESULTS

In 2012, 28,000 people with disabilities died, which accounted for 10% of the deaths in the total population. YLLs of people with disabilities were 517,337, which accounted for 23.4% of

YLLs of the total population. YLLs per 100,000 people in the total Korean population were 4,344, but those for people with disabilities were approximately 3.8 times higher (16,392). YLLs of men with disabilities were higher than those of women with disabilities, and YLLs per 100,000 were also higher in men with disabilities than women with disabilities. In terms of age, the population with disabilities demonstrated marked differences compared to the total population. Among the population with disabilities, YLLs were highest in the 70–79-year-old group (25.9%), followed by the 60–69-year-old group (22.2%) and 50–59-year-old group (20.4%). However, in the total population, YLLs were highest in the 50–59-year-old group (19.8%), followed by the 70–79-year-old group (18.0%) and 60–69-year-old group (16.2%). YLLs per 100,000 for the 0–9-year-old group were 14.9 times higher in population with disabilities (27,031) than the total population (1,815); this gap between total population and population with disabilities decreased with advancing age and was 14.0 times higher in population with disabilities than the total population in the 10–19-year-old group and 7.2 in the 20–29-year-old group. In contrast, in the ≥ 80-year-old group, YLLs per 100,000 were higher in the total population than in the population with disabilities (**Table 1**).

For causes of mortality, YLLs due to non-communicable diseases were highest (417,625), accounting for 80.7% of all YLLs, followed by YLLs due to injuries (78,545; 15.2%) and YLLs due to communicable, maternal, neonatal, and nutritional disorders (21,167; 4.1%). Among non-communicable diseases, cardiovascular and circulatory diseases were associated with the highest YLLs (YLLs, 130,856), followed by neoplasms (YLLs, 124,567) and diabetes, urogenital, blood, and endocrine diseases (YLLs, 60,909). For injuries, self-harm and interpersonal violence were associated with the highest YLLs (YLLs, 37,924), followed by unintentional injuries, except for transport injuries (YLLs, 26,404) and transport injuries (YLLs, 13,187). Among communicable, maternal, neonatal, and nutritional disorders, diarrhea, lower respiratory infections, meningitis, and other common infectious diseases were associated with the highest YLLs (YLLs, 9,568), followed by human immunodeficiency virus/acquired immune deficiency syndrome (HIV/AIDS) and tuberculosis (YLLs, 7,921). For level 2 diseases, cardiovascular and circulatory diseases were associated with the highest YLLs per 100,000 people (4,146), which accounted for 25.3% of the total YLLs. Neoplasms had the second-highest YLLs per 100,000 people (3,947; 24.1%), followed by diabetes, urogenital, blood, and endocrine diseases (1,930; 11.8%), and self-harm and interpersonal

Table 1. Total YLL by gender and age group for the total population and people with disabilities in 2012

Variables	Total population			People with disability		
	YLL	Total YLL, % ^a	YLL per 100,000	YLL	Total YLL, % ^a	YLL per 100,000
Gender						
Men	1,360,893	61.6	5,347	330,734	63.9	17,661
Women	847,187	38.4	3,338	186,602	36.1	14,540
Age group, yr						
0–9	84,504	3.8	1,815	6,673	1.3	27,031
10–19	53,824	2.4	819	8,580	1.7	11,500
20–29	123,716	5.6	1,859	13,077	2.5	13,418
30–39	194,959	8.8	2,342	24,946	4.8	12,503
40–49	352,223	16.0	3,977	59,369	11.5	15,607
50–59	436,541	19.8	5,726	105,630	20.4	17,207
60–69	357,998	16.2	8,337	114,831	22.2	18,183
70–79	398,053	18.0	13,844	134,113	25.9	18,363
≥ 80	206,263	9.3	19,233	50,119	9.7	12,412
Total YLL	2,208,080	100.0	4,344	517,337	100.0	16,392

YLL = years of life lost.

^aSum may not add up to exact total due to rounding.

Table 2. YLL for 21 causes of death in individuals with disabilities in 2012

Category	YLL	YLL per 100,000	Total YLL, %
Communicable, maternal, neonatal, and nutritional disorders			
HIV/AIDS and tuberculosis	7,921	251	1.5
Diarrhea, lower respiratory infections, meningitis, and other common infectious diseases	9,568	303	1.8
Neglected tropical diseases and malaria	431	14	0.1
Maternal disorders	203	6	0.0
Neonatal disorders	709	22	0.1
Nutritional deficiencies	929	29	0.2
Other communicable, maternal, neonatal, and nutritional disorders	1,406	45	0.3
Total communicable, maternal, neonatal, and nutritional disorders	21,167	671	4.1
Non-communicable diseases			
Neoplasms	124,567	3,947	24.1
Cardiovascular and circulatory diseases	130,856	4,146	25.3
Chronic respiratory diseases	22,212	704	4.3
Cirrhosis of the liver	20,695	656	4.0
Digestive diseases (except cirrhosis)	10,979	348	2.1
Neurological disorders	34,332	1,088	6.6
Mental and behavioral disorders	3,288	104	0.6
Diabetes, urogenital, blood, and endocrine diseases	60,909	1,930	11.8
Musculoskeletal disorders	3,743	119	0.7
Other non-communicable diseases	6,044	192	1.2
Total non-communicable diseases	417,625	13,233	80.7
Injuries			
Transport injuries	13,187	418	2.5
Unintentional injuries other than transport injuries	26,404	837	5.1
Self-harm and interpersonal violence	37,924	1,202	7.3
Forces of nature, war, and legal intervention	1,030	33	0.2
Total injuries	78,545	2,489	15.2
Total	517,337	16,392	100.0

YLL = years of life lost, HIV/AIDS = human immunodeficiency virus/acquired immune deficiency syndrome.

violence (1,202; 7.2%). YLLs due to these four diseases were 354,256, accounting for 68.5% of the total YLLs (Table 2).

We examined diseases that caused the greatest YLLs in each age group. For the 0–9-year-old group, neonatal disorders caused the greatest YLLs, accounting for 36.6% of the total YLLs (YLLs per 100,000 people, 9,896). Neurological disorders were responsible for the most YLLs in the 10–19-year-old group with disabilities (24.8% of YLLs in the 0–9-year-old group; YLLs per 100,000 people, 2,855), and self-harm and interpersonal violence caused the most YLLs in the 20–39-year-old group with disabilities (22.8% of YLLs in the 20–29-year-old group; 19.0% of YLLs in the 30–39-year-old group). Neoplasms caused the highest YLLs in the 40–69-year-old group with disabilities (21.0% of YLLs in the 40–49-year-old group; 26.4% of YLLs in the 50–59-year-old group; 29.1% of YLLs in the 60–69-year-old group). Cardiovascular and circulatory diseases were associated with the highest YLLs in the ≥ 70-year-old group with disabilities (32.8% of YLLs in the 70–79-year-old group; 38.4% of YLLs in the ≥ 80-year-old group). YLLs due to cardiovascular and circulatory diseases tended to increase with advancing age. Self-harm and interpersonal violence accounted for the highest proportion of YLLs in the 20–29-year-old group; however, this proportion demonstrated a decreasing trend from 20 years of age (Fig. 1).

We analyzed the burden of disease for each disease group. Ischemic stroke was associated with the highest YLLs per 100,000 people (1,329), followed by ischemic heart disease (YLLs per 100,000 people, 1,180), hemorrhagic and other non-ischemic stroke (YLLs per 100,000 people, 1,060), diabetes mellitus (YLLs per 100,000 people, 904), and self-harm (YLLs




Fig. 1. Percentage of YLL for people with disability by cause and age group in 2012. YLL = years of life lost, HIV/AIDS = human immunodeficiency virus/acquired immune deficiency syndrome.

per 100,000 people, 893). YLLs from these five causes of mortality were 169,337, which accounted for 32.7% of total YLLs. The highest cause of mortality in the total population was self-harm (YLLs per 100,000 people, 573), which was the fifth highest cause of mortality in the population with disabilities. Ischemic heart disease ranked second in the total population and population with disabilities, while ischemic stroke ranked seventh in the total population, but first in the population with disabilities. Chronic kidney disease due to hypertension was the cause of mortality with the greatest difference in rank between the total population and the population with disabilities; from 55th in the total population to 25th in the population with disabilities. Furthermore, epilepsy, brain and nervous system cancers, chronic kidney disease due to diabetes mellitus, and Parkinson's disease demonstrated a large increase in rank in the population with disabilities. In contrast, motorized vehicles with two wheels demonstrated the greatest fall in rank; from 22nd in the total population to 40th in the population with disabilities. Pedestrian injury by road vehicle, leukemia, and breast cancer also demonstrated huge decreases in rank in the population with disabilities (Fig. 2).

In terms of age-specific burden of disease, epilepsy was the greatest cause of YLLs in individuals aged 0–19 years, while self-harm, hemorrhagic and other non-ischemic stroke, and ischemic stroke were the greatest causes of YLLs in individuals aged 20–49 years, 50–59 years, and ≥ 60 years, respectively (Table 3). In the total population, preterm birth complications were the greatest cause of YLLs in individuals aged 0–9 years, while self-harm contributed the most among those aged 10–59 years. Among individuals aged 60–69 years lung cancer was the highest cause of YLLs, and for those over 70 years, ischemic heart disease contributed the most YLLs.¹⁰

We also analyzed the burden of disease based on specific types of disabilities. Among individuals with physical disabilities, which account for the highest proportion of disabilities in Korea, ischemic heart disease was associated with the highest YLLs. Furthermore, the




Fig. 2. YLL ranks for the top 25 causes of death for the total population and people with disability in 2012. YLL = years of life lost.

most common disabilities in Korea following ischemic heart disease are brain disorders, hearing disabilities, and visual disturbances. Ischemic stroke was associated with the highest YLLs in individuals with brain disorders, while ischemic heart disease and diabetes mellitus were associated with the highest YLLs in individuals with hearing impairments and visual disturbances, respectively. For individuals with intellectual disabilities, the next most

Table 3. Top cause of death and YLL by age group in individuals with disabilities in 2012

Age group, yr	Cause of death	YLL	Total YLL on age group, %	YLL per 100,000
0-9	Epilepsy	1,490	22.3	6,038
10-19	Epilepsy	1,055	12.3	1,414
20-29	Self-harm	1,886	14.4	1,935
30-39	Self-harm	4,005	16.1	2,008
40-49	Self-harm	6,776	11.4	1,781
50-59	Hemorrhagic and other non-ischemic stroke	8,337	7.9	1,358
60-69	Ischemic stroke	9,586	8.3	1,518
70-79	Ischemic stroke	17,942	13.4	2,457
≥ 80	Ischemic stroke	8,226	16.4	2,038

YLL = years of life lost.

Table 4. Top cause of death and YLL by type of disability in 2012

Type of disability	Cause of death	YLL	YLL per 100,000
External physical			
Limb impairment	Ischemic heart disease	14,209	907
Brain palsy and stroke	Ischemic stroke	25,294	6,166
Visual impairment	Diabetes mellitus	3,540	1,156
Auditory impairment	Ischemic heart disease	3,798	1,201
Lingual impairment	Larynx cancer	646	2,631
Facial deformity	Mouth cancer	45	256
Internal organ			
Renal failure	Chronic kidney disease due to diabetes mellitus	13,516	13,001
Heart failure	Ischemic heart disease	1,238	6,003
Liver failure	Cirrhosis of the liver	3,662	18,568
Respiratory failure	Chronic obstructive pulmonary disease	4,544	16,429
Intestinal and urinary failure	Colon and rectum cancers	4,994	21,764
Epilepsy	Epilepsy	422	3,810
Mental			
Mental retardation	Epilepsy	1,640	863
Mental diseases	Self-harm	5,046	4,337
Developmental disability	Epilepsy	65	376

YLL = years of life lost.

frequent disability in Korea, epilepsy caused the highest YLLs. For individuals with mental disabilities, which constitute 3.5% of the population with disabilities, self-harm caused the highest YLLs (Table 4).

DISCUSSION

We compared the disease burden of premature death (YLLs) in the population with disabilities and the total population. We also compared the causes of death associated with the highest YLLs for each type of disability. Our findings demonstrate that YLLs in the population with disabilities in 2012 was 517,337, which accounted for 23.4% of YLLs in the total population, and YLLs per 100,000 people were approximately 3.8 times higher in the population with disabilities than in the total population. Self-harm was associated with the highest YLLs in the total population, while ischemic stroke ranked first in the population with disabilities. The burden of disease was highest for ischemic heart disease, followed by hemorrhagic and other non-ischemic stroke, diabetes mellitus, and self-harm. YLLs from these five causes of mortality accounted for 32.7% of total YLLs in the population with disabilities. For diseases associated with the highest YLLs for each type of disability, ischemic heart disease resulted in the highest YLLs in the individuals with physical disabilities,

ischemic stroke in individuals with brain disorders, ischemic heart disease in individuals with hearing disabilities, diabetes mellitus in the visually impaired, and epilepsy in individuals with the intellectual disabilities.

In our study, the burden of disease was higher in the population with disabilities than the total population. YLLs per 100,000 people were 16,392 in the population with disabilities, which was approximately four times higher than those in the total population, suggesting that health problems are more significant in individuals with disabilities than in those without. This result was also supported by other studies on individuals with disabilities. According to Hosking et al.,⁵ mortality was 3.6 times higher in individuals with intellectual disabilities than in the general population in England. Another study also reported that all-cause mortality was three times higher in individuals with disabilities than in the general population.¹³ Mortality was especially high in younger individuals with disabilities, which according to the literature, is due to an increase in infant mortality or early neonatal death.^{14,15}

In this study, mean age of the population with disabilities was higher than that of the population without disabilities; high YLLs in individuals with disabilities may be partially affected by age. However, when we compared YLLs in specific age groups, it remained higher in the age groups with disabilities than in the corresponding general population groups. YLLs per 100,000 people were considerably higher in most age groups of the population with disabilities than in those without. For example, the burden of disease was 3.9 times and three times higher in the 30-year-old group with disabilities and 40-year-old group with disabilities, respectively, while this value increased to 14 times in the 10-year-old group. This is also because mortality and YLLs were lowest in the 10-year-old group in the general population. The only exception for higher YLLs in the group with disabilities was the ≥ 80 -year-old group, where YLLs per 100,000 people was lower in the population with disabilities than the general population. This result supports that of a previous study that reported that differences in mortality between the population with and those without disabilities decrease as age increases.¹⁶

Cardiovascular diseases and cancer had a higher burden of disease than other disease groups in the population with disabilities; this indicates that overall, non-communicable diseases have a higher burden of disease than communicable diseases or injuries in the population with disabilities. Relative importance was markedly different according to age. For example, neurological disorders were most significant in individuals aged 0–19 years, while self-harm and cardiovascular diseases were more significant in individuals in their 20s and 40s, respectively.

Among individual diseases, epilepsy had highest significance in individuals aged 0–19 years, while stroke was the most significant contributor to YLLs in individuals aged ≥ 50 years. This trend is different from the characteristics observed in the general population, where self-harm was the most significant in individuals aged 10–59 years, which may be due to the diseases present in individuals with physical and developmental disabilities. Another marked difference between the population with disabilities and general population is that YLLs due to cancer were consistently high in all age groups except the 0–9-year-old group in the population with disabilities, but the burden of disease from cancer was concentrated in the older age groups in the general population. Such differences indicate that the characteristics of health problems in individuals with disabilities not only differ from the

general population, but also vary among different age groups; therefore, it is necessary to consider age differences when evaluating disease characteristics and intervention policies in individuals with disabilities.

Meanwhile, the ranks of ischemic stroke, ischemic heart disease, and diabetes mellitus were elevated in the population with disabilities, which may be due to their association with underlying diseases in individuals with brain disorders. Similarly, chronic kidney disease from diabetes mellitus (8th) and epilepsy (16th) ranked high, which also may be associated with individuals with kidney impairments and intellectual disabilities. On the other hand, traffic accident-related diseases, such as those involving motorized vehicles with ≥ 3 wheels and pedestrian injuries by road vehicles, in addition to suicide, were lower in the individuals with disabilities than those without. Nevertheless, the absolute values of YLLs in most disease groups, including the abovementioned disease groups, were high in the individuals with disabilities. For example, YLLs per 100,000 people due to self-harm were 893 in the population with disabilities, higher than those in total population (573), alluding to the seriousness of the burden of disease among the individuals with disabilities.

For cause of mortality with each type of disability, ischemic stroke was found to be the leading cause of mortality in individuals with brain disorders, and underlying diseases causing disability were also the leading cause of mortality in people with visual disturbance and epilepsy. Meanwhile, self-harm, including suicide, was the leading cause of mortality in individuals with intellectual disabilities. This was similar to previous findings, where history of mental illnesses, such as depression, bipolar disorder, schizophrenia, and personality disorder, were greatest risk factors for suicide and where risk of suicide was 3–12 times higher in individuals with mental disabilities than in the general population.¹⁶⁻²³ Similarly, we noted that YLLs from suicide were highest for people with mental illnesses.

For people with physical disabilities or hearing impairments, YLLs from ischemic heart disease were higher than those from underlying diseases. This alludes to the poor health management of the individuals with disabilities, necessitating better chronic disease management for such patients. Our findings demonstrated that diseases associated with high YLLs varied with the type of disability, calling for differential health promotion policies according to the type of disability.

Death is a meaningful indicator of the burden of disease.^{10,24} It is also an important indicator of burden of disease in people with disabilities. For example, the number of registered people with disabilities in a given year is calculated by adding the number of new registrations to and subtracting the number of exclusions from the previous year's registration. Thus, the number of people with disabilities is determined by the number of new registrations and exclusions in a given year. Death accounts for the highest proportion of exclusions. Since death accounted for 85.4% of exclusions from the Korea disability registration system in 2014, the number of mortalities greatly influences the size of the population with disabilities. The mortality rate in the individuals with disabilities in Korea is on the rise, increasing from 2.16% in 2004 to 2.82% in 2014, which emphasizes the urgency of reducing the disease burden of premature death in individuals with disabilities to improve their health.

This study has a few limitations. First, we measured the burden of disease for people with disabilities based on YLLs. According to the burden of disease study in Koreans by Yoon et al.,⁹ years lived with disability (YLDs) and YLLs were analyzed and accounted for 88.7%

and 11.3% of the total DALYs in 2012, respectively. Since a significant proportion of DALYs account for YLDs, future studies should also calculate YLDs for people with disabilities. Second, this study only examined those who were registered in the disability registration system and not the entire population with disabilities. However, in Korea, individuals with disabilities who are issued disability certificates from healthcare institutions and who register as those having disabilities are entitled to disability pensions, tax exemption, and medical cost reimbursements, as stipulated by the Welfare Law for the Disabled. The 2014 disability survey revealed that approximately 91.7% of the estimated population with disabilities (as of December 2013) has been registered in the disability system.²⁵ The Korean disability registration data encompasses data from the majority of the population with disabilities.

The burden of disease related to premature death in people with disabilities is larger than that in the general population. This means that the health status of persons with disabilities is disadvantageous compared to the general population. This can be interpreted as the difference in health behaviors such as high smoking rates of the people with disabilities and the difficulty in using medical services like low primary care access. Therefore, in order to improve the health status of people with disabilities, it is necessary to improve accessibility to the system and the introduction of health promotion policies in accordance with the characteristics of persons with disabilities. As shown in this study, since the diseases with high YLL varied according to the type of disability, it is necessary to approach the policy for health promotion and prevention according to the type of disability. For people with disabilities, it is especially necessary to utilize services such as remote monitoring and home healthcare using information and communicating technology, as communication can be difficult or uncomfortable.

It is not known exactly how the risk factors such as smoking and drinking influence YLL. Therefore, future studies should focus on population attributable fractions according to risk factors in the population with disabilities group to better understand the effects of other risk factors on YLL.

REFERENCES

1. World Health Organization. World report on disability. http://www.who.int/disabilities/world_report/2011/en. Updated 2011. Accessed Jun 1, 2018.
2. Ministry of Health and Welfare. *Health and Welfare White Paper*. Sejong: Sejong Print; 2016.
3. National Law Information Center. <http://www.law.go.kr/lsInfoP.do?lsiSeq=178177&efYd=20160630#0000>. Updated 2016. Accessed Jun 1, 2018.
4. Kinne S. Distribution of secondary medical problems, impairments, and participation limitations among adults with disabilities and their relationship to health and other outcomes. *Disabil Health J* 2008;1(1):42-50. [PUBMED](#) | [CROSSREF](#)
5. Hosking FJ, Carey IM, Shah SM, Harris T, DeWilde S, Beighton C, et al. Mortality among adults with intellectual disability in England: comparisons with the general population. *Am J Public Health* 2016;106(8):1483-90. [PUBMED](#) | [CROSSREF](#)
6. Lauer E, McCallion P. Mortality of people with intellectual and developmental disabilities from select US state disability service systems and medical claims data. *J Appl Res Intellect Disabil* 2015;28(5):394-405. [PUBMED](#) | [CROSSREF](#)
7. McCarron M, Carroll R, Kelly C, McCallion P. Mortality rates in the general Irish population compared to those with an intellectual disability from 2003 to 2012. *J Appl Res Intellect Disabil* 2015;28(5):406-13. [PUBMED](#) | [CROSSREF](#)

8. Walker ER, McGee RE, Druss BG. Mortality in mental disorders and global disease burden implications: a systematic review and meta-analysis. *JAMA Psychiatry* 2015;72(4):334-41.
[PUBMED](#) | [CROSSREF](#)
9. Yoon J, Oh IH, Seo H, Kim EJ, Gong YH, Ock M, et al. Disability-adjusted life years for 313 diseases and injuries: the 2012 Korean burden of disease study. *J Korean Med Sci* 2016;31 Suppl 2:S146-57.
[PUBMED](#) | [CROSSREF](#)
10. Lee YR, Kim YA, Park SY, Oh CM, Kim YE, Oh IH. Application of a modified garbage code algorithm to estimate cause-specific mortality and years of life lost in Korea. *J Korean Med Sci* 2016;31 Suppl 2:S121-8.
[PUBMED](#) | [CROSSREF](#)
11. Murray CJ, Lopez AD; World Health Organization; World Bank & Harvard School of Public Health. *The Global Burden of Disease: a Comprehensive Assessment of Mortality and Disability from Diseases, Injuries, and Risk Factors in 1990 and Projected to 2020: Summary*. Geneva, Switzerland: World Health Organization; 1996.
12. Statistics Korea. http://kosis.kr/statHtml/statHtml.do?orgId=101&tblId=DT_1B42&conn_path=I2. Updated 2018. Accessed Jun 1, 2018.
13. Tyrer F, Smith LK, McGrother CW. Mortality in adults with moderate to profound intellectual disability: a population-based study. *J Intellect Disabil Res* 2007;51(Pt 7):520-7.
[PUBMED](#) | [CROSSREF](#)
14. Forsgren L, Edvinsson SO, Nyström L, Blomquist HK. Influence of epilepsy on mortality in mental retardation: an epidemiologic study. *Epilepsia* 1996;37(10):956-63.
[PUBMED](#) | [CROSSREF](#)
15. Patja K, Mölsä P, Iivanainen M. Cause-specific mortality of people with intellectual disability in a population-based, 35-year follow-up study. *J Intellect Disabil Res* 2001;45(Pt 1):30-40.
[PUBMED](#) | [CROSSREF](#)
16. De Wilde EJ, Kienhorst IC, Diekstra RF, Wolters WH. The specificity of psychological characteristics of adolescent suicide attempters. *J Am Acad Child Adolesc Psychiatry* 1993;32(1):51-9.
[PUBMED](#) | [CROSSREF](#)
17. Oquendo MA, Malone KM, Mann JJ. Suicide: risk factors and prevention in refractory major depression. *Depress Anxiety* 1997;5(4):202-11.
[PUBMED](#) | [CROSSREF](#)
18. Harris EC, Barraclough B. Suicide as an outcome for mental disorders. A meta-analysis. *Br J Psychiatry* 1997;170(3):205-28.
[PUBMED](#) | [CROSSREF](#)
19. Qin P, Agerbo E, Westergård-Nielsen N, Eriksson T, Mortensen PB. Gender differences in risk factors for suicide in Denmark. *Br J Psychiatry* 2000;177(6):546-50.
[PUBMED](#) | [CROSSREF](#)
20. Sadock BJ, Sadock VA. *Kaplan & Sadock's Synopsis of Psychiatry: Behavioral Sciences/Clinical Psychiatry*. 9th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2003.
21. Kim W, Jang SY, Chun SY, Lee TH, Han KT, Park EC. Mortality in schizophrenia and other psychoses: data from the South Korea national health insurance cohort, 2002–2013. *J Korean Med Sci* 2017;32(5):835-42.
[PUBMED](#) | [CROSSREF](#)
22. Kim MY, Ha SJ, Jun SS. Factors related to suicidal ideation in patients with chronic schizophrenia, depressive disorder, or bipolar disorder. *J Korean Acad Psychiatr Ment Health Nurs* 2015;24(4):217-25.
[CROSSREF](#)
23. Park S, Lee Y, Youn T, Kim BS, Park JI, Kim H, et al. Association between level of suicide risk, characteristics of suicide attempts, and mental disorders among suicide attempters. *BMC Public Health* 2018;18(1):477.
[PUBMED](#) | [CROSSREF](#)
24. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the global burden of disease study 2010. *Lancet* 2012;380(9859):2095-128.
[PUBMED](#) | [CROSSREF](#)
25. Kim SH, Lee YH, Hwang JH, Oh MA, Lee MK, Lee NH, et al. *2014 Disability Survey*. Sejong, Korea: Ministry of Health and Welfare; 2014.