

Original Article
Obstetrics & Gynecology



Correlation between Morphologic Grading and Euploidy Rates of Blastocysts, and Clinical Outcomes in In Vitro Fertilization Preimplantation Genetic Screening

Min Kyoung Kim ,¹ Jae Kyun Park ,^{1,2} Yunmi Jeon ,¹ Seung-Ah Choe ,³
Hee Jun Lee ,¹ Jayeon Kim ,³ Eun Mi Chang ,¹ Ji Won Kim ,¹ Sang Woo Lyu ,¹
Jin Young Kim ,¹ In Pyung Kwak ,¹ Woo Sik Lee ,¹ and Tae Ki Yoon ,³

¹Department of Obstetrics and Gynecology, Fertility Center of CHA Gangnam Medical Center, CHA University School of Medicine, Seoul, Korea

²Biomedical Science, CHA University, Seongnam, Korea

³CHA Fertility Center, Seoul Station, Seoul, Korea

OPEN ACCESS

Received: May 10, 2018

Accepted: Nov 28, 2018

Address for Correspondence:

Hee Jun Lee, MD, PhD

Division of Reproductive Endocrinology,
Department of Obstetrics and Gynecology,
Fertility Center of CHA Gangnam Medical
Center, CHA University School of Medicine,
569 Nonhyeon-ro, Gangnam-gu, Seoul 06125,
Korea.

E-mail: alonfo@chamc.co.kr

© 2019 The Korean Academy of Medical
Sciences.

This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>)
which permits unrestricted non-commercial
use, distribution, and reproduction in any
medium, provided the original work is properly
cited.

ORCID iDs

Min Kyoung Kim

<https://orcid.org/0000-0002-1753-2674>

Jae Kyun Park

<https://orcid.org/0000-0002-5333-4990>

ABSTRACT






Background: The standard morphological evaluation has been widely used for embryo selection, but it has limitations. This study aimed to investigate the correlation between morphologic grading and euploidy rate of in vitro fertilization (IVF) preimplantation genetic screening (PGS) and compare the pregnancy rates in young and old ages.

Methods: This is a retrospective study using the medical records of patients who underwent IVF procedures with PGS between January 2016 and February 2017 in a single center. The embryo grades were categorized into 4 groups: excellent, good, fair, and poor. Basic characteristics, euploidy rates, clinical pregnancy (CP) rates and ongoing pregnancy rates were analyzed.

Results: The excellent group had significantly higher rate of euploid embryos than fair group (47.82% vs. 29.33%; $P = 0.023$) and poor group (47.82% vs. 29.60%; $P = 0.005$). When the four groups were recategorized into two groups (excellent and good vs. fair and poor), they also showed significant difference in euploidy rates (44.52% vs. 29.53%; $P = 0.002$). When the patients were divided into two groups by age 35, the CP rates for those under and over 35 years old were 44.74% and 47.83%, respectively, which showed no significant difference.

Conclusion: The significant differences among the euploidy rates of different morphologic embryo grades demonstrated the positive correlations between the morphologic grading of the embryo and the euploidy rate of PGS. Additionally, there was no significant difference between the younger and older patients' CP rates. These findings emphasize the fact that old age patients might benefit from PGS whatever the indication of PGS is.

Keywords: PGS; Morphologic Grading; Euploidy Rate; Pregnancy Rate

Yunmi Jeon <https://orcid.org/0000-0003-2751-8718>Seung-Ah Choe <https://orcid.org/0000-0001-6270-5020>Hee Jun Lee <https://orcid.org/0000-0002-2652-9864>Jayeon Kim <https://orcid.org/0000-0001-7167-2304>Eun Mi Chang <https://orcid.org/0000-0001-9256-5057>Ji Won Kim <https://orcid.org/0000-0001-5066-1593>Sang Woo Lyu <https://orcid.org/0000-0001-5572-0898>Jin Young Kim <https://orcid.org/0000-0001-5156-0751>In Pyung Kwak <https://orcid.org/0000-0002-0812-6260>Woo Sik Lee <https://orcid.org/0000-0002-2329-1774>Tae Ki Yoon <https://orcid.org/0000-0001-6314-451X>**Funding**

This work was supported by grant No. 2018R1D1A1B07042308 from the National Research Foundation of Korea.

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Lee HJ. Data curation: Choe SA, Kim J, Chang EM, Kim JW, Lyu SW, Kim JY, Kwak IP, Lee WS, Yoon TK. Formal analysis: Park JK, Lee HJ. Investigation: Park JK, Jeon Y. Methodology: Kim MK, Lee HJ, Park JK. Validation: Kim MK, Lee HJ. Writing - original draft: Kim MK. Writing - review & editing: Kim MK, Lee HJ.

INTRODUCTION

The selection of embryo with best quality for transfer has been the major issue in in vitro fertilization (IVF) era. The standard morphological evaluation has been the most widely used method for embryo selection.¹ However, this method has some limitations. First, it is a very subjective assessment which results may vary among different observers. Second, there may be high variability in embryo scoring since the morphology can vary widely within a few hours.² Therefore, the clinical outcomes of patients with challenging infertility diagnoses such as recurrent pregnancy loss or recurrent implantation failure have not been improved much when their embryos were evaluated solely with morphological evaluations. Another method that may screen embryos with better implantation chances has been needed.

Preimplantation genetic screening (PGS) is derived from preimplantation genetic diagnosis, which was used for testing whether the embryo has the genetically inheritable gender-linked monogenic diseases.³ PGS analyzes all 24 chromosomes by comprehensive chromosomal screening with different techniques such as fluorescent in situ hybridization, array comparative genomic hybridization (aCGH), and next generation sequencing (NGS) to select euploid embryos that will most likely result in pregnancy. It was hypothesized that PGS may improve the pregnancy rates in women with advanced maternal age, recurrent implantation failure, or recurrent miscarriage.⁴⁻⁶

However, PGS also had limitations such as having large numbers of false-positive diagnoses due to mosaicism, disposal of normal euploid embryos, or excluding patients from access to ETs.^{7,8} Because of this limitation, the Preimplantation Genetic International Society defined new diagnostic criteria for PGS.⁹ Technical advancements such as blastocyst trophectoderm (TE) biopsy and NGS are aiming for better accuracy, but PGS is still not yet a perfect study for every patient. Selecting the right indication for PGS is beneficial for saving both time and money required for the challenging infertile patients to get pregnant.

The aims of this study are to find the correlation between morphologic grading and PGS euploid rates, and the indication of PGS according to age. The present study analyzed the relationship between morphological grading and PGS euploidy rates on blastocyst stage, and compared clinical efficiency of PGS among different age groups by calculating the ongoing pregnancy rate to see which category of patients may benefit the most by having PGS done.

METHODS**Patient details**

This retrospective study was performed using the medical records of 136 patients who underwent IVF procedures with PGS on blastocyst stage between January 2016 and February 2017 in Fertility Center of CHA Gangnam Medical Center. PGS on blastocyst stage was offered to infertile patients of advanced age (≥ 35 year; 61.0%), and/or with ≥ 3 unexplained recurrent pregnancy losses previously (27.9%) and/or ≥ 3 recurrent implantation failures previously (47.8%) (Table 1). The exclusion criteria were those who had 3-day embryo biopsied for PGS and those who did not undergo PGS (Fig. 1).




Fig. 1. Flowchart for patient allocation.
ET = embryo transfer.

Table 1. Baseline characteristics of the study participants

Parameters	No.
No. of cycles	136
Age, yr	
Women	35.93 ± 3.70
Men	38.94 ± 4.83
BMI, kg/m ²	20.40 ± 4.55
AMH concentration, ng/mL	4.14 ± 3.86
Antral follicle count, No.	13.47 ± 7.23
Basal FSH concentration, mIU/mL	5.30 ± 3.05
Basal LH concentration, mIU/mL	4.99 ± 2.58
Basal E2 concentration, pg/mL	54.38 ± 31.18
E2 concentration on hCG day, pg/mL	2,673.26 ± 1,701.83
Progesterone concentration on hCG day, ng/mL	0.85 ± 0.57
LH concentration on hCG day, mIU/mL	3.07 ± 4.59
Fertilization rate	1,577/2,194 (71.88)
Conventionally fertilized oocytes	292/369 (79.13)
ICSI fertilized oocytes	1,096/1,555 (70.48)
Half ICSI fertilized oocytes	189/270 (70.00)
Blastocyst formation rate	676/1,549 (43.60)
PGS indication	136
Advanced age (≥ 35 year)	83 (61.03)
≥ 3 unexplained recurrent pregnancy losses	38 (27.94)
≥ 3 recurrent implantation failures	65 (47.79)
Abortus chromosome abnormality	22 (16.18)
Others	11 (8.09)
Blastocysts analyzed for PGS	444
Euploid blastocysts	153 (34.46)
Aneuploid blastocysts	291 (65.54)
Fresh ET	20
Frozen ET	64

Values are presented as mean ± standard deviation (SD) or number (%).

BMI = body mass index, AMH = anti-Mullerian hormone, FSH = follicle stimulating hormone, LH = luteinizing hormone, E2 = estradiol, hCG = human chorionadotropin, ICSI = intracytoplasmic sperm injection, PGS = preimplantation genetic screening, ET = embryo transfer.

Clinical protocols

Gonadotropin-releasing hormone (GnRH) antagonist and agonist protocols were used in this study. Gonadotropin doses were formulated according to the patient's antral follicular count, antimüllerian hormone, and previous response to stimulation. Transvaginal ultrasound was performed to monitor follicular response to stimulation and gonadotropin doses were adjusted accordingly. Final oocyte maturation was triggered with hCG (Ovidrel; Merck, Kenilworth, NJ, USA) with or without GnRH agonist (Superfact; Sanofi-Aventis, Paris, France) when the mean diameter of ≥ 2 follicles sized ≥ 17 mm. Ultrasound-guided oocyte-retrieval under conscious sedation was performed 34–36 hours after the trigger.

Luteal phase support was provided either with progesterone vaginal suppositories, such as Utrogestan (Besins Healthcare, Paris, France), Lutinus (Ferring, Saint-Prex, Switzerland), Crinone gel 8% (Merck) or intramuscular injections (Sugest; Sanzyme Ltd., Banjara Hills, India) starting on the day of fresh ET. For frozen-thawed ET, luteal phase support was given starting from 5 days before the ET.

Embryo morphologic grading and biopsy

Blastocysts were graded according to the degree of expansion, inner cell mass (ICM) and TE morphology. The blastocoel expansion grading system is as follows: Early, the blastocoel filling $< 50\%$ of the nonexpanded embryo; Mid, the blastocoel filling $> 50\%$ of the embryo; Expanded, full blastocyst, cavity completely filling the embryo, or expanded blastocyst, cavity larger than the embryo with a thin zona pellucida; Hatching, a hatching blastocyst; Hatched, a blastocyst that has completely hatched out of the zona pellucida. The ICM grading was as follows: A, many, tightly packed cells; B, loosely grouped several cells; C, very few cells. Similarly TE grading was A, many cells creating cohesive epithelium; B, few cells forming loose epithelium; C, very few large cells.¹⁰ The embryo grades were categorized into 4 groups according to their morphologic grading before TE biopsy: Excellent (E; Expanded AA, Hatching AA, Hatched AA), Good (G; Early AA, Mid AA, Expanded AB, BA, Hatching AB, BA, Hatched AB, BA), Fair (F; Early AB, BA, Mid AB, AC, BA, BB, Expanded AC, BB, CA, Hatching AC, BB, CA, Hatched AC, BB, CA), and Poor (P; Early AC, BB, BC, CA, CB, CC, Mid BC, CA, CB CC, Expanded BC, CB, CC, hatching BC, CB CC, Hatched BC, CB CC).

All embryos were biopsied on day 5, during their blastocyst stages. TE biopsy was done from blastocysts that were cultured after conventional IVF or intracytoplasmic sperm injection (ICSI) using a pair of micromanipulators (NT-88; Narishige, Tokyo, Japan) with microscope (TE-2000; Nikon, Tokyo, Japan). The culture media consisted of G-PGD (Vitrolife, Göteborg, Sweden) with 5% HSA-solution (Vitrolife). For TE biopsy, the embryos were immobilized with a holding pipette (left side), and the zona pellucida was perforated using laser pulses ZILOS-tk noncontact laser (Hamilton Thorne Biosciences, Beverly, MA, USA). Using the right side biopsy pipette, gentle suction of TE cells was done through the zona pellucida perforation. The cells were eventually separated out of the zona pellucida (**Fig. 2**). The biopsied TE cells were washed with D-phosphate-buffered saline (D-PBS; Ca^{2+} and Mg^{2+} -removed) and contained in RNase–DNase-free 0.2 mL polymerase chain reaction (PCR) tubes for PGS referral to MG-MED Inc. (Seoul, Korea). The biopsied blastocysts were then cultured in sterile oil-covered media at 37°C , $6\% \text{CO}_2$, $6\% \text{O}_2$, and observations at 24–48 hour-interval was done to check development of the embryos.




Fig. 2. The process of blastocyst biopsy. **(A)** Expanded blastocyst (magnification $\times 200$). **(B)** TE cells were separated by biopsy pipette (magnification $\times 400$). Arrow indicates biopsied TE cells. **(C)** Shrunken blastocyst appears after TE biopsy (magnification $\times 400$). **(D)** Following further incubation for 24 hours after biopsy, the blastocyst undergoes a process of expansion and hatching from the zona pellucida (magnification $\times 200$). TE = trophectoderm.

PGS (aCGH) protocol

The biopsied cells were washed and collected into a PCR tube with 2.5 μL of PBS. Extraction and amplification of the DNA from the biopsied cells were performed according to the MG Flex single cell whole genome amplification kit system protocol (MG Flex; MGMED, Seoul, Korea). Approximately 3 μg of amplified DNA was used in the aCGH experiments (MGMED). Briefly, the amplified DNA was labeled with Cy-3 and Cy-5 dCTP for 3 hours using a random priming method. The labeled DNA was purified, dissolved in hybridization buffer, and hybridized overnight. The slides were washed several times and dried. Images of the slides were acquired with a GenePix4000B dual-laser scanner (Axon Instruments, Union City, CA, USA) and analyzed with MG Viewer analysis software (MGMED).

Statistical analysis

Continuous data are expressed as mean \pm standard deviation, and categorical variables are in absolute and percentage frequency. Pearson's χ^2 test, Fisher's exact test, student's t-test, and Mann-Whitney test were used for statistical analysis. A P value < 0.05 was considered statistically significant and the statistical analysis was performed with SPSS ver.23 (IBM, Armonk, NY, USA).

Ethics statement

The present study protocol was reviewed and approved by the Institutional Review Board (IRB) of CHA Gangnam Medical Center (IRB No. GCI-17-24). Consent to participate was waived due to its retrospective nature and medical records were only used in this analysis.

RESULTS

Baseline characteristics of the patients are described in **Table 1**. The number of developed blastocysts is 676 embryos in 136 cycles and actual number of embryos that underwent PGS is 444 embryos. PGS was done on all blastocysts that have developed in fresh ET cycles, but in frozen ET cycles, mostly only 2–3 embryos (maximum 5) per cycle were thawed and underwent PGS. Therefore, other 232 embryos were left frozen without PGS. For this reason, the number of developed blastocysts (676 embryos) and actual number of embryos that underwent PGS (444 embryos) is different. Total of 136 cycles which contain 444 PGS-blastocysts were included (**Fig. 1**). Among the total 444 blastocysts, there were 153 euploid blastocysts that were analyzed by PGS (34.46%) in **Table 1**. The euploid embryos were categorized into 4 groups according to their morphological grades and the euploidy rates are as follows: E (n = 33/69; 47.82%), G (n = 32/77; 41.55%), F (n = 22/75; 29.33%), and P (n = 66/223; 29.60%) in **Table 2** and **Fig. 3**. The E group had significantly higher rate of euploid embryos than group F ($P = 0.023$) and group P ($P < 0.01$) in **Table 2** and **Fig. 3**. However, the difference between groups E and G was without statistical significance ($P = 0.448$) in **Table 2** and **Fig. 3**. When the four groups were recategorized into two groups (E and G vs. F and P), they also showed significant difference in euploidy rates ($P = 0.002$) (**Fig. 3**).

In **Table 2**, when the embryos were divided by their degrees of blastocoel expansion, the euploidy rates showed a tendency to increase as the level of blastocoel expansion progressed, except the hatched group. However, there was no significant difference when compared to hatched degree since the number of hatched embryos were too small for statistical analysis. Therefore, we only compared the hatching group with Expanded, Mid and Early groups, excluding the hatched group in **Table 2**. ICM grades showed significant difference for the euploidy rates between A grade and C grade ($P = 0.022$). The TE grades were also significantly different between A grade and lower quality grade such as B and C ($P < 0.05$).

Only total of 84 cycles that have euploid embryos proceeded with ET and remaining 52 out of 136 cycles did not undergo ET due to no euploid embryos available (**Fig. 1**). When the

Table 2. Correlation between the human blastocyst morphological evaluations and their euploidy rates

Variables	Embryos (euploidy rate), No. (%)	P value
Blastocyst	All (n = 153/444)	vs. Excellent
Excellent	33/69 (47.82)	-
Good	32/77 (41.55)	NS
Fair	22/75 (29.33)	0.023
Poor	66/223 (29.60)	< 0.01
Expansion of the blastocoel	All (n = 153/444)	vs. Hatching
Hatched	1/4 (25.00)	NS
Hatching	13/22 (59.09)	-
Expanded	59/163 (36.19)	0.039
Mid	52/143 (36.36)	0.043
Early	28/112 (25.00)	< 0.01
ICM grade	All (n = 153/444)	vs. A
A (many, tightly packed cells)	84/212 (39.62)	-
B (loosely grouped several cells)	7/19 (36.84)	NS
C (very few cells)	62/213 (29.11)	0.022
TE grade	All (n = 153/444)	vs. A
A (many cells creating cohesive epithelium)	55/117 (47.01)	-
B (few cells forming loose epithelium)	35/118 (29.66)	0.014
C (very few large cells)	63/209 (30.14)	< 0.01

Values are presented as number (%).

NS = not specific, ICM = inner cell mass, TE = trophectoderm.




Fig. 3. Correlations between morphologic grades and euploidy rates. The excellent group had significantly higher rate of euploid embryos than group fair ($P = 0.023$) and group poor ($P = 0.005$). However, the difference between groups excellent and good was without statistical significance ($P = 0.448$). When the four groups were recategorized into two groups (excellent and good vs. fair and poor), they also showed significant difference in euploidy rates ($P = 0.002$).

^a $P < 0.01$; ^b $P < 0.05$.

84 cycles were divided into two groups by age 35 (**Table 3**), the clinical pregnancy (CP) rate for those under 35 was 44.74% and 47.83% for those over 35 years old, which showed no significant difference. The implantation rate and ongoing pregnancy rate also showed no significant difference between the age groups. Miscarriages were noted only from the older age group (31.82%) and none from those under 35 years old.

Multiple logistic regression analysis was done to find out which is the most influential morphologic grading factor that impacts euploidy status of the embryo (**Table 4**). Among the three grading factors, the blastocoel expansion grade was the most influential (odds ratio, 1.261; $P = 0.045$).

DISCUSSION

To summarize, the present study described the correlations between blastocyst morphologic grades and their euploidy rates. As the morphologic grades get better, the euploidy rates also increase. Among the morphologic grading criteria, all three parameters, such as the blastocoel expansion grade, ICM grade, and TE grade were relevant to euploidy rates. The embryos express their own genes after the third day of fertilization and their potential genetic abnormalities will result in embryo maldevelopment, which causes the lower morphologic grade of embryos.^{11,12} Lastly, the different age groups by age 35 did not show much difference in CP and ongoing pregnancy rates, suggesting that the PGS may be more useful for older women, over 35 years old with poorer chances of pregnancy.

Assessing the embryo morphology has been the classic tool for selecting the embryos for intra-uterine transfer. However, not all embryos with the best morphologic grade match the

Table 3. Comparisons of IVF outcomes between young and old age groups after PGS (Total of 84 cycles)

Variables	< 35 years old (n = 38)	≥ 35 years old (n = 46)	P value
Age, yr			
Women	32.05 ± 0.29	37.89 ± 0.38	< 0.01
Men	35.82 ± 0.50	41.02 ± 0.67	< 0.01
BMI, kg/m ²	21.00 ± 0.80	21.02 ± 0.64	NS
AMH concentration, ng/mL	3.89 ± 0.63	4.18 ± 0.55	NS
Antral follicle count	13.75 ± 2.36	8.33 ± 1.49	NS
Basal FSH concentration, mIU/mL	4.42 ± 0.48	5.80 ± 0.45	0.039
Basal LH concentration, mIU/mL	5.01 ± 0.68	3.67 ± 0.81	NS
Basal E2 concentration, pg/mL	54.74 ± 5.02	47.52 ± 6.30	NS
E2 concentration on hCG day, pg/mL	2,836.84 ± 392.14	1,612.83 ± 254.45	0.013
Progesterone concentration on hCG day, ng/mL	0.91 ± 0.13	0.80 ± 0.22	NS
LH concentration on hCG day, mIU/mL	5.54 ± 1.69	3.80 ± 2.13	NS
Blastocyst			
Excellent	12/17 (70.59)	14/24 (58.33)	NS
Good	12/25 (48.00)	12/29 (41.38)	NS
Fair	12/31 (38.71)	7/21 (33.33)	NS
Poor	27/63 (45.86)	29/77 (37.66)	NS
Expansion of the blastocyst			
Hatched	0/1 (0)	9/32 (28.13)	NS
Hatching	7/11 (63.64)	23/58 (39.66)	NS
Expanded	22/43 (51.16)	26/56 (46.43)	NS
Mid	21/46 (45.65)	3/4 (75.00)	NS
Early	13/35 (37.14)	1/1 (100.00)	NS
ICM grade			
A (many, tightly packed cells)	34/68 (50.00)	33/72 (45.83)	NS
B (loosely grouped several cells)	3/7 (42.86)	2/6 (33.33)	NS
C (very few cells)	26/61 (42.62)	27/73 (36.99)	NS
TE grade			
A (many cells creating cohesive epithelium)	18/31 (58.06)	25/47 (53.19)	NS
B (few cells forming loose epithelium)	19/45 (42.22)	8/31 (25.81)	NS
C (very few large cells)	26/60 (43.33)	29/73 (39.73)	NS
Fertilization rate	474/677 (70.01)	554/730 (75.89)	NS
Conventionally fertilized oocytes	104/142 (73.24)	145/167 (86.83)	NS
ICSI fertilized oocytes	250/357 (70.03)	350/485 (72.16)	NS
Half ICSI fertilized oocytes	120/178 (67.42)	59/78 (75.64)	NS
Euploidy rate	46.32 ± 0.04 (63/136)	41.06 ± 0.04 (62/151)	NS
Embryos transferred	1.34 ± 0.08	1.20 ± 0.06	NS
Implantation rate	18/51 (35.29)	25/55 (45.45)	NS
CP rate	17/38 (44.74)	22/46 (47.83)	NS
Miscarriage rate	0	7/22 (31.82)	0.015
Ongoing pregnancy rate	17/38 (44.74)	15/46 (32.61)	NS

Values are presented as mean ± standard deviation (SD) or number (%).

IVF = in vitro fertilization, PGS = preimplantation genetic screening, BMI = body mass index, NS = not significant, AMH = anti-Müllerian hormone, FSH = follicle stimulating hormone, LH = luteinizing hormone, E2 = estradiol, hCG = human chorionic gonadotropin, ICM = inner cell mass, TE = trophectoderm, ICSI = intracytoplasmic sperm injection, CP = clinical pregnancy.

Table 4. Multiple logistic regression analysis of the correlation between morphologic grading and euploidy status

Variables	Euploidy status	
	OR (95% CI)	P value
Morphologic grading	1.314 (1.109–1.557)	< 0.01
Expansion	1.261 (1.005–1.582)	0.045
ICM	1.003 (0.654–1.537)	0.989
TE	1.335 (0.809–2.205)	0.258

OR = odds ratio, CI = confidence interval, ICM = inner cell mass, TE = trophectoderm.

euploid status as this study demonstrated about 52% of excellent blastocysts being aneuploid and in contrast, about 30% of poor grades being euploid. Previous studies also showed that several aneuploid embryos scored high morphologically.^{11,13,14} This emphasizes that it may

be hazardous to select embryos solely by morphological assessment and underlines the importance of utilizing PGS for best embryo selection.

PGS methods have evolved continuously throughout the past decades. To mention the aspect of biopsy methods, cleavage stage biopsy was first utilized. However, Fragouli et al.¹² demonstrated that PGS biopsy during cleavage stage showed high rates of aneuploidy even in reproductively younger women (73% for average 33.6 years of women's age). Other studies also reported similar results.¹⁵⁻¹⁷ This phenomenon might be attributed to the sperm-derived men's meiotic errors, oocyte abnormalities and post-zygotic chromosome malsegregation during mitosis.¹⁸ On the other hand, blastocyst biopsy has improved the chances of selecting euploid embryos among the top morphology-scored embryos.¹² TE biopsy has been the preferred method due to its extraembryonic origin rather than ICM which develops into embryo itself. It is also known to predict the ICM karyotype status without impairing the embryonic implantation.¹⁹ Alfarawati et al.¹¹ described that declining TE grades was correlated with increasing aneuploidy rates. Furthermore, the study by Minasi et al.²⁰ reported that euploid blastocysts showed a higher percentage of top grade ICM and TE, higher blastocoel expansion grades and shorter time to beginning of blastulation, expansion, and hatching, compared to aneuploid blastocysts. Another recent study demonstrated that embryos with higher grade of blastocyst expansion, ICM and TE had greater chances of being euploid.²¹ All these data underlines our result showing correlations between blastocyst morphology and euploidy rates on blastocyst stage.

PGS has been debated in terms of clinical efficiency. Not only it is imperfect as errors may occur, it may also burden patients with extra costs. Moreover, the incidence of mitotic mosaicism is not known, which can result in sampling errors. Therefore, it is unavoidable that some of normal embryos will be thrown away. Paulson²² presented an estimation of normal embryos lost due to PGS as high as 40% and 50% of the remaining embryos achieving implantation. A recent study also demonstrated that PGS for women under 40 years of age at their first IVF attempt is unlikely to be beneficial for most women.²³ But, on the contrary, Harton et al.²⁴ showed that implantation and pregnancy rates were not significantly different between reproductively younger and older patients by selective transfer of euploid embryos, and also describes that chromosome screening was able to decrease the maternal age effect in pregnancy. Since older patients had greater risks of developing aneuploid embryos, PGS sorting out euploid embryos for transfer may significantly increase the chances of pregnancy. Another study by Kang et al.²⁵ demonstrates that only the women older than 37 years old benefitted from PGS, having improved implantation rates and live birth rates. Therefore, PGS might be useful to improve the IVF outcomes in older aged patients, which is similar to the result of this study.

There are some limitations in this study. The selection bias might have affected our results. In **Table 3**, the miscarriage rate in old age group (≥ 35 years old) is 31.82%, which is rather high. But, the unexpected high miscarriage rate could be attributed to selection bias. Significant number of patients with history of recurrent pregnancy loss might be included in this study. Also, the retrospective nature of this study also remains as a limitation. The present study's patient group with over 35 years of age had relatively good ovarian reserve (mean level of anti-Mullerian hormone [AMH], 4.18 ng/mL in **Table 3**), and it is probably because patients aged 35 or older who can produce more number of embryos are more feasible to undergo PGS. Many patients over 35 years of age who have decreased ovarian reserve undergo IVF treatment. In order to perform PGS for patients over 35 years of age

who have decreased ovarian reserve, we need to collect their small number of embryos several times using embryo cryopreservation, and better start PGS when there are enough embryos to be evaluated. These procedures take much longer time and were not appropriate for retrospective short term follow-up study like this. Further prospective long-term studies including more number of patients with decreased ovarian reserve should be performed to analyze the cost-effectiveness of PGS for older patients.

In conclusion, the different age groups by age 35 did not show much difference in CP rates and ongoing pregnancy rates, suggesting that the PGS may be more useful for older women who are over 35 years old with poorer chances of pregnancy.

REFERENCES

- Gardner DK, Sakkas D. Assessment of embryo viability: the ability to select a single embryo for transfer --a review. *Placenta* 2003;24 Suppl B:S5-12.
[PUBMED](#) | [CROSSREF](#)
- Gardner DK, Meseguer M, Rubio C, Treff NR. Diagnosis of human preimplantation embryo viability. *Hum Reprod Update* 2015;21(6):727-47.
[PUBMED](#) | [CROSSREF](#)
- Handyside AH, Kontogianni EH, Hardy K, Winston RM. Pregnancies from biopsied human preimplantation embryos sexed by Y-specific DNA amplification. *Nature* 1990;344(6268):768-70.
[PUBMED](#) | [CROSSREF](#)
- Greco E, Bono S, Ruberti A, Lobascio AM, Greco P, Biricik A, et al. Comparative genomic hybridization selection of blastocysts for repeated implantation failure treatment: a pilot study. *BioMed Res Int* 2014;2014:457913.
[PUBMED](#) | [CROSSREF](#)
- Rubio C, Simón C, Vidal F, Rodrigo L, Pehlivan T, Remohí J, et al. Chromosomal abnormalities and embryo development in recurrent miscarriage couples. *Hum Reprod* 2003;18(1):182-8.
[PUBMED](#) | [CROSSREF](#)
- Kahraman S, Bahçe M, Samli H, Imirzalioglu N, Yakisn K, Cengiz G, et al. Healthy births and ongoing pregnancies obtained by preimplantation genetic diagnosis in patients with advanced maternal age and recurrent implantation failure. *Hum Reprod* 2000;15(9):2003-7.
[PUBMED](#) | [CROSSREF](#)
- Orvieto R, Shuly Y, Brengauz M, Feldman B. Should pre-implantation genetic screening be implemented to routine clinical practice? *Gynecol Endocrinol* 2016;32(6):506-8.
[PUBMED](#) | [CROSSREF](#)
- Gleicher N, Vidali A, Braverman J, Kushnir VA, Barad DH, Hudson C, et al. Accuracy of preimplantation genetic screening (PGS) is compromised by degree of mosaicism of human embryos. *Reprod Biol Endocrinol* 2016;14(1):54.
[PUBMED](#) | [CROSSREF](#)
- Preimplantation Genetic Diagnosis International Society. PGDIS position statement on chromosome mosaicism and preimplantation aneuploidy testing at the blastocyst stage. http://www.pgdis.org/docs/newsletter_071816.html. Updated 2016. Accessed July 19, 2016.
- Gardner DK, Weissman A, Howles CM, Shoham Z. *Textbook of Assisted Reproductive Techniques: Laboratory Perspectives*. 4th ed. London: Informa Healthcare; 2012.
- Alfarawati S, Fragouli E, Colls P, Stevens J, Gutiérrez-Mateo C, Schoolcraft WB, et al. The relationship between blastocyst morphology, chromosomal abnormality, and embryo gender. *Fertil Steril* 2011;95(2):520-4.
[PUBMED](#) | [CROSSREF](#)
- Fragouli E, Alfarawati S, Spath K, Wells D. Morphological and cytogenetic assessment of cleavage and blastocyst stage embryos. *Mol Hum Reprod* 2014;20(2):117-26.
[PUBMED](#) | [CROSSREF](#)
- Munné S. Chromosome abnormalities and their relationship to morphology and development of human embryos. *Reprod Biomed Online* 2006;12(2):234-53.
[PUBMED](#) | [CROSSREF](#)

14. Hardarson T, Caisander G, Sjögren A, Hanson C, Hamberger L, Lundin K. A morphological and chromosomal study of blastocysts developing from morphologically suboptimal human pre-embryos compared with control blastocysts. *Hum Reprod* 2003;18(2):399-407.
[PUBMED](#) | [CROSSREF](#)
15. Gutiérrez-Mateo C, Colls P, Sánchez-García J, Escudero T, Prates R, Ketterson K, et al. Validation of microarray comparative genomic hybridization for comprehensive chromosome analysis of embryos. *Fertil Steril* 2011;95(3):953-8.
[PUBMED](#) | [CROSSREF](#)
16. Northrop LE, Treff NR, Levy B, Scott RT Jr. SNP microarray-based 24 chromosome aneuploidy screening demonstrates that cleavage-stage FISH poorly predicts aneuploidy in embryos that develop to morphologically normal blastocysts. *Mol Hum Reprod* 2010;16(8):590-600.
[PUBMED](#) | [CROSSREF](#)
17. Voullaire L, Slater H, Williamson R, Wilton L. Chromosome analysis of blastomeres from human embryos by using comparative genomic hybridization. *Hum Genet* 2000;106(2):210-7.
[PUBMED](#) | [CROSSREF](#)
18. Fragouli E, Alfarawati S, Spath K, Jaroudi S, Sarasa J, Enciso M, et al. The origin and impact of embryonic aneuploidy. *Hum Genet* 2013;132(9):1001-13.
[PUBMED](#) | [CROSSREF](#)
19. Johnson DS, Cinnioglu C, Ross R, Filby A, Gemelos G, Hill M, et al. Comprehensive analysis of karyotypic mosaicism between trophectoderm and inner cell mass. *Mol Hum Reprod* 2010;16(12):944-9.
[PUBMED](#) | [CROSSREF](#)
20. Minasi MG, Colasante A, Riccio T, Ruberti A, Casciani V, Scarselli F, et al. Correlation between aneuploidy, standard morphology evaluation and morphokinetic development in 1730 biopsied blastocysts: a consecutive case series study. *Hum Reprod* 2016;31(10):2245-54.
[PUBMED](#) | [CROSSREF](#)
21. Barash OO, Ivani KA, Willman SP, Rosenbluth EM, Wachs DS, Hinckley MD, et al. Association between growth dynamics, morphological parameters, the chromosomal status of the blastocysts, and clinical outcomes in IVF PGS cycles with single embryo transfer. *J Assist Reprod Genet* 2017;34(8):1007-16.
[PUBMED](#) | [CROSSREF](#)
22. Paulson RJ. Preimplantation genetic screening: what is the clinical efficiency? *Fertil Steril* 2017;108(2):228-30.
[PUBMED](#) | [CROSSREF](#)
23. Scriven PN. Towards a better understanding of preimplantation genetic screening for aneuploidy: insights from a virtual trial for women under the age of 40 when transferring embryos one at a time. *Reprod Biol Endocrinol* 2017;15(1):49.
[PUBMED](#) | [CROSSREF](#)
24. Harton GL, Munné S, Surrey M, Grifo J, Kaplan B, McCulloh DH, et al. Diminished effect of maternal age on implantation after preimplantation genetic diagnosis with array comparative genomic hybridization. *Fertil Steril* 2013;100(6):1695-703.
[PUBMED](#) | [CROSSREF](#)
25. Kang HJ, Melnick AP, Stewart JD, Xu K, Rosenwaks Z. Preimplantation genetic screening: who benefits? *Fertil Steril* 2016;106(3):597-602.
[PUBMED](#) | [CROSSREF](#)