

The Prevalence and Associated Risk Factors of Renal Artery Stenosis in Patients Undergoing Cardiac Catheterization

Hyun Yong Song¹, Jae Ha Hwang¹, Hyunjin Noh¹, Sug Kyun Shin¹, Dong Hoon Choi², Won Hum Shim², Ho Yung Lee¹, Seung Yun Cho², Dae Suk Han¹, and Kyu Hun Choi¹

Abstract

Renal artery stenosis may be a cause of hypertension and a potential contributor to progressive renal insufficiency. However, the prevalence of renal artery disease in a general population is poorly defined. The purposes of this study were to evaluate the prevalence of angiographically-determined renal artery narrowing in a patient population undergoing routine cardiac catheterization, and to identify the risk factors for renal artery stenosis. After left ventriculography, abdominal aortography was performed to screen for the presence of renal artery stenosis. A total of 427 patients (274 males, 153 females) were studied and the mean age was 59 years. Renal artery narrowing was identified in 10.5% of patients. Significant ($\geq 50\%$ diameter narrowing) renal artery stenosis was found in 24 patients (5.6%) and insignificant stenosis was found in 1.4%. The stem of the renal artery was a more common site of stenosis in 62.2% of patients than in the ostium (37.8%), but the severity of stenosis was not significantly different according to the site of stenosis. By univariate and multivariate logistic regression analysis, the association of clinical variables with renal artery stenosis was assessed. Multivariable predictors included age, hypertension and peripheral vascular disease ($p < 0.05$). The variables such as sex, smoking history, hyperlipidemia, renal insufficiency, as well as the presence of obesity, severity of coronary heart disease and D.M., were not associated. In conclusion, the prevalence of angiographically-determined renal artery narrowing in a patient population undergoing cardiac catheterization is 10.5%. Old age, hypertension and evidence of peripheral vascular disease represent the predictors of renal artery stenosis.

Key Words: Renal artery stenosis, prevalence, risk factor, cardiac catheterization

INTRODUCTION

Renovascular disease as a cause of end-stage renal disease has been mostly overlooked until recently.¹ As the overall population ages, the number of individuals with atherosclerotic renovascular disease will increase, as will the number of patients with renovascular hypertension and end-stage renal disease. The anatomic presence of atherosclerotic renovascular disease as a threat to renal function has become a more pressing concern than that of hypertension secondary to renovascular disease.^{2,3} A review of comorbid atheros-

clerotic disease in end-stage renal disease patients indicated that occlusive disease of the renal arteries may contribute to progressive renal failure.¹

Stenosis of the renal arteries usually results from atheromatous lesions, and this prevalence then increases with age and with atherosclerosis at extrarenal sites.⁴ Renal artery stenosis (RAS) is a potentially correctable problem because revascularization techniques including surgery and percutaneous renal artery angioplasty have been shown to be effective in treating RAS and in preserving renal function.^{3,5} However, the prevalence and predictors of renal artery disease are poorly defined. The population of patients referred for cardiac catheterization exhibits many clinical features that could be associated with atherosclerotic renal artery disease. Thus, abdominal aortography at the time of cardiac catheterization provides a readily performed means of screening for the presence of RAS in a population likely to have renal artery disease.

Received September 22, 1999

Accepted December 18, 1999

Departments of ¹Internal Medicine and ²Cardiovascular Center, Institute of Kidney Disease, Yonsei University College of Medicine, Seoul, Korea.

Address reprint request to Dr. K. H. Choi, Department of Internal Medicine, Yonsei University College of Medicine, C.P.O. Box 8044, Seoul 120-752, Korea. Tel: 82-2-361-5437, Fax: 82-2-393-6884, E-mail: khchoi6@yumc.yonsei.ac.kr

To evaluate the prevalence of angiographically-determined renal artery narrowing in a patient population undergoing routine cardiac catheterization and to identify the risk factors for RAS, we reviewed the medical records and angiographic findings in this population.

MATERIALS AND METHODS

Patient population

Four hundred and twenty seven consecutive patients undergoing cardiac catheterization were screened for the presence of renal artery disease between July 1998 and December 1998 through the use of abdominal aortography. Before the patients had cardiac catheterization, demographic data, medical history, physical findings, and blood chemistries were recorded. Peripheral vascular disease was defined as a history of claudication, previous vascular procedure, a history of stroke or transient ischemic attack, or physical evidence of carotid, femoral, or abdominal bruits. Congestive heart failure was diagnosed by echocardiography or ventriculography. Hypertension was defined as diastolic blood pressure >90 mmHg when subjects were measured as outpatients before catheterization, or while the patient was taking antihypertensive medications. Renal insufficiency defined as a creatinine level was above 1.5 mg/dL.

Angiographic analysis

The cardiac catheterization was performed via the femoral artery approach known as Judkins' technique. The presence and severity of coronary atherosclerotic lesions were determined in a routine fashion. Coronary artery lesions graded as $>70\%$ narrowing of the luminal diameter were classified as significant. After left ventriculography, the pigtail catheter was withdrawn into the abdominal aorta and positioned a few centimeters superior to the renal arteries. Aortography was performed in the anterior-posterior projection with ioxaglate de sodium (Guerbet S.A., Paris, France) power-injected at a rate of 20 mL/s to a total volume of 30 mL. The injection was recorded on 35 mm cine film or compact disc at 30 frames per second.

Aortograms were reviewed by two cardiologists blinded to the clinical information. The adequacy of the study as well as the presence or absence of renal artery disease were noted. The percentage of RAS was classified as a minor irregularity ($<25\%$ narrowing) or as 25, 50, 75 and 100% luminal diameter narrowing. By convention, an angiographically-significant lesion was defined as a $\geq 50\%$ luminal diameter narrowing of a major renal artery. Selective arteriography with a 5 F renal catheter was done if the renal artery lesion was significantly narrowed. Accessory renal arteries with disease were felt to be significant if more than one-third of the renal mass was estimated to be supplied by the vessel. Lesion location was classified as ostial, main artery, or branch vessel. Ostial lesions were defined as stenotic if the segment of the renal artery lumen immediately contiguous to the aorta was compromised. Lesions of the main segment of the renal artery began at least 2 to 3 mm beyond the ostial segment. Branch lesions were defined as stenotic lesions originating beyond the first bifurcation of the renal artery.

Statistical analysis

Statistical analysis was performed using the Chi-square test, stepwise univariate and multivariable logistic regression in order to identify the clinical and angiographic variables predictive of RAS. The variables that were significant in the univariate model were then entered into a multivariate logistic regression model to identify the best set of independent predictors of RAS. Data were expressed as mean \pm S.D. Statistical significance was defined as $p < 0.05$.

RESULTS

Patient characteristics

The demographic data from the study patients are summarized in Table 1. The mean age of the group was 59.2 ± 10.3 yr. About half of the patients had a history of smoking. Of 427 patients, 216 (51%) had hypertension, and 92 (22%) had diabetes. Peripheral vascular disease and congestive heart failure were observed in 75 (18%) and 74 (17%) patients, respectively. Angiographically-significant coronary artery

Table 1. Demographic Data

| | N=427 (%) |
|-----------------------------|------------------|
| Mean age, yr | 59.2 ± 10.3 |
| Men : Women | 284 : 143 |
| Smoking | 218 (51) |
| % ideal body weight | 102.6 ± 19.8 |
| Hypertension | 216 (51) |
| Diabetes Mellitus | 92 (22) |
| Peripheral vascular disease | 75 (18) |
| Congestive heart failure | 74 (17) |
| Coronary artery disease | |
| 1 Vessel | 131 (31) |
| multi-vessel | 171 (40) |
| Serum creatinine, mg/dL | |
| Mean (range) | 1.19 (0.40–4.70) |
| ≥ 1.5 mg/dL | 36 (8.4) |

disease was noted in 302 (70.7%) patients.

Prevalence and location of RAS

The prevalence of RAS in this population is illustrated in Fig. 1. Bilaterally normal renal arteries were identified in 89.5% of patients screened. Insignificant RAS of at least one renal artery was seen in 21 (4.9%) patients. Significant unilateral RAS was identified in 18 (4.2%) patients, and 6 (1.4%) had significant bilateral RAS. Thus, 45 (10.5%) patients in the study group had angiographically RAS observed at the time of cardiac catheterization. All lesions were designated as being atherosclerotic. The majority of lesions were located in the ostium or the main segment of the renal artery (Fig. 2). The stem of the renal artery was a more common site of stenosis in 62.2% than the ostium in 37.8% ($p < 0.05$), but the severity of stenosis was not significantly different according to the site of stenosis.

Predictors of RAS

By univariate logistic regression, the clinical and biochemical variables were analyzed to identify associations that were more common in patients with RAS. The predictive clinical variables (shown in descending order according to χ^2 values) are summarized in Table 2. The evidence of peripheral vascular disease was the most important predictor of RAS. Next in order of importance was age and then hypertension. The mean age of patients with RAS was




Fig. 1. The prevalence of renal artery stenosis (RAS) (N=427). Insignificant RAS is defined as <50% luminal narrowing, and significant RAS is defined as ≥50% luminal narrowing.




Fig. 2. Location and distribution of renal artery stenosis. The stem of renal artery was more common site of stenosis in 62.2% than the ostium in 37.8%. Definitions of insignificant and significant lesions were same as Fig. 1.

63.2 ± 13.5 yr compared with 58.6 ± 9.7 yr for patients with no RAS ($p < 0.05$). The level of blood urea nitrogen was significantly higher in patients with RAS, compared to those with no RAS, though the creatinine level was not much different between the two groups. There was also no significant difference between the two groups in other variables including obesity, gender, smoking history, lipid levels and the presence of diabetes. The coronary artery disease tended to be more severe in patients with RAS, compared to those with no RAS (Table 3).

In univariate analysis for predictors of significant RAS, hypertension, peripheral vascular disease, and renal insufficiency defined as a serum creatinine level of 1.5 mg/dL were significantly more common in patients with significant RAS compared to those with

Table 2. Univariate Logistic Regression Analysis of Clinical and Biochemical Variables to Identify Predictors of RAS

| Variables | No RAS (N=382) | RAS (N=45) | χ^2 | p value |
|--------------------------|----------------|-------------|----------|---------|
| PVD | 54 (14.1%) | 21 (46.7%) | 22.73 | 0.0001 |
| Mean age, yr | 58.6±9.7 | 63.2±13.5 | 8.52* | 0.0035 |
| Hypertension | 181 (47.5%) | 35 (77.8%) | 6.02 | 0.0142 |
| BUN, mg/dL | 16.8±8.6 | 20.3±10.1 | 4.59 | 0.0322 |
| Congestive heart failure | 67 (17.5%) | 7 (15.6%) | 0.92 | 0.0397 |
| Female | 127 (33.2%) | 16 (35.5%) | 0.82 | 0.3663 |
| Lp (a), mg/dL | 25.3±23.1 | 29.4±30.3 | 0.67 | 0.4133 |
| Smoking | 197 (51.6%) | 21 (46.7%) | 0.40 | 0.5255 |
| Coronary heart disease | 270 (70.7%) | 32 (71.1%) | 0.38 | 0.8453 |
| Total cholesterol, mg/dL | 190.4±40.9 | 182.4±37.6 | 0.23 | 0.6347 |
| Diabetes mellitus | 83 (21.7%) | 9 (20.0%) | 0.19 | 0.6628 |
| Triglyceride, mg/dL | 149.6±120.7 | 161.2±167.6 | 0.04 | 0.8340 |
| % IBW | 102.5±20.3 | 103.4±14.7 | 0.03 | 0.8719 |

PVD, peripheral vascular disease; BUN, blood urea nitrogen; Lp(a), lipoprotein (a); % IBW, percentile ideal body weight; RAS, renal artery stenosis.

* Analyzed by 10-year increment.

Table 3. Relationship between the Severity of Coronary Artery Disease and RAS

| | No RAS (%) | RAS (%) | χ^2 | p value |
|-------------------|------------|----------|----------|---------|
| No. of vessels | | | | |
| 0 | 120 (29.8) | 5 (20.8) | 0.05 | 0.834 |
| 1 | 126 (31.3) | 8 (32.0) | 0.00 | 0.956 |
| 2 | 70 (17.4) | 4 (16.7) | 0.05 | 0.826 |
| 3 | 87 (21.6) | 9 (37.5) | 2.32 | 0.062 |
| Left main disease | 17 (4.2) | 1 (4.2) | 0.00 | 0.987 |

RAS, renal artery stenosis.

no RAS (Table 4). Multivariate logistic regression of all variables identified three risk factors predictive of RAS (Table 5). The strongest multivariable predictors of RAS were: older age, hypertension, and the presence of peripheral vascular disease ($p=0.0069$, $p=0.0324$, $p=0.0499$, respectively).

DISCUSSION

The incidence of RAS was 10.5% and significant RAS was found in 5.6% of patients. Significant unilateral disease was found in 18 (4.2%) patients and bilateral disease was present in 6 (1.4%). Our finding of a 10.5% prevalence of renal artery narrowing in a cardiac catheterization population is consistent with findings from a previous smaller study which demon-

strated a prevalence of 13.7%.⁶ Ramirez et al.⁶ reported that only 4.9% had more than 50% narrowing of the arterial lumen, which was also similar to our finding. Harding et al.⁴ noted RAS in 30% of patients undergoing cardiac catheterization. Based on these findings, the prevalence of unsuspected RAS seems to be 10 to 30% in a population having cardiac catheterization. As well, during the catheterization, aortograms helped in the evaluation of renal vasculature in this group.

The location of renal artery narrowing has important clinical implications in regard to the approach to potential revascularization. Ostial lesions are generally the results of atherosclerotic aortic plaques that overhang the orifice of the renal artery, producing inflow obstruction.⁷ In our study, the stem of the renal artery was a more common site of stenosis in 62.2% of the patients than the ostium in 37.8%. Among patients who had significant RAS, 20.8% had ostial lesions. The severity of stenosis was not significantly different according to the site of stenosis. On the other hand, Harding et al.⁴ observed that 48% of the lesions were in the ostium and 49% were in the main segment of the renal artery. Based on these findings, renal artery narrowing may occur more frequently in the main segment of the renal artery than in the ostium.

The lesion type was presumed to be atherosclerotic on the basis of its angiographic appearance. This is in contrast to findings from a report of renovascular

Table 4. Univariate Logistic Regression Analysis of Clinical and Biochemical Variables to Identify Predictors of Significant RAS

| Variables | No RAS (N=402) | RAS (N=25) | χ^2 | p value |
|-----------------------------|-------------------|------------------|----------|---------|
| Creatinine ≥ 1.5 mg/dL | 29 (2.2%) | 7 (29.2%) | 20.20 | 0.0001 |
| PVD | 62 (15.4%) | 13 (54.2%) | 17.29 | 0.0001 |
| Hypertension | 198 (24.3%) | 18 (75.0%) | 5.95 | 0.0147 |
| Lp (a), mg/dL | 25.2 \pm 22.8 | 36.3 \pm 41.5 | 2.39 | 0.0665 |
| Diabetes mellitus | 88 (21.8%) | 4 (16.0%) | 1.57 | 0.2095 |
| Coronary heart disease | 283 (70.2%) | 19 (79.2%) | 1.34 | 0.2471 |
| % IBW | 102.9 \pm 20.1 | 97.4 \pm 10.2 | 1.29 | 0.2551 |
| Smoking | 204 (50.6%) | 14 (58.3%) | 0.32 | 0.5733 |
| Female | 137 (34.0%) | 6 (25.0%) | 0.22 | 0.6386 |
| Mean age, yr | 59.2 \pm 9.9 | 58.7 \pm 16.1 | 0.18* | 0.6695 |
| Triglyceride, mg/dL | 151.9 \pm 127.4 | 129.9 \pm 95.8 | 0.05 | 0.8320 |
| Total cholesterol, mg/dL | 190.1 \pm 40.4 | 178.6 \pm 42.2 | 0.00 | 0.9622 |
| Congestive heart failure | 68 (16.9%) | 6 (25.0%) | 0.00 | 0.9869 |

PVD, peripheral vascular disease; Lp (a), lipoprotein (a); % IBW, percentile ideal body weight; RAS, renal artery stenosis.

No RAS: no or insignificant RAS.

* Analyzed by 10-year increment.

Table 5. Multivariate Logistic Regression Analysis of Clinical and Biochemical Values to Identify Predictors of RAS

| Variables | β | p value | Odds ratio |
|-----------------------------|---------|---------|------------|
| Age* | 0.0762 | 0.0069 | 1.08 |
| Hypertension | 0.2739 | 0.0324 | 1.03 |
| Peripheral vascular disease | 0.5862 | 0.0499 | 1.01 |

RAS, renal artery stenosis.

* Analyzed by 10-year increment.

disease where roughly 60% of atherosclerotic lesions and 40% of fibromuscular dysplasia were found in patients investigated for renovascular hypertension.⁸ This discrepancy is likely because of the vastly different populations being studied. Renovascular hypertension was not suspected in this population referred for evaluation of cardiac disease. Olin et al.⁹ and Vetrovec et al.¹⁰ confirmed the overwhelming preponderance of atherosclerotic rather than fibroplastic renovascular disease in similar subsets of patients with advanced age and evidence of multi-organ atherosclerosis, respectively.

Both univariate and multivariate analyses of the clinical and biochemical variables showed that old age, hypertension, and the presence of peripheral vascular disease are the most important markers for increased risk of coexistent angiographic RAS. It is

not surprising that advancing age is the strongest predictor because it is commonly accepted that atherosclerosis in general, as well as atherosclerotic renovascular disease, most often occurs later in life.^{11,12} Schwartz and White¹³ found severe stenosis ($>50\%$ renal artery diameter reduction) in only 5% of autopsy material from patients younger than 64 years. By contrast, 18% of autopsy material from patients 65 to 74 years and 42% from patients over 75 years demonstrated severe RAS.

Almost half of the patient population studied had a history of hypertension, and of the 216 hypertensive patients, 16.2% had RAS, while significant RAS was identified in 8.3%. This finding is similar to another finding of 15.3% incidence of RAS in hypertensive patients who underwent cardiac catheterization.⁶ Renovascular hypertension was not suspected in this population referred for the evaluation of cardiac disease, but based on these findings, it appears that about 15% of hypertensive patients may have atherosclerotic RAS.

The presence of peripheral vascular disease, known as a marker for renal artery disease, was also a good predictor of RAS. In our study, 28% of patients with evidence of peripheral vascular disease had RAS and 54.2% of significant RAS patients had peripheral vascular disease. Patients with atherosclerosis elsewhere, especially with abdominal aortic aneurysms, aorto-occlusive disease, or lower-extremity occlusive

disease, have previously been shown to have a high prevalence of significant RAS.^{9,14,15} This finding supports the belief that atherosclerotic renovascular disease is often a manifestation of generalized atherosclerosis involving the abdominal aorta. Therefore, physical evidence of carotid, femoral, or abdominal bruits seems to raise suspicion of RAS.

Angiotensin-converting enzyme inhibitors have recently been proposed as the treatment of choice in patients with hypertension and peripheral vascular disease since they are free of many of the side effects of other antihypertensive agents. They can, however, induce acute renal failure in patients with RAS.¹⁶ Evidence of RAS probably indicates that those patients are at high risk of renal complications during treatment with angiotensin-converting enzyme inhibitors. We believe that in patients with evidence of peripheral vascular disease, angiotensin-converting enzyme inhibitors should be used with caution. Consideration should be given to the possibility of underlying RAS.

Considering the diffuse nature of atherosclerotic disease, patients with coronary artery disease would also be more likely to have RAS. It has been previously shown that 58% of patients with unsuspected RAS had clinically overt coronary artery disease.¹⁷ Harding et al.⁴ reported that the greater the number of coronary arteries involved, the greater was the likelihood of significant RAS. Our data revealed that 71.1% of patients with RAS had coronary artery disease. However, there was no significant correlation between the severity or the number of coronary artery disease and the presence of RAS. This finding suggests that coronary artery disease is not a strong predictor of unsuspected RAS. Other cardiovascular risk factors such as obesity, diabetes, hyperlipidemia, and congestive heart failure were not correlated with RAS.

By univariate analysis, an elevated creatinine concentration was found to be more common in patients with significant RAS, though it was not a significant predictor of significant RAS by multivariate analysis. The incidence of 29.2% in our study is less than that of Hansen et al.,¹⁸ who found 65% of renal insufficiency (creatinine ≥ 1.3 mg/dL) in patients on surgical management of renovascular disease. Renal function seems to be significantly influenced by the site, type and severity of lesion. In addition, progressive deterioration of renal function in patients

with atherosclerotic RAS is common and occurs even in the presence of blood pressure control.¹⁹ It could be argued that the changes of renal function are evaluated regularly in patients with an elevated creatinine concentration and angiographic evidence of RAS.

In conclusion, the prevalence of angiographically renal artery narrowing, primarily due to atherosclerotic disease, in a population undergoing cardiac catheterization is 10.5%. Old age, hypertension and evidence of peripheral artery occlusive disease represent the important predictors of RAS. Abdominal aortography at the time of cardiac catheterization offers a safe and effective means for evaluating the renal vasculature in this potentially high-risk group.

REFERENCES

1. Textor SC. Atherosclerotic renovascular disease as a cause of end-stage renal disease: cost considerations. *Blood Purif* 1996;14:305-14.
2. Libertine JA, Zinman L, Bresslin DJ, Swinton NW, Legg MA. Renal artery revascularization: restoration of renal function. *JAMA* 1980;244:1340-2.
3. Novick AC, Textor SC, Bodie B, Khauli RB. Revascularization to preserve renal function in patients with atherosclerotic renovascular disease. *Urol Clin North Am* 1984;11:477-90.
4. Harding MB, Smith LR, Himmelstein SI, Harrison K, Phillips HR, Schwab SJ, et al. Renal artery stenosis: Prevalence and associated risk factors in patients undergoing routine cardiac catheterization. *J Am Soc Nephrol* 1992;2:1608-16.
5. Hayes JM, Risius B, Novick AC, Geisinger M, Zelch M, Gifford RW Jr. Experience with percutaneous transluminal angioplasty for renal artery stenosis at the Cleveland Clinic. *J Urol* 1988;139:488-92.
6. Ramirez G, Bugni W, Farber SM, Curry AJ. Incidence of renal artery stenosis in a population having cardiac catheterization. *South Med J* 1987;80:734-7.
7. Jacobson HR. Ischemic renal disease: An overlooked clinical entity? *Kidney Int* 1988;34:729-34.
8. Novick AC. Atherosclerotic renovascular disease. *J Urol* 1981;126:567-72.
9. Olin JW, Melia M, Young JR, Graor RA, Risius B. Prevalence of atherosclerotic renal artery stenosis in patients with atherosclerosis elsewhere. *Am J Med* 1990;88:46N-51N.
10. Vetrovec GW, Landerwehr DM, Edwards VL. Incidence of renal artery stenosis in hypertensive patients undergoing coronary angiography. *J Intervent Cardiol* 1989;2:69-76.
11. Horvath JS, Waugh RC, Tiller DJ, Duggin GG. The

- detection of renovascular hypertension: A study of 490 patients by renal angiography. *QJ Med* 1982;51:139-46.
12. Choudhri AH, Cleland JGF, Rowlands PC, Tran TL, McCarty M, Al-Kutoubi MAO. Unsuspected renal artery stenosis in peripheral vascular disease. *Br Med J* 1991;14:1197-8.
 13. Schwartz CJ, White TA. Stenosis of renal artery: An unselected necropsy study. *Br Med J* 1964;2:1415-21.
 14. Duston HP, Humphries AW, de Wolfe VG, Page IH. Normal arterial pressure in patients with renal arterial stenosis. *JAMA* 1964;187:138-9.
 15. Brewster DC, Retana A, Waltman AC, Darling RC. Angiography in the management of aneurysms of the abdominal aorta: Its value and safety. *N Engl J Med* 1975;292:822-5.
 16. Hricik DE, Browning PJ, Kopelman R, Goorno WE, Madias NE, Dzau VJ. Captopril-induced functional renal insufficiency in patients with bilateral renal artery stenosis in a solitary kidney. *N Engl J Med* 1983;308:373-6.
 17. Valentine RJ, Clagett GP, Miller GL, Myers SI, Martin JD, Chervu A. The coronary risk of unsuspected renal artery stenosis. *J Vasc Surg* 1993;18:433-40.
 18. Hansen KJ, Starr SM, Sands E, Burkart JM, Plonk GW, Dean RH. Contemporary surgical management of renovascular disease. *J Vasc Surg* 1992;16:319-31.
 19. Dean RH, Kieffer RW, Smith BM, Oates JA, Nadeau JHJ, Hollifield JW, et al. Renovascular hypertension: Anatomic and renal function changes during drug therapy. *Arch Surg* 1981;116:1408-15.
-