

## Serological Observation of *Toxoplasma Gondii* Prevalence in *Apodemus Agrarius*, a Dominant Species of Field Rodents in Korea

Soung-Hoo Jeon and Tai-Soon Yong

### Abstract

Field rodents involved in ecological food chains and which are the prey of carnivores in the natural environment may serve as reservoir hosts for *Toxoplasma gondii* infection in humans, however, no data has been published to date in Korea. A total of 1,008 *Apodemus agrarius*, a dominant species of field rodents in Korea, were trapped at various locations around the country, and their serum antibody (IgG) levels to *T. gondii* were examined by ELISA. The mean absorbance was 0.11, and fifteen samples (1.49%) showed positive titers from 0.18 to 0.59. The seropositive samples were analyzed by immunoblot. Five of them showed reactive bands to *T. gondii* water soluble antigens of 30, 35, and 43 kDa. This immunoblot analysis showed very similar patterns to that obtained using sera of experimentally infected mice with *T. gondii*. The present study presents indirect evidence of the existence of *T. gondii* in field rodents in Korea.

**Key Words:** *Toxoplasma gondii*, *Apodemus agrarius*, prevalence, antibody

### INTRODUCTION

*Toxoplasma gondii* is an obligate intracellular parasite that causes significant morbidity and mortality in humans throughout the world. *T. gondii* also infects all kinds of mammals and some birds. The organism has been isolated from herbivorous, carnivorous and omnivorous animals.<sup>1</sup>

In Korea, Soh et al. reported *T. gondii* skin test positive rate in 5.6% of 373 examinees for the first time.<sup>2</sup> And also reported on another seroepidemiological study which indicated that *T. gondii* infection was related with neurological or physical disorders in humans.<sup>3</sup> Choi et al. described two cases of chorioretinitis patients, a six-year-old child with congenital toxoplasmosis and a 26-year-old female with acquired toxoplasmosis.<sup>4</sup> Choi et al. reported two outbreaks of acute toxoplasmosis involving 8 adult patients, who had eaten raw pork.<sup>5</sup>

Dubey et al. conducted a survey on swine and wild

lives in 47 swine farms, to identify sources and reservoirs of *T. gondii* infection, and found that the *T. gondii* antibody positive rate was 2.1–68.3%.<sup>1</sup> The presence of *T. gondii* was confirmed in the brains and hearts of rodents trapped on farms, therefore, the rodents were confirmed as reservoirs of *T. gondii* infection.

Field rodents are herbivores and therefore, probably become infected by ingesting food or water contaminated with oocysts. Since the existence and distribution of *T. gondii* in field rodents in Korea have not been studied until now, the present study was performed to elucidate the seroprevalence rate of *T. gondii* in *Apodemus agrarius*, a dominant species of field rodents in Korea.

### MATERIALS AND METHODS

#### Trapped sites of *Apodemus agrarius*

Thirteen regions and 49 sites in Korea were selected to capture *Apodemus agrarius* during December 1990 and November 1997 (Fig. 1, Table 1). They were trapped at the levee of a rice paddy after harvest and at ridges of ordinary fields using Sherman live traps. Each trap was baited with 20 g of oats-peanut butter ball set up with 2–3 m intervals at 5–6 P.M.

Received December 21, 1999

Accepted May 22, 2000

Department of Parasitology, Yonsei University College of Medicine, Seoul, Korea.

This study was supported by a research grant for research instructors of Yonsei University College of Medicine for 1998.

Address reprint request to Dr. S. H. Jeon, Department of Parasitology, Yonsei University College of Medicine, C.P.O. Box 8044, Seoul 120-752, Korea. Tel: 82-2-361-5290, Fax: 82-2-363-8676, E-mail: shjeon1201@yahoo.co.kr




Fig. 1. Geographic regions, sites (●) and 8 seropositive sites (★) for capture of *A. agrarius* between December 1990 and November 1997 in Korea.

and at 6–7 A.M. the next morning. Live trapped *A. agrarius* were transported to the nearest field stations. After anesthetizing the rodents, blood samples were collected.

#### Preparation of *T. gondii* lysates

The tachyzoite lysate of *T. gondii* (RH strain) was prepared for ELISA and immunoblot analysis. Briefly, the ascitic fluid containing tachyzoites was obtained from ICR mice, which were infected intraperitoneally 4 days earlier with *T. gondii*. Washing tachyzoites several times with 0.01 M phosphate buffered saline (PBS), pH 7.2, by centrifugation at 1,000 rpm. The precipitated host cells were discarded, and suspended tachyzoites, which contained a limited number of contaminated host cells were separated. The tachyzoites were sonicated for 30 sec to prepare the homogenate, which was centrifuged at 4°C, at 15,000 g

using a high-speed centrifuge, and the supernatant was retained. The protein concentration of the lysate was determined using a Bio-Rad protein assay kit (Bio-Rad Labs, Richmond, USA), and stored at -70°C.

#### Serum samples and control sera

Serum samples were collected from *A. agrarius*. The positive control sera were obtained from ICR mice immunized intraperitoneally with *T. gondii* (RH) lysate at 6 weeks post-immunization, and also from mice infected intraperitoneally with 10 to 15 *T. gondii* tissue cysts (ME49 strain, 15–60 μm in diameter) at 3 weeks and 6 weeks postinfection. The negative control sera were obtained from mice injected intraperitoneally with 0.1 M PBS, pH 7.2.

#### ELISA

Anti-*T. gondii* IgG antibody titers in serum samples were measured by ELISA. Optimal conditions for dilutions of reagents were determined by checkerboard titration. Polystyrene plates (Costar, Cambridge, MA, USA) were coated with the lysate (5 μm/ml) diluted in 0.05 M carbonate buffer (pH 9.6), and incubated overnight at 4°C. After rinsing the plates 3 times with washing buffer (0.05% Tween 20 in 0.01 M PBS, pH 7.2), blocking was performed with 1% bovine serum albumin (BSA) in washing buffer for 1 hr at room temperature (RT). Test samples (1 : 200) and the control sera (1 : 200) were then diluted in dilution buffer (0.2% BSA in washing buffer) and incubated for 1 hr at 37°C. The wells were washed, and 1 : 2,500 diluted alkaline phosphatase (AP) conjugated goat anti-mouse IgG antibody (Sigma, St. Louis, MO, USA) in dilution buffer was added and incubated for 1 hr at 37°C. After a final wash, the enzyme substrate (*p*-nitrophenylphosphate, 1 mg/ml in diethanolamine buffer, pH 9.8) was added, and incubated for 10 min at RT. The reaction was stopped by the addition of 1 N NaOH. The absorbance was read at 410 nm using an ELISA Reader (Dynatech MR-5000, Chantilly, VA, USA).

#### SDS-PAGE and immunoblot analysis

SDS-PAGE was performed by the method of Laemmli with 10% acrylamide separating gels.<sup>6</sup> *T. gondii* (RH) tachyzoite extract separated by SDS-

Table 1. The Prevalence Acute of Anti-*Toxoplasma Gondii* Antibodies of *Apodemus Agrarius* Trapped between December 1990 and November 1997 at 49 sites in Korea

Locality	Seropositive/Tested	Prevalence (%) rate	
Seoul:	(1) Kaehwa-dong, Kangso-gu	0/2	0
Inchon:	(2) Oryu-dong, Kyesan-gu	0/10	0
	(3) Yungjong-do	0/6	0
Kyonggi-do:	(4) Tokchong-ri, Yangju-gun	1/19	5.3
	(5) Kwangtan 3-ri, Paju-gun	0/49	0
	(6) Pugok 1-dong, Puchon-shi	0/7	0
	(7) Dorae 5-ri, Koyang-gun	3/112	2.7
	(8) Namsan-ri, Kanghwa-gun	0/22	0
	(9) Kyopo-ri, Pyongtaek-shi	0/43	0
	(10) Sanchong-ri, Pochon-gun	0/12	0
Taejon:	(11) Tandong, Yosong-gu	0/4	0
Chungchongbuk-do:	(12) Shimok-ri, Chong-won-gun	1/40	2.5
	(13) Sosong 2-dong, Chungju-shi	0/4	0
Chungchongnam-do:	(14) Pongan-ri, Kongju-shi	0/10	0
	(15) Toyang-ri, Nonsan-shi	0/13	0
	(16) Toksan-myon, Yesan-gun	0/1	0
	(17) Yongpo-ri, Yon-gi-gun	0/4	0
	(18) Kosan-ri, Tangjin-gun	0/20	0
	(19) Naejang-ri, Hongsong-gun	0/13	0
Kang-won-do:	(20) Kundok-ri, Samchok-gun	0/15	0
	(21) Chongok-dong, Tonghae-shi	3/16	8.8
	(22) Chonchong-ri, Chunchon-shi	0/19	0
	(23) Munhye 1-ri, Chorwon-gun	0/5	0
Chollabuk-do:	(24) Haechon-ri, Wanju-gun	0/9	0
	(25) Shinyong-dong, Iksan-shi	0/91	0
	(26) Changshin-ri, Iksan-shi	0/64	0
	(27) Unsan-ri, Puan-gun	0/2	0
Chollanam-do:	(28) Migok-ri, Hwasun-gun	0/4	0
	(29) Kunso-myon, Yonggwang-gun	0/10	0
	(30) Yong-wol-ri, Hampyong-gun	0/9	0
	(31) Namsan-ri, Changhung-gun	0/10	0
	(32) Chohwa-ri, Yochon-gun	1/21	4.8
Kwangju:	(33) Won-dong, Kwangsan-gu	1/11	9.1
	(34) Uchi-dong, Puk-gu	0/10	0
Kyongsangbuk-do:	(35) Yopae 2-ri, Kimchon-shi	0/19	0
	(36) Chiksan-ri, Yechon-gun	0/22	0
	(37) Chinhyon-dong, Kyongju-shi	0/7	0
	(38) Unsu 2-ri, Kimchon-shi	0/12	0
	(39) Chikha-ri, Andong-shi	0/29	0
	(40) Cho 1-dong, Sangju-shi	0/32	0
	(41) Chinan-ri, Mun-gyong-shi	0/21	0
	(42) Pukyong-ri, Yongdok-gun	0/19	0
Kyongsangnam-do:	(43) Ka-po-dong, Masan-shi	0/24	0
	(44) Toson-ri, Tong-yong-gun	0/15	0
Cheju-do:	(45) Tongkwi-ri, Pukcheju-gun	0/36	0
	(46) Kumdok-ri, Pukcheju-gun	1/42	2.4
	(47) Otung-dong, Cheju-shi	0/2	0
	(48) Ara 1-dong, Cheju-shi	0/31	0
	(49) Kosong-ri, Pukcheju-gun	4/10	40.0
Total (%)	15/1,008	1.49	

PAGE and then electrophoretically transferred to a nitrocellulose (NC) membrane using the method of Towbin et al.<sup>7</sup> Membranes were incubated with each serum sample with positive ELISA titers and sera of controls. NC strips were blocked for 1 hr with 3% BSA in 20 mM PBS, pH 7.4 containing 0.05% Tween 20. After several washes, the strips were incubated with 1 : 100 diluted primary antisera, and then with AP conjugated goat anti-mouse IgG (1 : 1000) antibody. The substrate (NBT/BCIP: Promega, WI, USA) for AP was added and incubated for 5 min at RT. The reaction was stopped by rinsing the strips in water.

**RESULTS**

The results of ELISA for 1,008 serum samples of *A. agrarius* collected at 49 trapped sites are given in Table 1. The mean absorbance of 1,008 serum samples was 0.11 ( $\pm 0.033$ ). The mean absorbance of antibody titers of 22 mice infected with *T. gondii* ME49 tissue cysts at 3 weeks and 6 weeks postinfection, 22 mice immunized with *T. gondii* RH tachyzoite lysate at 6 weeks postimmunization and 22 negative control mice were 0.23 ( $\pm 0.08$ ), 0.57 ( $\pm 0.15$ ), 0.85 ( $\pm 0.17$ ) and 0.08 ( $\pm 0.05$ ), respectively

(Fig. 2). Absorbances greater than the mean plus two standard deviations of the absorbance of seronegatives was considered positive. Fifteen of 1,008 serum samples were found to be antibody positive (0.18–0.59) against *T. gondii* RH strain lysate. Table 1 shows the prevalence rate of anti-*Toxoplasma* antibodies of *A.*




Fig. 2. The geometric mean value ( $\pm$ SD) of ELISA absorbance of the 1,008 sample group (S) was 0.11 ( $\pm 0.033$ ). The mean absorbance of the antibody titers of 22 mice infected with ME49 tissue cysts at 3 weeks (C1) and 6 weeks (C2) postinfection, 22 mice immunized with RH tachyzoite lysate at 6 weeks (C3) postimmunization and 22 negative control mice (C4) were 0.23 ( $\pm 0.08$ ), 0.57 ( $\pm 0.15$ ), 0.85 ( $\pm 0.17$ ) and 0.08 ( $\pm 0.05$ ), respectively.




Fig. 3. Immunoblot analysis of *A. agrarius* sera (1–15) showing reactive bands against *T. gondii* (RH strain) lysate, especially at 30, 34 and 41 kDa. Note a similar reactive band pattern in the lane C1 of ME49 strain of *T. gondii* infected. The band pattern using immune serum with *T. gondii* (RH strain) lysate was different in lane C2. M: Marker, P: *T. gondii* lysate, 1–15: Seropositive serum samples of *A. agrarius*, C1: Mouse serum infected with ME49 tissue cyst, C2: Mouse serum immunized with RH tachyzoite lysate, C3: Seronegative serum sample of *A. agrarius*, C4: Control serum of normal mouse. Arrows of 5 and 6 strips indicate 81, 43, 41, 37, 35, 34, 30 and 28 kDa from top to bottom (C1: 65, 59, 41, 37, 35, 34, 30; C2: 116, 105, 65, 59, 43, 41, 30 kDa).

*agrarius*. Out of 49 trapped sites, seropositive animals were found at 8 sites. The overall positive rate for *T. gondii* was 1.49% (n=15/1,008). The positive rate by locality was highly variable, with the highest being found at Kosong-ri, Cheju-do (40.0%, n=4/10), and the next at Chongok-dong, Kang-won-do (18.8%, n=3/16), whereas the lowest rate was shown at Kumdok-ri, Cheju-do (2.4%, n=1/42).

Of 1,008 samples, 15 positive sera and the extract from RH and ME49 strain were analyzed by immunoblotting (Fig. 3). Of 15 positive samples, 5 showed prominent band patterns, and significantly strong IgG responses to proteins with molecular weights (MWs) of 28, 30, 34, 35, 37, 41, 43 and 81 kDa were found in five sera from Cheju-do (lanes: 5, 6, 7) and Kang-won-do (lanes: 11, 12). The negative control sera and negative *A. agrarius* sera measured by ELISA showed no bands or only a few very faint bands. The ME49 strain infected sera reacted to 30, 34, 35, 37, 41, 59 and 65 kDa proteins. The RH strain immunized sera reacted to 30, 41, 43, 59, 65, 105 and 116 kDa proteins.

## DISCUSSION

From an epidemiological point of view, studies on population densities of field rodents are very important because they are involved in the ecological food chains of mammals in natural environment and may serve as one of the reservoir hosts in the transmission of *T. gondii* to humans. In this study, *A. agrarius*, which is very widely distributed, was found to be the predominant (94.3%, n=1,008/1,068) species of field rodents, followed by *Mus musculus* (1.9%), *Crocidura laisura* (1.8%), *Microtus fortis* (1.1%), *Micromys minutus* (0.4%), *Rattus norvegicus* (0.2%), *Eothenomys rufocanus regulus* (0.1%), *Mustela siberica* (0.1%) and *Rattus rattus* (0.1%). Nineteen *Crocidura laisura*, insectivores, were captured. Twenty *Mus musculus*, two *Rattus norvegicus* and one *Rattus rattus*, which were originally domestic rodents, were collected in the field. *A. agrarius*, a predominant species of field rodent in Korea could serve as an important reservoir host as subordinate of ecological niche of mammals for *T. gondii* transmission to human. In particular, high Anti-*Toxoplasma* IgG level samples were found primarily in animals captured near farmhouses. So herbivores, such as, *A. agrarius* seemed to have been infected by ingesting

farm products or water contaminated with the oocysts in cats' faeces.

A total of 216 animals in captivity at the Seoul Grand Park in Korea were examined by Choi et al., based on antibody titers to *T. gondii* by indirect latex agglutination test.<sup>8</sup> Twenty out of 131 mammals (15.3%) and 2 out of 75 birds (2.7%) showed positive antibody titers, and none among rodents (Indian giant squirrel). With the exception of the exercises mentioned in this paper, the existence and distribution of *T. gondii* have not been investigated to date in the Korean environment. In this study, the authors carried out an epidemiological study on *T. gondii* in nature.

ELISA is one of the most useful serological methods, because a large number of sera can be examined in a relatively short time.<sup>4</sup> Although, the cut-off value in ELISA was determined as mean + 2SD of negative controls in this study, 5 (0.54), 6 (0.44), 7 (0.59), 11 (0.32), and 12 (0.45) absorbance values of the sera of strips with prominent bands on immunoblotting were greater than the mean + 3SD (0.224).

From 1981 to 1990, the prevalence of *T. gondii* infection in small mammals was investigated at various locations in the Czech Republic, and the prevalence rate was reported as 7.4% (n=7/94) in *A. agrarius*.<sup>9</sup> As compared with the results of our study, the prevalence rate in *A. agrarius* in the Czech Republic was much higher. The difference in the seroprevalence rate of *T. gondii* infection in rodents between the Czech Republic and Korea may reflect an actual difference in its prevalence. The difference also might have been caused by differences between the trapping locations or reasons.

Immunoblot analysis showed reactive bands to *T. gondii* water-soluble antigens from 28 to 81 kDa in 5 samples sera (Fig. 3). In particular, the reactive bands at 30, 34, 35, 37 and 41 kDa were from field mouse sera, which were also recognized by the sera of ME49 strain infection. The reactive band of 43 kDa was also recognized in the sera of RH strain immunization. The best characterized of these are the three MWs of 30, 35 and 43 kDa associated with the surface antigens of similar size, described by Couvreur et al.<sup>10</sup> Kasper et al.<sup>11</sup> and Handman et al.<sup>12</sup> The MW of 41 kDa seemed similar to the immunoblot of the 5 sample sera and mouse antisera to ME49 and RH strain. This molecule, the stage-specific protein of tachyzoite, which was associated with

internal organelles, has previously been reported.<sup>13</sup> Tomavo et al. have described 4 bradyzoite surface molecules with MWs of 18, 21, 34 and 36 kDa. The present study also showed reactive bands to proteins with MWs of 34 and 37 kDa, although it is not clear whether these proteins are corresponding bradyzoite surface antigens.<sup>14</sup> The 28 kDa molecule reacts only weakly, and may be identical to cytoplasmic molecules of similar size, as described by Weiss et al.<sup>15</sup> On the other hand, there were no bands, or only few very faint bands with sera of seronegative samples or the serum of the negative control.

The present study presents indirect evidence of the existence of *T. gondii* in field rodents in Korea. Further studies are required to provide definite evidence for the existence of *T. gondii* in field rodents, for an example, by the detection of *T. gondii* by nested polymerase chain reaction technique using species-specific primers or the primary isolation of the parasite.

## ACKNOWLEDGEMENTS

We thank Professor Han-Il Ree, Department of Parasitology, College of Medicine, Yonsei University for identifying field rodents and his critical reading of the manuscript. We also acknowledge Professor Ho-Woo Nam, Department of Parasitology, School of Medicine, Catholic University for kindly providing *Toxoplasma gondii* RH and ME49 strains.

## REFERENCES

- Dubey JP, Weigel RM, Siegel AM, Thulliez P, Kitron UD, Mitchell MA, et al. Sources and reservoirs of *Toxoplasma gondii* infection on 47 swine farms in Illinois. *J Parasitol* 1995;81:723-9.
- Soh CT, Lee SJ, Aha YK. Latent infection by *Toxoplasma gondii* in Korea. *Yonsei Med J* 1960;1:52-4.
- Soh CH, Chung PR, Chung SO, Lew JD. Serological Observation of *Toxoplasma* antibody among neurologically and physically deficient groups in the Seoul area of Korea. *Yonsei Rep Trop Med* 1975;6:23-30.
- Choi JS, Choi CS, Soh CT. Isolation of *Toxoplasma gondii* from congenital and acquired chorioretinitis cases. *Yonsei Rep Trop Med* 1980;11:39-42.
- Choi WY, Nam HW, Kwak NH, Huh W, Kim YR, Kang MW, et al. Foodborne outbreaks of human toxoplasmosis. *J Infect Dis* 1997;175:1280-2.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:681-5.
- Towbin H, Staehelin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci USA* 1979;76:4350-6.
- Choi WY, Yoo JE, Nam HW, Oh CH, Kim SW, Katakura K, et al. *Toxoplasma* antibodies by indirect latex agglutination tests in 200 animals. *Korean J Parasitol* 1987;25:13-23.
- Hejlíček K, Literák I. Long-term study of *Toxoplasma gondii* prevalence in small mammals (Insectivora and Rodentia). *Folia Zool* 1998;47:93-101.
- Couvreur G, Sadak A, Fortier B, Dubremetz JF. Surface antigens of *Toxoplasma gondii*. *Parasitology* 1988;97:1-10.
- Kasper LH, Crabb JH, Pfefferkorn ER. Purification of a major membrane protein of *Toxoplasma gondii* by immunoadsorption with a monoclonal antibody. *J Immunol* 1983; 130:2407-12.
- Handman E, Goding JW, Remington JS. Detection and characterization of membrane antigens of *Toxoplasma gondii*. *J Immunol* 1980;124:2578-83.
- Charif H, Darcy F, Torpier G, Cesbron-Delauw MF, Capron A. *Toxoplasma gondii*: Characterisation and localisation of antigens secreted from tachyzoites. *Exp Parasitol* 1990; 71:114-24.
- Tomavo S, Fortier B, Soete M, Ansel C, Camus D, Dubremetz JF. Characterization of bradyzoite-specific antigens of *Toxoplasma gondii*. *Infect Immun* 1991;59:3750-3.
- Weiss LM, Laplace D, Tanowitz HB, Wittner M. Identification of *Toxoplasma gondii* bradyzoite-specific monoclonal antibodies. *J Infect Dis* 1992;166:213-5.