

Department of Internal, Chonnam National University Medical School, Gwangju, Korea

[illegible]

739

. Heparin glycoprotein IIb/
 IIIa receptor inhibitor taxol coating stent
 stent 가
 가
 가 가 가
 ,
 가
 가
 ent 가
 . 가
 aspirin, ticlopidine, clopidogrel ,
 ,
 . stent
 GP IIb/IIIa
 가
 stent
 가
 가
 가
 가
 가
 가

REFERENCES

- 1) Furman MI, Dauerman HL, Goldberg RJ, Yarzbeski J, Lessard D, Gore JM. *Twenty-two year (1975 to 1997) trends in the incidence, in-hospital and long-term case fatality rates from initial Q-wave and non-Q-wave myocardial infarction: A multi-hospital, community-wide perspective.* *J Am Coll Cardiol* 2001;37:1571-80.
- 2) Zahn R, Schiele R, Schneider S, Gitt AK, Wienbergen H, Seidl K, et al. *for the maximal individual therapy in acute myocardial infarction (MITRA) and the myocardial infarction registry (MIR) study groups. Decreasing hospital mortality between 1994 and 1998 in patients treated with primary angioplasty but not in patients treated with intravenous thrombolysis. Results from the pooled data of the maximal individual therapy in acute myocardial infarction (MITRA) registry and the myocardial infarction registry (MIR).* *J Am Coll Cardiol* 2000;36:2064-71.
- 3) Zahn R, Schiele R, Schneider S, Gitt AK, Wienbergen H, Seidl K, et al. *for the maximal individual therapy in acute myocardial infarction (MITRA) and the myocardial infarction registry (MIR) study groups. Primary angioplasty versus intravenous thrombolysis in acute myocardial infarction: Can we define subgroups of patients benefiting most from primary angioplasty?* *J Am Coll Cardiol* 2001;37:1827-35.
- 4) Every NR, Lehmann KG. *The effectiveness of primary PTCA: Does patient risk matter?* *J Am Coll Cardiol* 2001;37:1836-8.
- 5) The global use of strategies to open occluded coronary arteries in acute coronary syndromes (GUSTO IIb) angioplasty substudy investigators. *A clinical trial comparing primary angioplasty with tissue plasminogen activator for acute myocardial infarction.* *N Engl J Med* 1997;336:1621-28.
- 6) Weaver WD, Simes RJ, Betriu A, Grines CL, Zijlstra F, Garcia E, et al. *Comparison of primary coronary angioplasty and intravenous thrombolytic therapy for acute myocardial infarction.* *JAMA* 1997;278:2093-8.
- 7) Antoniucci D, Santoro GM, Bolognese L, Valenti R, Trapani M, Fazzini PF. *A clinical trial comparing primary stenting of the infarct-related artery with optimal primary angioplasty for acute myocardial infarction: results from the Florence Randomized Elective Stenting in acute coronary occlusions (FRESCO) trial.* *J Am Coll Cardiol* 1998;31:1234-9.
- 8) Suryapranata H, van't Hof AW, Hoorntje JC, deBoer MJ, Zijlstra F. *Randomized comparison of coronary stenting with balloon angioplasty in selected patients with acute myocardial infarction.* *Circulation* 1998;97:2502-05.
- 9) Stone GW, Brodie BR, Griffin JJ, et al. *Clinical and angiographic follow-up after primary stenting in acute myocardial infarction; the primary angioplasty in acute myocardial infarction (PAMI) stent pilot trial.* *Circulation* 1999;99:1548-54.
- 10) Zhu MM, Feit A, Chadow H, Alam M, Kwan T, Clark LT. *Primary stent implantation compared with primary balloon angioplasty for acute myocardial infarction: a meta-analysis of randomized clinical trials.* *Am J Cardiol* 2001;88:297-301.
- 11) Le May MR, Labinaz M, Davies RF, Marquis JF, Laramee LA, O'Brien ER, et al. *Stenting versus thrombolysis in acute myocardial infarction trial (STAT).* *J Am Coll Cardiol* 2001;37:985-91.
- 12) Vogt A, Niederer W, Pfafferott C, Engel H-J, Heinrich KW, Merx W, et al. *On behalf of the study group of the Arbeitsgemeinschaft leitender kardiologischer Krankenhausärzte (ALKK). Direct percutaneous transluminal coronary angioplasty in acute myocardial infarction: predictors of short-term outcome and the impact of coronary stenting.* *Eur Heart J* 1998;19:917-21.