

4

- 2 -

. .

[ ]

:

가

가

가

.

45

가

4

2

K

:

1: 15

가

가

K

2: 17

가

45

가

가

K

:

가

45

,  
4

:

,


:

682-714,

290-3

: (052) 250-7129, Fax: (052) 235-2823


e-mail: twpark@uuh.ulsan.kr



**Fig. 1.** Preoperative both clavicle AP & Lat X-ray show distal clavicle fracture.



**Fig. 2.** Preoperative shoulder axillary X-ray shows posterior displacement of distal clavicle.



**Fig. 3.** On computerized tomographic finding, posterior displacement of distal clavicle is marked. 3D reconstruction image.



**Fig. 4.** Preoperative both clavicle AP & Lat X-ray show distal clavicle fracture.



**Fig. 5.** Preoperative shoulder axillary X-ray shows posterior displacement of distal clavicle.



**Fig. 6.** On computerized tomographic finding, posterior displacement of distal clavicle is marked. 3D reconstruction image.

**Fig. 5.** Preoperative shoulder axillary X-ray shows posterior displacement of distal clavicle.

4

13

## REFERENCES

- 1) **Black GB, McPherson JAM and Reed MH:** Traumatic pseudodislocation of the acromioclavicular joint in children: a fifteen year review. *AM J Sports Med*, 19: 644-646, 1991.
- 2) **David P Richard MD and Andrew Howard MD:** Disatal clavicle fracture mimicking type IV acromioclavicular joint injury in the skeletally immature athlete. *Clin J Sports Med*, 11: 57-59, 2001.
- 3) **Eidman DK, Stiff SJ and Tullous HS:** Acromioclavicular lesions in children. *Am J Sports Med*, 9: 150-154, 1981.
- 4) **F Alan Barber:** Complete Posterior Acromioclavicular Dislocation-A case Report. *Orthopedics*, Vol. 10, No. 3: 493-396, 1987 Mar.
- 5) **Havaraneck P:** Injuries of distal clavicular physis in children. *J Pediatr Orthop*, 9: 213-215, 1989.
- 6) **Lemos MJ:** The evaluation and treatment of the injured acromioclavicular joint in athletes. *Am J Sports Med*, 26: 137-144, 1998.
- 7) **Rogers LF:** The radiology of epiphyseal injuries. *Radiology*, 96: 289-299, 1970.
- 8) **Sanders JO, Rockwood CA and Curtis RJ:** Fractures and dislocations of the humeral shaft and shoulder. IN: Rockwood CA, Wilkins KE, Beaty JH, eds. *Fractures in children*. Philadelphia: J. B. Lippincott, 970-977, 1991.

**Abstract****Distal Clavicle Fracture in Adolescence Mimicking Type IV  
Acromioclavicular Joint Injury****Tae-Woo Park, M.D., Sung-Do Cho, M.D., Chae-Chil Lee, M.D.***Department of Orthopedic Surgery, Ulsan University Hospital,  
University of Ulsan College of Medicine, Ulsan, Korea*

Distal clavicle fracture in children may mimic acromioclavicular joint separation, but have very different prognosis and treatment. We are reporting two cases of distal clavicle fracture in adolescence mimicking type IV acromioclavicular injury. This report demonstrates the importance of shoulder axillary view, computerized tomography as well as physical examination on diagnosis of the distal clavicle fracture in adolescence.

**Key Words:** Distal clavicle fracture, Adolescence

**Address reprint requests to** \_\_\_\_\_

Tae-Woo Park

290-3 Cheonha-Dong, Dong-Ku, Ulsan 682-714 Korea

Department of Orthopedic Surgery, Ulsan University Hospital, University of Ulsan College of Medicine

Tel : +82-52-250-7129, Fax : +82-52-235-2823

E-mail : twpark@uuh.ulsan.kr